



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

LANE MEDICAL LIBRARY STANFORD
G175 .K64 1884
Micro-organisms and disease : an introdu



24503330116

G175
K64
1884

LANE

MEDICAL



LIBRARY

LEVI COOPER LANE FUND

Net 95





MICRO-ORGANISMS AND DISEASE.



MICRO-ORGANISMS AND DISEASE

AN
INTRODUCTION INTO THE STUDY OF
SPECIFIC MICRO-ORGANISMS.

LANE LIBRARY

BY
E. KLEIN, M.D., F.R.S.,
*Joint Lecturer on General Anatomy and Physiology in the Medical School of
St. Bartholomew's Hospital, London.*

WITH 108 ENGRAVINGS.



MACMILLAN AND CO.

1884.

M.

YASUJIMA

London:
R. CLAY, SONS, AND TAYLOR,
BREAD STREET HILL, E.C.

G 110
K 64
1884

TO
JOHN SIMON, C.B., D.C.L., LL.D., F.R.S.,

THIS BOOK

Is Respectfully Dedicated

BY

THE AUTHOR.

P R E F A C E.

THE following book, with few additions, is a reprint of a series of articles that have appeared in the *Practitioner* (1884). Not pretending to be in any sense an exhaustive treatise it will, I hope, nevertheless serve to illustrate most of the important points bearing on investigations into the life history of micro-organisms connected with infectious diseases.

I must not omit to mention that most of the investigations recorded here were carried out for the Medical Department of the Local Government Board during the past ten years.

E. KLEIN.

CONTENTS.

	PAGE
INTRODUCTION	1
CHAPTER I.	
MICROSCOPIC EXAMINATION	3
CHAPTER II.	
PREPARATION OF CULTURE MATERIAL	9
CHAPTER III.	
VESSELS AND INSTRUMENTS USED IN CULTIVATION . . .	17
CHAPTER IV.	
PREPARATION OF CULTURE-MEDIA FOR INOCULATION . . .	21

CONTENTS.

CHAPTER V.	
	PAGE
METHODS OF INOCULATION	25
CHAPTER VI.	
MORPHOLOGY OF BACTERIA	34
CHAPTER VII.	
MICROCOCCUS	37
CHAPTER VIII.	
BACTERIUM	60
CHAPTER IX.	
BACILLUS	66
CHAPTER X.	
BACILLUS: NON-PATHOGENIC FORMS	75
CHAPTER XI.	
BACILLUS: PATHOGENIC FORMS	83

CONTENTS.

xi

CHAPTER XII.

	PAGE
VIBRIO	129

CHAPTER XIII.

SPIROBACTERIUM	131
--------------------------	-----

CHAPTER XIV.

YEAST FUNGI: TORULACEÆ, SACCHAROMYCES	136
---	-----

CHAPTER XV.

MOULD FUNGI: HYPHOMYCETES OR MYCELIAL FUNGI	140
---	-----

CHAPTER XVI.

ACTINOMYCES	148
-----------------------	-----

CHAPTER XVII.

ON RELATIONS OF SEPTIC TO PATHOGENIC ORGANISMS	151
--	-----

CHAPTER XVIII.

VITAL PHENOMENA OF NON-PATHOGENIC ORGANISMS	168
---	-----

CONTENTS.

CHAPTER XIX.

	PAGE
VITAL PHENOMENA OF PATHOGENIC ORGANISMS . . .	177

CHAPTER XX.

VACCINATION AND IMMUNITY	184
------------------------------------	-----

CHAPTER XXI.

ANTISEPTICS	187
-----------------------	-----

MICRO-ORGANISMS AND DISEASE.

INTRODUCTION.

THE relation of micro-organisms to the infectious diseases is admitted to be very intimate; and although it may not be quite so universal as some are inclined to assume, it is nevertheless definitely proved to exist as regards some of the infectious maladies affecting man and brutes. In order to pass in review all the ascertained facts and observations in this vast and constantly-growing field of pathology, and to appreciate and to assign their true value to the many observations bearing on this relation of micro-organisms to disease, it is necessary that the reader, and still more the worker in this field, should be enabled to criticise the observations and facts brought forward by the numerous writers on the subject, for otherwise he would probably take as proved what has really not passed beyond the stage of possibility. And it is this point which requires the most careful attention, viz., to be able to see at a glance that, owing to the imperfect or faulty method of investigation employed, or that, owing to certain inferences incompatible with the general laws and general tendency of the well-founded and experimentally proved facts, the statements set forth in a particular observation or series of observations are not to be accepted.

In all investigations of the relation of micro-organisms to disease it is necessary to bear in mind that, as Koch¹ has pointed out, no observation can be said to be complete, or, one should rather say, in no instance can it be said to have been *satisfactorily proved*, that a particular infectious disease is due

¹ *Die Milzbrand-impfung*, Cassel and Berlin, 1888.

to a particular micro-organism if any one of the following conditions remains unfulfilled :—(1) It is absolutely necessary that the micro-organism in question is present either in the blood or the diseased tissues of man or of an animal suffering or dead from the disease. In this respect great differences exist, for in some infectious diseases the micro-organisms, although present in the diseased tissues, are not present in the blood ; while in others they are present in large numbers in the blood only or in the lymphatics only. These points will be considered hereafter in the special cases. (2) It is necessary to take these micro-organisms from their nidus, from the blood or the tissues as the case may be, to cultivate them artificially in suitable media, *i.e.* outside the animal-body, but by such methods as to exclude the accidental introduction into these media of other micro-organisms ; to go on cultivating them from one cultivation to another for several successive generations, in order to obtain them free of every kind of matter derived from the animal body from which they have been taken in the first instance. (3) After having thus cultivated the micro-organisms for several successive generations it is necessary to re-introduce them into the body of a healthy animal susceptible to the disease, and in this way to show that this animal becomes affected with the same disease as the one from which the organisms were originally derived. (4) And, finally, it is necessary that in this so affected new animal the same micro-organisms should again be found. A particular micro-organism may probably be the cause of a particular disease, but that really and unmistakably it is so can only be inferred with certainty when every one of these desiderata has been satisfied.

It will be my aim in the following pages, first to describe the methods that may be employed with success in investigations bearing on the relation of micro-organisms to disease ; secondly, to describe in conformity with reliable observations the morphology and physiology of the micro-organisms that bear any relation to disease ; and thirdly, to enumerate the observations that have been made in recent years to prove the existence of such an intimate relation. Last, but not least, we shall consider the precise relation of the particular micro-organisms to the causation of disease.

CHAPTER I.

MICROSCOPIC EXAMINATION.

FOR the examination of micro-organisms good high powers are essential, at the least a power magnifying 300 to 400 linear diameters. Zeiss' D or E and Zeiss' or Powell and Lealand's oil immersion 1-12th or 1-16th inch will be found sufficient for all purposes. In the case of tissues stained with aniline dyes a good substage-condenser such as Abbé's or Powell and Lealand's, is invaluable. I use Zeiss' stand with Abbé's condenser, open diaphragm, and plane mirror. As Koch¹ pointed out, and what is now universally acted upon, stained specimens mounted in Canada-balsam solution or Dammar varnish, when examined on an Abbé's condenser, show the micro-organisms with extreme clearness and sharpness.

The examination of the morphological characters of an organism is carried out on fresh unstained, as well as on fresh stained microscopic specimens. Although the latter method is, for reasons hereafter to be mentioned, by far the most perfect and reliable one, it is nevertheless important to ascertain as far as possible the appearances, chemical reactions, and general morphology of perfectly fresh specimens. Blood, juices, tissues, and fluids in which the micro-organisms have been growing, are subjected directly, without any previous preparation, to microscopic examination. With artificial nourishing media in which micro-organisms have been growing, the examination of fresh specimens is of great importance, for the reason that the organisms can be easily identified and their size and general morphological characters be more correctly ascertained than after drying, hardening, and staining. Besides, the chemical reactions can be satisfactorily studied in fresh

¹ *Die Aetiologie d. Wundinfektionskrankheiten.* p. 34, Leipzig, 1879. Translated as *Traumatic Infective Diseases* (New Syd. Soc.), London, 1880.

specimens only. All one has to do is to draw up with a capillary pipette or to take up with the point of a needle a drop or particle of the material, to place it on an object-glass, and to cover it up with a thin cover-glass. Where one has to deal with liquids, such as artificial nourishing fluids, blood, serum, tissue-juices, secretions, transudations, and exudations, no addition is required. In the case of more solid material, such as solid artificial nourishing material, bits of tissue, &c., the addition of a drop of neutral previously well-boiled saline solution (of 0.6 to 0.75 per cent.) is advantageous although not absolutely necessary, since by pressing down the cover-glass a layer of the material sufficiently thin for examination can be obtained. In some instances a bit of tissue can be teased out into fine particles by means of two clean needles. Where it is a question of micro-organisms sufficiently conspicuous by their shape, size, and general appearance, their identification in the fresh condition is not difficult; this is the case with bacilli, actinomyces, and mycelia, but in the case of micrococci, especially when isolated or in couples, and lying in blood, juices, or tissues, their recognition is often extremely difficult. When in large clumps, such as larger or smaller masses of zoogloea, or when in the shape of chains, the identification is not difficult; but in the more isolated state they are not easily recognised owing, as a rule, to the presence of granules or particles of various kinds, from which morphologically their distinction is well-nigh impossible. In such cases there are certain rules of thumb, if I may say so, which assist, although they do not absolutely insure, the diagnosis. These are the micro-chemical reactions. The addition of liquor potassæ leaves micro-organisms quite unaltered, whereas fatty and most albuminous granules alter or altogether disappear by it. Acetic acid from 5 to 10 per cent. strong does not affect micro-organisms, but albuminous and other granules become in most instances altered. These two re-agents, I think, are as reliable as any others; if they fail, then others like alcohol, chloroform, sulphuric ether, &c., are not of any greater help, but the latter re-agents may be used, for instance, when it is a question between fat-granules and micrococci, or crystals and bacilli.

Micro-organisms have a great affinity for certain dyes, especially aniline dyes, and therefore these are used with great success to demonstrate their presence, and to differentiate in many instances morphological details which in the unstained condition are not discernible. The staining is effected on fresh *unaltered* organisms, or after they have been dried. In the

first instance the process is carried out thus :—A microscopic specimen is made, and to it is added afterwards drop after drop of the dye, passing it through the specimen in the usual way of applying fluids to a microscopic specimen, *i.e.* by adding with a capillary pipette the dye at one margin of the cover-glass and sucking it up with a strip of filter-paper applied to the opposite margin of the cover-glass. When the staining has taken place the excess of the dye is washed away with salt solution, water, or alcohol, or both, as the case may be (see below). Unless the organisms are embedded in continuous masses of solids, this method gives good results. In the latter case, say if they are embedded in a microscopic lump of tissue, or in a particular spot of a fine section of a fresh tissue, it is necessary, after having placed the lump or section on an object-glass, to drop the dye on to this previous to putting on the cover-glass. After some minutes the dye is allowed to run off by inclining the object-glass, and then the washing is proceeded with till all the excess of the dye is removed ; the mounting is then done by placing a drop of water or salt solution on the specimen and covering it with a cover-glass. In the case of sections through fresh and hardened tissues containing micro-organisms, the method of staining and of permanently mounting them as a whole is more complicated, and will be detailed presently.

When one has to deal with coherent masses of micro-organisms, present either in natural media (*i.e.* animal tissue), or artificial cultivations, such as zoogloea and pellicles of *micrococcus* or *bacterium*, these can be bodily transferred to a watch-glass, stained, washed, and mounted without much difficulty, either for immediate or permanent use. The permanent specimens are made in this way :—Place the sections or pellicle in a watch-glass containing the dye, leave it there till deeply tinted, take out with a needle or the like, wash in water, then in alcohol, leave here for five minutes or more till most of the excess of the colouring-matter is removed, then lift it on to an object-glass, spread well out, place on it a drop of clove-oil, and after a minute or two drain off the clove-oil, add a drop of Canada-balsam solution (in chloroform or benzol), and cover with a cover-glass. In some special instances such as the bacilli of leprosy and tuberculosis, double staining is required. With other organisms, such as the bacilli of glanders or tuberculosis, the washing is carried out, not with water but with acid (acetic acid and nitric acid respectively). All the details will be stated when dealing with these special organisms,

The method extensively and successively used for the demonstration and preservation of microscopic specimens of micro-organisms in fluids, as blood, pus, and juices, is that of Weigert and Koch, which consists in spreading out on a glass slide or cover-glass a very thin film—the thinner the better—of the fluid (artificial or natural culture medium), blood, pus, or juice, and drying it rapidly by holding it for ten to twenty seconds over the flame of a spirit-lamp or gas-burner. The most successful preparations are obtained when the heating is carried on for such a time that the film, having become opaque at first, rapidly turns transparent. Several drops of the aniline dye to be used are then poured over the specimen, and after remaining on it from five to thirty minutes or more are poured off. The aniline dyes used are those that are commonly known as having a great affinity for cell-nuclei (Hermann); they are also known as the neutral or basic, but not the acid ones. The most useful aniline dyes amongst them are those that are soluble in water; these are preferable to those soluble in alcohol, but in certain special instances (to be mentioned hereafter) some of them are of definite use. Methyl-blue, methyl-violet, vesuvin, Bismarck-brown, magenta, fuchsin, rosanilin, gentian-violet, Spiller's purple, eosin, dahlia, purpurin, iodine-green, are the aniline dyes commonly used. The washing is carried out in all cases, except with tubercle-bacilli and the bacilli of glanders, with distilled water, then with alcohol; this latter as a rule is to wash out the dye from all parts except the micro-organisms, and it is therefore necessary not to carry the washing further than this, but to carry it as far as practicable. After this, wash again with distilled water, dry, and mount in a drop of Canada-balsam solution. Throughout this process it is of course necessary always to remember on which side of the glass the specimen had been dried.

Double staining is carried out with any two of the above dyes; as a rule a brown and blue or violet, or a red and blue, are preferred. Some of the violet and purple dyes have the peculiarity that in some—not in all instances—they give to the preparation a double tint—some things appear blue, others more pink.

The process of double staining is carried out either for each dye separately—i.e. we first apply one dye, after some minutes wash in distilled water, and then apply the second dye—or the two dyes to be used are mixed and then used like a single dye.

In the case of tubercle-bacilli the staining is first done with a magenta mixture (Ehrlich's, Weigert's, or Gibbes' see

below), then washed for a few seconds with a 10 per cent. solution of nitric acid, then for a few minutes with distilled water. After this the preparation is stained with methyl-blue in the ordinary way. Or, after Koch's method, the specimen is first stained in alkaline methyl-blue (mixed with a 10 per cent. solution of caustic potash) for twenty-four hours, or for half an hour to one hour at a temperature of 40° C., and then stained in a concentrated solution of vesuvin. Wash it next with water, then with alcohol, dry, and mount in Canada-balsam solution.

In leprosy, the specimen on the glass is stained with magenta, then washed in distilled water, then stained with methyl-blue, washed, and mounted. With such organisms—as *e.g.* the micrococcus in the sputum of acute croupous pneumonia, and the micrococcus in gonorrhœal discharge—the staining is best carried out with a mixture of methyl-blue and vesuvin.

Weigert's double stain is very excellent for many purposes ; it is prepared thus :—

Saturated watery solution of aniline, 100 ccm.

[This is made thus :—Aniline oil, 1 part, dist. water, 3 parts. Shake every half hour for four hours, and decant the water as the oil settles to the bottom.]

Saturated alcoholic solution of fuchsin, 11 ccm. Mix.

The sections are well stained in this mixture, then washed in distilled water ; after this they are immersed for a few seconds in alcohol, and then transferred for one, two, or three hours to the following solution [Watson Cheyne, *Practitioner*, April 1883, p. 258]:—

Distilled water, 100 ccm.

Saturated alcoholic solution of methyl-blue, 20 ccm.

Formic acid, 10 minima.

After this, wash in alcohol, pass through clove-oil, and mount in Canada-balsam solution.

In examining fresh or hardened tissues for micro-organisms it is necessary to make thin sections, which can be easily done with the aid of any of the microtomes in common use, amongst which Williams's microtome, and especially Dr. Roy's ether-spray freezing microtome, are no doubt the best and easiest to manipulate. As regards hardened material, it is necessary to remember that the hardening must be carried

out properly, small bits—about a half to one cubic inch—of tissue being placed in alcohol, or better in Müller's fluid, and kept there; in the first instance, for two to five days; in the second, for from one to three weeks or more. Then small bits are cut out, of which it is desired to make sections. Those hardened in spirit must be soaked well in water to enable the material to freeze, then superficially dried with blotting-paper, and then used for cutting sections with Roy's microtome. Those hardened in Müller's fluid are at once superficially dried with blotting-paper and cut. When making sections with Williams's freezing microtome it is necessary to soak the material first in gum mucilage and then to freeze and to cut. Fresh tissues are at once cut with the freezing microtome, the sections placed in 0.6 per cent. of saline solution, floated out and well spread out, and then stained by transferring them in this condition, *i.e.* well spread out, into a watch-glass containing the dye. The sections of hardened tissues are floated out in water, well spread out, and then transferred to the dye or dyes as the case may be.

It is necessary to prevent too much shrinking of the sections, especially those of fresh tissues; for this reason it is advisable to float the sections in the saline solution or water, as the case may be, on a broad lifter or spatula, to spread them well out upon it, and to transfer them carefully into the dye, then into the dish with water used for washing off the excess of the dye, to transfer them, well spread out on the lifter, to alcohol, then after several minutes to oil of cloves, and finally on to a glass slide, on which they are mounted in the usual manner with Canada-balsam solution, the excess of clove-oil being previously drained off.

It is advisable, although not absolutely essential, to keep the sections in a well spread-out condition for a few seconds in alcohol before placing them into the dye.

CHAPTER II.

PREPARATION OF CULTURE MATERIAL.

ARTIFICIAL cultivations of micro-organisms in suitable nourishing media in the incubator (Fig. 1) at temperatures




FIG. 1.—INCUBATOR, WITH PAGE'S REGULATOR.

A. Page's Regulator.—This consists of a tube filled with mercury, and immersed in the water surrounding the chamber of the incubator. In the upper part of the tube, above the mercurial column, is a fine open glass tube, having

near the lower end a fine hole. When the temperature of the water rises, the mercurial column rises, and at a certain temperature rises above the lower open end of the small inner glass tube just mentioned. If this point is reached, then the burner at C receives only the amount of gas that passes through the fine lateral hole of that inner glass tube. If the temperature of the water falls, the mercury falls, and the lower end of the inner glass tube becomes again free, and now the burner at C receives a much greater supply of gas. If so, the temperature of the water again rises, the mercury rises, obstructs the lower end of the inner glass tube, the supply of gas is reduced to what can pass through the fine lateral hole, and consequently the temperature again falls, and so on. To adjust the regulator it is necessary when the thermometer indicates the required degree of temperature to push the outer large glass tube, and with it the inner tube, of the regulator so far down that the top of the mercurial column just obstructs the free end of the inner glass. The temperature then regulates itself for the reasons stated previously. These regulators are sufficient for all practical purposes when it is not a question of small differences in temperature, since they are tolerably constant within one or two centigrades. The trouble one experiences in the working of these and other similar regulators arises from the inconstancy of the main gas supply, this, as is well known, varying within wide limits. The stopcock, E, obviates this to a limited extent, when this is put at an angle of 45° only a limited amount of gas passes from the main supply tube to the regulator, and therefore the variations in pressure of the gas are not felt to their full extent. A Sugg's regulator interposed between E and the main supply tap is very useful.

B. Thermometer to indicate the temperature in the chamber.

C. Gas burner

D. Chamber of incubator.

E. Stopcock to regulate when required the supply of gas.

F. Main supply.—The upper, lower, right and left walls of the incubator are made of a double layer of tin; between the two is water. The front and back of the chamber are closed by a movable glass plate.

An excellent incubator for constant temperature is made by the Cambridge Scientific Instrument Company. It has a double gas supply: one small permanent flame, and a second one subject to the regulator.

varying between 30° and 40° C., are necessary in order to study more accurately the life-history of the septic as well as the pathogenic organisms. Moreover, large numbers of them become available in a short time, and their relation to disease can be tested more conveniently. For if it should be found that, having carried on outside the animal body successive cultivations of a particular organism, the re-introduction of this cultivated organism into the animal body is again productive of the same disorder as before, then the conclusion becomes inevitable that this organism is intimately related to the causation of the disease. It must be conceded that after several successive cultivations in fluids any hypothetical substance supposed to be the *materies morbi*, and introduced at first from the blood or tissues, being in a very diluted condition in the first cultivation, would after several cultivations be practically lost. But if this last cultivation should be found to act in the same manner pathogenically, *i.e.* if every droplet of it, charged with the new brood of the

organism, nevertheless possesses full pathogenic power, then it is logical to say that this pathogenic property rests with the organism. For this and other reasons it is of essential importance to be able to carry on successive cultivations of one and the same organism without any accidental contamination or admixture, *i.e.* it is necessary to carry on *pure cultivations*.

ARTIFICIAL CULTIVATION MEDIA.

A.—FLUIDS.

As fluid nourishing material the following are used with preference :—

1. *Broth made from meat—pork, beef, rabbit, chicken.*—The connective tissue and fat are first cut out from the fresh meat—in the case of rabbit or chicken the whole animal without head or viscera is used—and then placed in water and boiled. Generally for each pound half an hour's good boiling is allowed. With regard to the quantity of water, each pound of meat ought to yield ultimately at least one pint of broth. When boiled, the broth is allowed to stand, the fat is skimmed off, and the broth well neutralised by adding liquor potassæ or, better still, carbonate of sodium.

The fresher the meat the less acid (sarcolactic acid) is in the broth before neutralisation. The broth is then filtered through a filter, previously overheated (see below), into flasks previously sterilised (see below). If the broth is not clear after once filtering it is filtered again. If not clear then, it is allowed to stand for several hours. A fine sediment is found at the bottom of the vessel, and from this the clear supernatant fluid is decanted into a sterilised vessel. The broth, if not clear after the first filtering, can be cleared by boiling it with the white of an egg. The now clear fluid is filtered again. The flasks which receive the broth are well plugged with sterilised cotton-wool (see below). In this state the flask is placed over a Bunsen burner (Fig. 2) on a wire netting and boiled for half an hour or more ; during the boiling the cotton-wool plug is lifted out for half its length. The flask ought not to contain more broth than about one-half or two-thirds of its volume, to prevent the broth from rising too much and wetting the plug. When turning off the flame the plug is pushed down so as fully to plug the neck and mouth of the flask ; a beaker with sterile cotton-wool cap is placed over

the mouth of the flask (Fig. 3), and this is allowed to stand for one night. Next day the boiling is repeated for half an hour or more in the same manner as before. If the meat has been fresh and the vessels and cotton-wool have been sterile, twice boiling is found sufficient to destroy every impurity. But to make sure, the broth is placed in the incubator and kept there for twenty-four hours at a temperature of 32° – 38° C., and then boiled on the next day for half an hour in the usual way. The supposition is made, that if by any chance after twice boiling the broth it should contain unchanged spores of bacilli—the only organisms that will resist boiling, although




FIG. 2.—A BUNSEN BURNER, WITH ROSE FOR BOILING FLUIDS IN TEST TUBES.



FIG. 3.—A FLASK CONTAINING STERILE STOCK FLUID.

they do not resist boiling for more than half an hour—the spores would germinate into bacilli when kept for twenty-four hours in the incubator at 32° – 38° , and these would then be killed by the third boiling. As a matter of fact I have not as a rule found any contaminating germs survive the second boiling. It is of course to be borne in mind that during the first as well as second and subsequent boiling the cotton-wool plug is not removed from the mouth of the flask, but is only raised out half its length from the neck. The cotton-wool and the cotton-wool cap and beaker are replaced immediately or

simultaneously with the turning off of the burner. This broth so prepared is placed in the incubator at 32°—38° C. and kept there from one to three weeks. If, as is generally the case, it remains limpid, it is considered completely sterile.

2. *Peptone and Sugar Solution*.—Beef peptone (Savory and Moore's) is dissolved in distilled water, over a burner, to the amount of about 2 per cent. ; to the solution is added cane sugar to the amount of about 1 per cent. ; so that every 100 ccm. of the fluid contains two grammes of peptone and one gramme of sugar. When dissolved it is well neutralised and then filtered (the vessels being of course also in this, as in all other cases, sterilised by overheating) into flasks, and treated in the same manner as the broth.

The same fluid can be used without the addition of the cane-sugar ; or peptone dissolved to the amount of 1 to 2 per cent. in broth.

3. *Buchner's Fluid*.—10 parts of Liebig's extract, and 8 parts of peptone, in 1,000 parts of water.

4. *Hydrocele Fluid* (Koch).—A new or well sterilised (by over-heating) trocar and cannula are used for the tapping ; to the cannula is fixed an india-rubber tube that has been soaking in strong carbolic acid solution for forty-eight hours. The distal end of the tube is introduced carefully and rapidly into the neck of a sterilised flask plugged with sterile cotton-wool, and the fluid thus allowed to flow into the flask to about two-thirds of its volume. This is then exposed in a water- or sand-bath to a temperature of from 58° to 62° C. for three to five hours on five or six consecutive days. Placed then into the incubator at 32°—38° C. for from one to three weeks, the fluid remains limpid.

5. *Blood Serum* (Koch).—A glass cannula and india-rubber tubing are soaked for forty-eight hours in strong carbolic acid ; the cannula is tied into the carotid artery of a healthy sheep, and the arterial blood, after opening the clip at the proximal end of the artery, is allowed to flow into a sterile flask, the distal end having been introduced into the neck of the flask as above. After letting it stand for twelve to twenty-four hours the clear serum is taken off by means of a large sterilised glass pipette or glass syphon ; this is carefully introduced between the cotton-wool plug and glass neck, and then discharged into a sterile plugged flask or large test-tube, the pipette or syphon being also here carefully introduced between cotton-wool plug and glass.

This stock serum is then heated for successive days in the same manner as the hydrocele fluid.

Of less common use are :

6. *Pasteur's Fluid*.—In 100 parts of distilled water are dissolved 10 parts of pure cane-sugar, 1 part of ammonium tartrate, and the ash of 1 part of yeast.

7. *Cohn's Fluid*.—100 ccm. of distilled water, 1 gramme of ammonium tartrate, no sugar, and instead of the ash of yeast are substituted (A. Mayer) 0·5 gramme of potassium phosphate, 0·5 gramme of crystallised magnesium sulphate, 0·05 gramme of (tribasic) calcium phosphate. These two fluids are sterilised in the same manner as the broth and peptone solutions. Pathogenic organisms do not thrive in either of these two fluids.

B.—SOLIDS.

The solid media have the great advantage over the fluids that in the former artificial cultures can be carried out more easily ; as, owing to the resistance the solid basis offers to the growth of the organisms, they remain more limited to the spot or spots on which they are sown, and therefore can be watched more easily ; besides, an accidental contamination, *i.e.* a growth appearing at a spot at which no sowing was made, can be recognised at once. These advantages are perhaps of the greatest use when it is intended to grow the organisms on a surface exposed to the influence of air—of course protected from contamination with other organisms.

These advantages of solid media have been very minutely pointed out by Koch in his researches on pathogenic bacteria.¹

As solid media are used :

1. *Slices of Boiled Potato or Boiled White of Egg or Paste* (Fokker, Schröter, Cohn, Wernich).—Although these are of great use in the study of hyphomycetes, and especially of pigment-bacteria, they are not generally used in the case of other bacteria and pathogenic organisms. The progress of the growth of a particular organism sown out at a particular spot or line on the surface of these substances can be easily watched with the unaided eye. These substances when quite fresh are placed on flat glass dishes ; these are covered with a well-fitting bell-glass, the space being kept moist by a piece of moist blotting-paper placed in the dish.

2. *Gelatine* (Brefeld, Grawitz, Koch).—This is used advantageously as a mixture with broth, peptone, beef-extract, blood-serum, or hydrocele fluid. Koch, who introduced this mixture, used it for the cultivation of bacteria on solids, to be exposed

¹ *Mittheilungen d. k. Gesundheitsamtes*, 1. 1881.

to the air ; the proportion of gelatine in the mixture was 2 to 3 per cent. But this mixture, although solid at ordinary temperature, does not keep solid in the incubator, not even at 20° C. I have found that at least 7.5 per cent. of gelatine must be contained in the mixture to keep it solid at 20° to 25° C. Above this temperature not even 11 per cent. gelatine will keep solid.

The finest (gold label) gelatine, in thin tablets, is cut up in small strips ; these are soaked in distilled water (1 in 6) overnight, and then dissolved over a water-bath, well neutralised with carbonate of sodium, and filtered hot. If not clear, it is boiled with white of egg, and passed hot through sterilised fine calico. Then this fluid is mixed with half its bulk of broth, peptone solution, or beef-extract solution, so that there is 1 part of gelatine in 9 parts of fluid, or 11½ per cent. of gelatine. This mixture is boiled repeatedly and treated like broth, as described above. The mixture can, when cast solid, be liquefied by melting it on the water-bath, can be easily decanted into sterilised plugged test-tubes (see below), and can then be used as a good solid nourishing material for the cultivation of organisms up to 25° C.

The above gelatine solution without admixture can be boiled once or twice, and thus made sterile and kept as a stock. This can be used as an addition to blood-serum or hydrocele fluid ; the mixture must be sterilised in the same way as serum or hydrocele fluid alone, *i.e.* exposed for five to seven days to a temperature of 58° to 62° C. Of course, whatever the proportions are in which the two are mixed, the mixture does not keep solid above 25° C. But by exposing it for from several days to several weeks to the heat of the incubator, the mixture can by evaporation be rendered practically solid for higher temperatures also.

3. More satisfactory, because capable of remaining solid at any temperature, is solid serum of blood, solid hydrocele fluid (G. Makins), and Agar-Agar (Koch).

The first, *i.e.* the serum of blood, and the second, *i.e.* the hydrocele fluid, can be made solid by heating the above sterile serum or hydrocele fluid respectively (see page 13) *gradually* up to 68°—70° C. In the course of an hour or two the material becomes solid, losing slightly its limpidity, but is sufficiently transparent for all practical purposes. By heating it rapidly, or heating it above 70°, it becomes solid, granular, and opaque. Of course, once thus made solid it cannot be liquefied again, and therefore must be already contained in the vessels (test-tubes and small flasks) in which

the growth of organisms is to be carried on. Or blood serum and hydrocele fluid can be rendered solid by exposing the sterilised material (see above), in sterile plugged test-tubes, to a moderate heat—*e.g.* in the incubator at 32° to 38° C.—for several weeks. Through evaporation the material is rendered solid. Thus treated it retains its limpidity in a perfect manner.

Agar-Agar, or Japan isinglass, is very difficult to obtain,¹ it is sold in the shape of very thin, shrivelled, transparent lamellæ, or narrow bands. It is soaked overnight in salt water (one part of Agar-Agar to five or six of salt water), and then dissolved on the water-bath; well neutralised with carbonate of sodium, filtered and mixed with a third of its bulk of broth, peptone, or beef-extract solution. I use now as a rule peptone solution, as described above. Well boiled on two or three successive days, each time for thirty minutes to an hour, in sterile flasks, a sterile material is obtained, which is quite transparent, and remains solid up to a temperature of 45°–50° C., *i.e.* a temperature much higher than is ever used for the cultivation of micro-organisms. It becomes liquid at higher temperatures, and in case of necessity can be again subjected to boiling. Before considering it as perfectly sterile it ought to be kept like all other materials for from several days to several weeks in the incubator at 32°–38° C. If quite limpid after this time it may safely be considered as sterile.

Amongst all the solid media, I have found this mixture of Agar-Agar and peptone sugar solution to be the best in many respects. It is beautifully limpid and solid, and an excellent nourishing material. Agar-Agar alone without the admixture of peptone is not satisfactory as a culture medium.

¹ Messrs. Christy and Co., of 155 Fenchurch Street, have succeeded in obtaining for me large quantities of this material from Paris. I understand from my friend Dr. R. Maddox that this substance is in reality what the French call *Gelose*.

CHAPTER III.

VESSELS AND INSTRUMENTS USED IN CULTIVATIONS.

All vessels (flasks, test-tubes, beakers, filters, calico), to be used are first thoroughly sterilised by overheating. In the case of flasks and test-tubes, this can be done by exposing them thoroughly in *all parts* to the open flame of a large Fletcher's burner; *while thoroughly heated* the mouth is plugged with a good long plug (1 to 2 inches) of sterile cotton-wool, this being pushed in by means of overheated forceps. The plug in all cases must not be loose, but also not too firm—an error in the latter direction being of course preferable to one in the former. The cotton-wool plug may, if long enough, be single; or, if short ones are used, double. Or the flasks and test-tubes are placed in an air-chamber (see Fig. 4) heated by a large Fletcher's burner for several hours, up to between 130° and 150° C. In the case of small flasks and test-tubes this process is of course much more convenient, since a large number can be heated simultaneously. Beakers and glass filters to be used merely for a temporary operation are placed over a wire net on a tripod and heated by the flame of a Bunsen's burner. In the case of test-tubes which are to receive cultivation-fluids, I generally expose them, after having been cleaned and dried, in the air-chamber for several hours (three to six) to a temperature of from 130°—150° C.: while hot they are taken out *seriatim*, plugged with the sterile cotton-wool, and replaced in the air-chamber, and heated again for several hours. All this, and other operations to be described below, may appear to some rather tedious and unnecessarily complicated, but it cannot be too strongly insisted on that in these matters one cannot be too scrupulous. A slight relaxation may, and occasionally is.

followed by disastrous consequences in the shape of accidental contamination, and consequent loss of material prepared at the cost of much labour and time. Long experience in these matters has taught me that, although in some instances less scrupulous care has not been followed by bad results, still I have had also many unpleasant failures owing to slight laxity in these matters.




FIG. 4.—HOT-AIR CHAMBER FOR STERILISING TEST-TUBES AND COTTON-WOOL.

An iron chamber with double wall, the inner chamber having separate folding doors. In the inner chamber are placed the test-tubes, glasses, &c., and the cotton-wool, the latter in a loose condition. Both sets of doors are closed, and the apparatus heated by a large Fletcher's burner. A thermometer passing from the inner chamber through the upper wall indicates the temperature of the chamber.

Several weeks' work may be annihilated by a single omission. Sometimes one is perhaps in a slight hurry, and does not think the want of an additional heating of the test-tube or cotton-wool or an additional boiling of the fluid will be followed by any bad consequences. But, alas, nature does not take into account our convenience, and failure is our reward. If in any kind of experiments "overdoing" is an error in the right direction, it is in these very experiments in *the cultivation of micro-organisms*.

The *cotton-wool* used for plugging flasks and test-tubes is prepared by pulling up loosely a quantity of good cotton-wool and exposing it in a loose state in the air-chamber to a temperature of 130°—150° C. for several hours, for several successive days. The cotton-wool ought to be just brown, *i.e.* just singed. Too much charring makes it very brittle, and it is then difficult to make of it a satisfactory plug. The plug used should not be too firm and not too loose: in the former case it is not easy to lift it up quickly, and in the latter it does not close sufficiently well. Cotton-wool that has been kept, say only for a day or two in the air-chamber for three or four hours is not absolutely sterile; nor is cotton-wool that has been kept in a compressed state in the air-chamber for any number of days. The central portions remain under these conditions quite white and are not sterile. No cotton-wool that is not just brown, *i.e.* just singed, is safe from risk of impurity. No cotton-wool steeped in absolute alcohol, strong carbolic acid, or any other disinfecting fluid, for ever so many days or weeks, can be absolutely relied on.

As stated above, a plug of sterile cotton-wool tolerably firm, of about one to two inches, or two plugs of about one inch each, are used for the plugging of the flasks and test-tubes. An assertion such as that made by Dr. Williams at the British Association (Biological Section, September 1883), that cotton-wool plugs are not reliable, because they do not protect the fluids in the vessels plugged with them from accidental air-contamination, is to be accepted only as applying to very loose plugs and to cotton-wool not properly sterilised. To good firm plugs of sterile cotton-wool it evidently cannot apply, since all the results of all workers in this field (Pasteur, Sanderson, Cohn, Koch, Klebs, Buchner, and many others) are against it.

Instruments, such as the points of needles, and forceps, used in the processes of cultivation, lifting up cotton-wool plugs, making cotton-wool plugs, inoculations, &c., must be heated in the open flame of a Bunsen burner, if they are to be absolutely relied on for cleanliness. Scissors and knives used



FIG. 5.—A CAPILLARY PIPETTE, DRAWN OUT INTO FINE POINTS: LENGTH ABOUT 12 TO 14 INCHES.

for cutting tissues which are intended for inoculation, ought to be likewise scrupulously clean. One ought to keep a special set of instruments, the blades of which are capable of being heated in the open flame without being spoilt.

Syringes used for cutaneous, subcutaneous, or other inoculations, ought to be capable of being overheated. The ordinary Pravaz syringe of vulcanite not being capable of undergoing this process, Koch has devised a glass syringe similar to the Pravaz syringe. I do not use any syringe for inoculation, but prefer using each time a fresh *capillary glass pipette* made just before the inoculation. Into this pipette I draw the droplet to be used for inoculation, and having made a very small incision—about $\frac{1}{8}$ of an inch—through the skin, the pointed end of the pipette is pushed forward into the subcutaneous tissue for about half an inch or one inch and then the fluid is blown out into the tissue. In this way I am always absolutely safe from any contamination with a previously used virus, which might possibly adhere to one or other part of a syringe.

The fine point of capillary pipettes (Fig. 5), used for inoculation of animals, or for drawing out a drop of fluid of a cultivation in a flask or test-tube, or for inoculating material contained in a test-tube or flask, are thus made: while one hand holds the bulb of the pipette, the other holds one end, and putting at some distance from this end the tube into an ordinary flame and quickly drawing it out, a point of extreme fineness can be made. The same is done with the other end. Such a pipette can be considered as practically closed at both ends.

CHAPTER IV.

PREPARATION OF CULTURE-MEDIA FOR INOCULATION.

WE have on a former page described the methods to obtain sterile stock of nourishing media suitable for artificial cultivations. The solids, as serum gelatine, serum, and hydrocele fluid, must, before solidification, be placed in test-tubes and small flasks, and then sterilised in the manner above described, to be made ready for establishing cultures, *i.e.* for inoculation. The Agar-Agar mixture however can, like broth, peptone mixture, beef extract solution, and gelatine mixtures, be kept as stock in large flasks. When thus sterile, these latter can be decanted into a number of test-tubes or small flasks, in which the cultivation is to be carried out. Gelatine mixtures (gelatine and broth, gelatine and peptone, gelatine and beef extract) and the Agar-Agar mixture, must of course be liquefied over a flame before being ready for decanting. The test-tubes most suitable are about six inches long, and should not be less than about one inch broad; the flasks are about of the capacity of one to two ounces, and ought to have a neck of comparatively good width. The test-tubes receive the fluids for about one and a half to two and a half inches in depth, the flasks for about one-fourth to one-third of their bulk. All these test-tubes and flasks with their cotton-wool plugs, before receiving the material, should be thoroughly sterilised by overheating. As I mentioned in the previous chapter, this ought to be well borne in mind, for starting with a sterile nourishing fluid—*i.e.* one that has been kept in the stock flask for several days to several weeks in the incubator at a temperature of from 32°—38° C. and that has remained perfectly clear and limpid—and working with thoroughly sterilised test-tubes and cotton-wool plugs—very

little care is required to obtain sterile material ready for inoculation. To start with a stock of nourishing material, however well sterilised, and to decant it into test-tubes with cotton-wool plugs not absolutely sterile must lead to failure. I have seen this happen over and over again, and all the material decanted became consequently contaminated and thereby useless for inoculations. The test-tubes and flasks must be well cleaned, then dried, placed in the air-chamber, and kept there exposed for several hours to a temperature of from 130°—150° C. on several successive days, or they may be thoroughly heated in all parts over the open flame of a gas-burner. The same applies to the cotton-wool, as mentioned in a former chapter. The test-tubes and flasks are plugged by means of clean forceps with the cotton-wool which is just brown, and then replaced in the air-chamber and again heated for several hours on two or three occasions up to a temperature of 130°—150° C., or they may be well heated over the open flame of the burner. To decant sterile stock fluid into these test-tubes and flasks, I proceed thus: A clean beaker with spout, covered with a clean glass plate, is placed on a wire net on a tripod over the flame of a Bunsen burner, and thoroughly heated for half an hour or so; then it is allowed to cool, and when cool, the plug of the stock flask is lifted with forceps, and some of the sterile fluid quickly poured from the flask into the beaker. The plug is replaced in the neck of the stock flask and the beaker covered with the glass plate. Of course the quantity poured into the beaker should be large enough to supply the required number of test-tubes or small flasks. The stock flask containing still some fluid, having been opened for however short a time, has of course been exposed to air-contamination, and therefore must be treated accordingly, if the fluid left in it is to serve as sterile nourishing material on a future occasion. Consequently it is subjected to boiling for from fifteen to thirty minutes, then placed in the incubator, boiled again the next day and put back in the incubator, where it is left at a temperature of from 32°—38° C. for several days. If after a week or so the fluid remains limpid, it is of course to be considered sterile.

Next, the fluid that has been poured into the beaker (covered with the glass plate) is poured as quickly as possible into the test-tubes, one after the other, by lifting with clean forceps the plug and pouring in the fluid to a depth of one and a half to two and a half inches, and the plug replaced.

During this procedure, contamination with air-organisms, if

there be any about, becomes inevitable. To lessen this chance as much as possible, it is necessary to lift the plug with clean forceps, to pour the fluid as rapidly as is practicable into the test-tube or flask, and to replace immediately the cotton-wool plug. Further it is necessary to bear in mind, that the atmosphere is not at all times and everywhere equally contaminated (see Prof. Tyndall's observations). I generally avoid undertaking this process on windy days, and when I do it, I generally close windows and doors and keep the air in the room as still as possible. I do not do it in a room in which




FIG. 6.—A BEAKER CONTAINING A NUMBER OF CULTURE-TUBES PLUGGED WITH COTTON-WOOL.

recently (say an hour or two previously) the floor, walls, or tables have been swept.

I have opened under these conditions the plugs of test-tubes containing sterile material, and kept them so for a time varying from one to ten seconds, and in some instances I have not seen more than from 10 to 20 per cent. contaminated.

Now, having filled the required number of test-tubes and flasks with the desired quantity of fluid, I subject these *seriatim* to boiling. By means of an ordinary test-tube

holder I hold them above a very small flame until the fluid boils, and keep it so boiling for from two to five minutes. During this process of boiling the cotton-wool is only slightly pulled up, and immediately before ceasing to boil the plug is again replaced, and pushed down with a clean glass rod. Then the test-tube is placed (of course upright) in a beaker at the bottom of which a layer of cotton-wool—a sort of cushion—has been placed. When finished, the test-tubes in the beaker are all transferred to the incubator and kept there for from twelve to twenty-four hours at a temperature of 32°—38° C. Then the boiling is repeated once more. After this they are kept in the incubator for several days to several weeks. I generally keep them there for two to three weeks, and all those in which the fluid has remained limpid and clear are considered sterile and ready for use. As a rule, starting with sterile stock fluid, and using thoroughly sterile test-tubes and cotton-wool plugs, after once or twice boiling after decanting, there ought to be no loss of tubes through accidental contamination with air-organisms (during decanting). Sometimes, however, I have had loss to the amount of 5 per cent. or more, but then there was always a hitch of some kind traceable. To decant under carbolic acid spray is not practicable and possesses many unpleasant drawbacks, besides, in some instances when I used it, there was really a greater percentage of contaminated tubes than without it. I therefore do not use the spray.

Test-tubes containing solid nourishing material are generally kept sufficiently inclined during solidification of the material (see a former chapter) to allow the material to spread into a layer of large area, although this is not essential.

CHAPTER V.

METHODS OF INOCULATION.

HAVING now in test-tubes and small flasks sterile material ready for inoculation, it is necessary to describe the mode of inoculating the same.

1. *Inoculations from Artificial Cultures.*—The first and simplest is the case where it is required to inoculate a new tube or flask with a definite organism that has been growing previously in a culture-tube; that is to say, where it is required to establish from an artificial cultivation a new and further artificial cultivation. Take a freshly drawn-out capillary pipette, with a fine point, as described in a former chapter; draw up with clean forceps slightly the top part of the cotton-wool plug of the old tube or flask, push carefully and gently one of the pointed ends of the capillary pipette—the other can be broken off blunt—through the remaining part of the cotton-wool plug, and push it downwards till it emerges into the culture-fluid, or, if this be solid material, till it reaches the spot or place where the organism is growing; allow a small droplet to ascend into the capillary pipette, which it readily does by capillarity; or if a larger quantity is required draw it up by gently sucking at the outer end of the capillary pipette. Then draw the capillary pipette altogether out of the tube and cotton-wool plug, and push this latter down with the forceps into its former position. Immediately after this proceed to inoculate the new culture-tube by doing exactly the same as before, viz., draw up slightly with the forceps the top part of its cotton-wool plug, push through the remainder of this plug the pointed end of the capillary pipette, *i. e.* the one containing the droplet of the material to be sown, and push it into the material at the bottom of the test-tube or flask. A trace of the sowing material flows out by itself, or, if a large

quantity is required, it is carefully blown from the pipette. If the sowing is to be carried out on the surface of the solid nourishing material, the inoculation is of course performed by depositing the seed on the surface; if in the depth, the end of the pipette is pushed down into the depth of the material and the seed there deposited. The pipette is then altogether withdrawn and the plug replaced as before. The new tube is then placed in a beaker on a cushion of cotton-wool, and exposed to the required temperature in the incubator.

If we have, however, a culture-fluid or any fluid that contains, as the microscopical examination proves, various species of organisms, which we wish to isolate, *i.e.* if we wish to introduce into a new culture-tube only one species, then the method of Klebs of "fractional cultivation," or the method of Lister and v. Nägeli of "dilution," is resorted to.

The "fractional cultivation" consists in the attempt to isolate by successive cultivations the different organisms that have been growing previously in the same culture. If we take up by means of a capillary pipette a trace of the culture-fluid, and inoculate with traces of it in the manner above described a series of new culture-tubes containing various nourishing materials, and expose these tubes in the incubator to a definite temperature, say 35° C., then the chances are that in the first twenty-four or thirty-six hours not all the different species of organisms sown out will have increased equally in numbers in all tubes; most probably only one species in each tube, *i.e.* the one that grows best in this particular medium and at this particular temperature, will be found to have increased to an enormous extent, while the others have made little or no progress as yet. The nourishing fluid appears turbid, and filled chiefly with the one kind of organism. Now draw out with a fresh capillary pipette a minute droplet of this new culture and inoculate with a trace of it a new culture-tube. The chances are that you inoculate only one kind, that is, the one which is most abundant or perhaps is solely present. After twenty-four hours' incubation this new tube contains now probably only one kind of organism. To make it quite certain, inoculate from this a new culture-tube in the same manner, and now you probably have sown only a single species. In this manner by continued transference it is possible to obtain cultures with only one species of organisms. Many conditions, such as naked-eye appearances of a particular kind, coloration of the culture-medium, formation of a pellicle, the quantity of growth in a given time, soon indicate whether we have the desired single

species; in some instances it is, however, extremely difficult to isolate after this method.

The method of "dilution" means diluting the culture-fluid containing the various species to a very large extent with some sterile indifferent fluid, such as well-boiled saline solution of 0.6 per cent., and then inoculating new tubes with a droplet of this greatly diluted material. For this purpose draw into a rather large pipette a tiny droplet of the old culture-fluid, then pass the pointed end of this pipette into a test-tube or flask (plugged) containing well-boiled saline solution, and draw up a quantity of this solution so as to greatly dilute (1000-fold or more) the droplet of culture-fluid, and with this inoculate then a series of new culture-tubes containing different nourishing material, using always only a trace for inoculation. In this way it is probable that, owing to the great dilution, the trace of a droplet of this mixture used for the new inoculation contains only one species. Using a series of new culture-tubes and inoculating them thus, after twenty-four hours of incubation it will be found that some tubes have not received any seed, others only one species. If it be required to dilute the original fluid greatly, say if it teems with different organisms, then a droplet of this is placed into a large flask containing the well-boiled saline solution, so that a dilution of 1 in 1,000,000 or more can be effected.

The two methods *i.e.* that of fractional culture and of dilution, may be successfully combined in this way: from the first or second new culture, established after the method of fractional cultivation, in which after twenty-four or thirty-six hours one species greatly predominates, draw out with a large capillary pipette a droplet, and dilute this to a great extent with the saline solution, as described above, and now inoculate with a trace of this mixture a new culture-tube. Or, if after twenty-four hours' incubation the microscope reveals in this further culture more than one species, continue the process of dilution and inoculation for a further generation. Thus it is possible to obtain cultures of only one species, although the original fluid contained several species of organisms.

2. *Inoculations with Blood, Juices, and Tissues.*—To establish a cultivation from blood of a dead animal, cut open the thorax by removing the sternum with clean scissors, cut open the pericardial sac, pierce with the pointed end of a fresh capillary pipette the wall of the right ventricle or right auricle, and allow a drop or two of blood to ascend into the pipette, or if a larger quantity is required suck it up. Withdraw the pipette and inoculate new culture-tubes as above. Or, if blood of a

large vein is required, separate the vessel with clean instruments, and make a small incision with clean scissors and push the pointed end of the capillary pipette well forward. If juice of a lymphatic gland, or spleen, or other parenchymatous organ be required, pierce the organ after having washed its surface with solution of perchloride of mercury (Koch), with the pointed end of a capillary pipette, then push it into the part required for a little distance, and squeezing the organ press a drop or two of the juice into it. The same procedure is adopted when the pus of an abscess is required, the wall of which can be pierced with the pointed end of the capillary pipette. If not, a slight incision is made and the pipette introduced through this into the abscess. If blood of a living animal is required, expose a vessel with clean instruments, make a small incision with clean scissors, push through this incision the pointed end of the capillary pipette well forward, and allow the blood to rise into the capillary tube. If blood of a living human being is required, clean well with soap and water and then with strong carbolic acid or perchloride of mercury solution the tip of a finger, make a venous congestion in the last phalanx by compressing it with a corner of a handkerchief, prick the volar skin of the phalanx with a clean (overheated and cooled) needle, and plunging the pointed end of the pipette into the drop of blood allow a droplet to ascend into the capillary tube of the pipette.

If solid tissues or parts of tissues are required, *e.g.* the base of an ulcer, a tubercle of the liver, spleen, or lung, it is possible to squeeze into the capillary tube of a pipette, after pushing its pointed end into the part, a small droplet of juice of the part required; but if this be not practicable, *i.e.* if a solid particle be required, then follow Koch's method. This is as follows: cut with clean scissors or scalpel into the part, dig out rapidly with the point of a needle or platinum wire previously overheated in the flame of a burner a small particle, and quickly introduce this into the culture-tube to the place required, *e.g.* surface or depth of a solid or fluid nourishing material. Of course in this case the cotton-wool must be altogether lifted, and therefore contamination with organisms is possible. But inoculating several tubes at once and performing the operation quickly, one always succeeds in getting some of the tubes without any air-contamination. I have made numerous inoculations with solid particles (tubercles) in this manner, and like Koch have seen only a small percentage of tubes going bad through contamination with air-organisms.

The same plan, *i.e.* of using the clean point of a needle or

platinum wire for taking up the material to be used for inoculation, is resorted to if one has to deal with the culture of solid nourishing material, on or in which the organisms are growing that we want to transplant either for inoculation of a new tube or of an animal. A useful method, which does not require the lifting out of the plug at all, and which can easily be employed in the last case, is this: deposit from the pointed end of a capillary pipette a droplet of some sterile fluid (broth or thoroughly boiled saline solution) on the spot of the solid medium on which the organisms are growing, then scratch this spot with the end of the capillary pipette in order to get the organisms off from the solid basis and mixed with the drop of fluid deposited there, then let this drop again ascend into the end of the capillary pipette, and withdraw this altogether. All this can be done without lifting out the cotton-wool plug of the test-tube or flask in which the growth is proceeding.

If one has to use a particle of tissue the surrounding portions of which are probably contaminated by putrefactive organisms, e.g. a tubercle in the lung or a tubercle in the spleen, it is well to follow Koch, and to disinfect the surrounding parts by just washing them with a dilute solution of corrosive sublimate, and then to remove these parts with clean scissors so as to obtain the central particle which one wishes to use for inoculation: of course one must not steep the whole organ in sublimate solution, since this would naturally destroy all organisms.

All these methods can be easily modified according to the requirements of the special cases, and it is not necessary here to give more than what has already been described in the preceding.¹

In order to observe in a microscopic specimen the gradual changes in the growth of a micro-organism, there are several methods employed. In all of them it is of course necessary to keep the specimen heated up to the desired temperature.

The simplest method consists in sowing the organisms on a suitable nourishing material in a small glass cell, fit to be placed on the stage of a microscope and to be there observed even with high powers, similar to those cells which Koch has used in his studies on bacillus anthracis. Such a glass cell consists of a glass slide, in its centre a concave pit, not too large, and capable of being quite closed up by an ordinary cover-glass, the edges of which fasten by means of clean paraffin or olive oil. Place with a clean needle a speck of spleen pulp

¹ Compare also Koch, *Untersuchungen über pathogene Bacterien*, in *Berichte aus dem k. Gesundheitsamte*, Berlin, 1881; and *Die Aetiologie d. Tuberculose*, *Berlin. klin. Wochenschrift*, No. 15, 1882.

of an animal dead of anthrax into a drop of nourishing material, fluid or solid, on the centre of a clean cover-glass, the edges of which have been prepared as just mentioned, and fasten this on the above slide so that the specimen faces the concave pit : expose this so prepared specimen to a constant temperature, either by placing it in the incubator and examining it with the microscope from hour to hour, or on the warm stage (Stricker, Ranvier) used in histological work for directly observing the influence of temperature on the various cells and tissues ; or, place it simply on the stage of the microscope and expose the whole (*i.e.* microscope and all) in a suitable warm chamber (after Klebs), but so that the chamber allows light to pass by means of a small window to the mirror of the microscope, while the eyepiece is so arranged as to project through a hole in the




FIG. 7.—A GLASS CELL, FOR OBSERVING UNDER THE MICROSCOPE THE PROGRESS OF GROWTH OF MICRO-ORGANISMS.

The upper figure shows the cell in perspective ; the lower figure in profile or cross section.

- A. Glass slide.
- B. Cover-glass.
- C. Glass ring forming the wall of the chamber.
- P. Drop of nourishing material in which the micro-organisms grow.

upper wall of the chamber. The plan which I generally follow is with slight modifications that of Koch.

A glass cell (Fig. 7) is made by cementing a glass ring, $\frac{3}{4}$ — $\frac{7}{8}$ inch in diameter and about $\frac{1}{8}$ — $\frac{1}{4}$ inch high, on to an ordinary glass slip. The chamber of this cell is well cleaned with absolute alcohol. A thin cover-glass, square or round, about one inch in breadth, is well heated by holding it for a few seconds over the flame of a gas-burner or spirit-lamp. On the upper edge of the above glass ring is placed with a camels' hair brush a thin layer of clean olive oil ; a droplet of water is deposited on the bottom of the cell in order to keep this

afterwards well supplied with moisture ; a drop of the sterile nourishing material (broth, aqueous humour, hydrocele fluid, blood serum, liquefied gelatine mixture, liquefied Agar-Agar mixture, &c.) is then deposited by means of a capillary pipette on to the centre of the cover-glass ; then the point of a capillary pipette or needle containing the material it is desired to sow is rapidly plunged into the drop of the nourishing material (or if this is solidified is deposited in lines or points on the drop of nourishing material), the cover-glass inverted and placed on to the glass ring : the layer of olive oil keeps the edges of the cover-glass air-tight on the glass ring. This cell is then placed into the incubator and exposed there to the desired temperature. Microscopic examination is carried out from time to time to watch the progress made. This can be done with high powers, since the growth is taking place on the lower surface of the cover-glass.

Although contamination with air-organisms is not excluded, still it is possible by making several specimens at the same time and operating rapidly, to obtain pure cultures. This glass cell can be also watched on a warm stage, or in a Klebs' warm chamber.

M. Nacet of Paris has designed a glass cell, in which the drop of nourishing material is deposited on to the bottom of the cell, the glass slip being here replaced by a very thin glass ; but then there is a peculiar arrangement in the microscope, by which the lower surface of the glass cell, *i.e.* the one nearest to the growth, is directly subjected to microscopic observation.

After what has been said above about inoculation of solid and fluid nourishing media with solid matter, it is not necessary to dwell specially on the method of inoculation with earth or similar substances.

3. *Examination of Water for Micro-organisms.*—Most water contains bacteria of some kind, as has been shown by direct experiment by Burdon Sanderson.¹ If any sample of water is to be examined for micro-organisms, particularly bacterial forms, it is allowed to stand for a few hours, till most of the particulate matter is settled, and then with a capillary pipette a little of the fluid and sediment is drawn out and used for (a) microscopic specimens to be examined fresh ; (b) microscopic specimens prepared after the Weigert-Koch method, *i.e.* by spreading out on a cover-glass a thin layer, drying it, staining it with suitable aniline dyes, *e.g.* Spiller's purple, gentian violet,

¹ *Reports of the Medical Officer of the Privy Council, 1870.*

methyl blue, or magenta, washing with water, then spirit, then distilled water, then drying, and finally mounting it in Canada-balsam solution. (c) Test-tubes containing sterile nourishing material (broth, Agar-Agar mixture, gelatine mixture, Cohn's or Pasteur's fluid) are inoculated in the manner described previously, i.e. by piercing the cotton-wool plug with the pointed end of the capillary pipette. These test-tubes are then exposed in the incubator, and after one or two days or more, a sample is withdrawn with a capillary pipette, and used for microscopic examination. As a rule, after a day or two of incubation we can already distinguish with the unaided eye whether there are any organisms present, the nourishing fluid either being uniformly turbid—this is generally the case—or there being a growth at the bottom of the fluid. But of course the microscopic examination only shows what kind of organisms are present. New cultivations are made from this one, if any are required. (d) A good plan of recognising easily that there are present various kinds of organisms in such cultures is one similar to that recommended by Professor Angus Smith.¹ Sterile gelatine broth or gelatine only, contained in sterile test-tubes plugged with sterile cotton-wool, is liquefied, but of course not heated to more than about 35°—40° C., then inoculated with the water (to be tested), by means of the capillary pipette; after inoculation the gelatine is mixed by shaking the test-tube slightly. In this way the organisms present in the water are distributed in the gelatine. Then the gelatine is allowed to set and is kept in this solid state. The organisms being distributed in the gelatine, after some days' growth are noticeable as clusters which gradually increase in extent and are distributed in various parts of the medium. The various species, owing to difference of growth, form clusters differing in aspect, size, and arrangement.

4. *Examination of Air.*—The simplest plan to test for the presence of organisms in the air is to draw out the cotton-wool plug of several test-tubes or flasks containing the sterile nourishing material, or, if this be boiled, potato, paste, or gelatine (see p. 14), to expose their surface, and to leave it thus for variable periods, from a few seconds to several minutes. Then replace everything and expose the material to incubation, or keep it only at the ordinary temperature of the room. Another method is to collect the particles present in the air on glasses moistened with pure glycerine (Maddox), and then to make microscopic specimens or inoculate tubes with this glycerine.

¹ *Sanitary Record*, p. 344, 1888.

A method which is very useful is the one recommended by Cohn and Miflet.¹ The principle of it is, that by means of an aspirator, an air-pump of any kind—*e.g.* a Sprengel pump, or simply the fall of water—air of a particular locality is drawn into one, two, or more Wolff's bottles (each with the ordinary two bent glass tubes), connected with one another by short pieces of india-rubber tubing, and containing the sterile material in which the organisms are required to grow. All bottles and tubes being of course sterile, the plugging of the tubes after the air has passed is done with sterile cotton-wool. Any quantity of air for any length of time can thus be passed through a series of such bottles, the one that receives the air first being of course most contaminated.

The bottles are after the experiment placed in the incubator if required, the outer end of their tubes being plugged with cotton-wool.

Miquel² has carefully described many ingenious methods for the study of air-organisms

¹ *Zeitschr. f. Biol. d. Pf.* iii. 1, p. 119.

² *Les Organismes vivants de l'Atmosphère*, Paris, 1863.

CHAPTER VI.

MORPHOLOGY OF BACTERIA.

BACTERIA are minute organisms not containing chlorophyll, and multiplying by fission—hence the term *schizomycetes* (v. Nägeli). They are composed of a kind of protoplasm, the mycoprotein of Nencki, and are invested with a membrane, which is composed chiefly of cellulose and a certain amount of mycoprotein (Nencki).

Their contents are transparent and clear, but sometimes contain minute bright granules of sulphur (Beggiatoa). Owing to the cellulose membrane they resist the action of acids and alkalies. Many species of bacteria—micrococcus, bacterium, spirillum—are able by rapid multiplication to form colonies; the individuals are then embedded in a hyaline gelatinous matrix produced by them, this is also mycoprotein. Some species are possessed of one or two straight or slightly spiral cilia or flagella, and thereby they are capable of locomotion, darting through, or spinning round, in the fluid in which they are suspended. Such is the case with many kinds of bacteria, bacilli, and spirilla.

Bacteria grow best when left undisturbed; movement of the vessel in which they grow is not advantageous. Light and electricity do not appear to have a decided influence, since most of them grow well in the light. According to Cohn and Mendelssohn,¹ strong electric currents have a noxious influence on the growth of micrococci.

Some bacteria require free access of oxygen, and are called aerobic (Pasteur); others grow without free oxygen, and are anaërobic (Pasteur). All require for their growth certain nourishing materials containing carbon and nitrogen. Water

¹ Cohn's *Beitr. z. Biol. d. Pfl.* Bd. iii. 1.

is an essential element for them, and a certain temperature is in many instances a stimulant of their growth. Most pathogenic bacteria require for their propagation a temperature varying in the different cases between 18° and 40° C. The bacteria obtain their nitrogen from organic compounds; some are capable of obtaining it from compounds as simple as ammonium tartrate; others, especially pathogenic organisms, require much more complex combinations, such as occur in the animal body. Carbon they obtain likewise from organic compounds, such as carbohydrates, amongst which sugar is the chief, and vegetable acids combined as salts are also to be mentioned. It is essential for all that certain inorganic salts, phosphates, potassium and sodium salts, should be present, since their own substance contains a large percentage of it—4 to 6 per cent.

While all are capable of disintegrating organic combinations containing nitrogen, they in their turn help to produce certain chemical products, which in some cases are definite for a definite species (see below). Such is the case with the various bacteria connected with the fermentations producing lactic acid, butyric acid, and acids belonging to the aromatic series. On many bacteria connected with putrefaction, and also on some pathogenic organisms, these chemical products have a deleterious effect. Small quantities impede their growth, and sufficiently large quantities kill them altogether.

Most bacteria are killed by heat below the temperature of boiling water, many of them when exposed for several hours to a temperature above 50°—60° C. Exceptions are the spores of bacilli, which in some instances (spores of hay bacillus, Cohn) require exposure to the heat of boiling water for as much as half an hour. By raising the boiling point above 100°, it does not require more than a few minutes to kill them (Sanderson).

Drying destroys most bacteria, except the spores of bacilli. Freezing destroys likewise most bacteria, except the spores of bacilli, which survive exposure to as low a temperature as -15° C., even when exposed for an hour or more. No spores survive exposure to a temperature of 120° C.

Amongst those substances which inhibit the growth of, or altogether destroy the bacteria, are carbolic acid, salicylic acid, thymol, &c.; corrosive sublimate is the most powerful (Koch) since even solutions as weak as 1:300,000 are said to inhibit the growth of bacillus anthracis.

The best classification of bacteria is that given by Cohn,¹ and this we shall adopt: (1) spherobacteria micrococci;

¹ *Be tr. s. B. d. Pfl. Bd. I.*

(2) bacteria or microbacteria; (3) bacilli or desmobacteria; (4) spirilla, (5) spirochætæ. There are also various kinds which approach one or the other of these, *e.g.* ascococcus, sarcina, leptothrix (Beggiotoa), cladothrix, streptothrix, &c. (see below).

I shall not attempt to give an exhaustive description of the morphological characters of all micro-organisms, but shall limit myself to those forms which are related in some way or other to diseases.

CHAPTER VII.

MICROCOCCUS (Hallier, Cohn).

By the specific term micrococcus is understood a minute spherical or slightly oval organism (spherobacterium, Cohn), that like other bacteria divides by fission (schizomycetes), and that does not possess any special organ, cilium or flagellum, by using which it would be capable of moving freely about. Micrococci, like other granules when suspended in a fluid medium, show (Brownian) molecular movement. Micrococci propagate always by simple division, never by any other means, *e.g.* gemmation and spores. All assertions to the contrary are based on incorrect observations. All micrococci possess a delicate membrane of cellulose, and owing to this resist the action of alkalies and acids. The contents are homogeneous and highly refractive while active, pale when inactive. They consist like those of other bacteria of mycoprotein (Nencki). The size of micrococci varies within considerable limits, say 0.0008—0.002 millimetres, or even a little more. Micrococci vary greatly as regards both size and mode of growth. All multiply by slightly elongating and then dividing by a transverse constriction into two: a dumb-bell; each of these again divides into two, either transversely or in the same direction as before. The new elements of successive divisions may remain connected, and thus form a chain (or mycothrix, Itzigsohn and Hallier; torulaform string, Cohn), or they separate into single organisms or dumb-bells. In some species there is a pre-eminent tendency to form chiefly dumb-bells, in others to form shorter or longer chains generally more or less curved.

Such exquisite chains one meets with sometimes in serum of blood exposed to the air for some days, and in pleural and

peritoneal exudations of animals dead for a few days. I have seen in an artificial culture made by my friend Mr. A. Lingard



FIG. 8.—MICROCOCCI OF PUTRID HUMAN SPUTUM.

1. Single micrococci and dumb-bells.
2. Short chains.
3. A long chain.
4. A zoogloea.

This and all subsequent figures are drawn under a magnifying power of about 700 diameters except stated otherwise.




FIG. 9.—FROM THE SAME PUTRID SPUTUM AS IN PREVIOUS FIGURE. THE MICROCOCCI ARE LARGER.

1. Dumb-bells.
2. Sarcinae.
3. A small zoogloea, in reality consisting of four sarcina-groups.

from a blister in a rabbit's ear the most exquisite convolutions of threads of micrococci. (See Fig. 10.)



FIG. 10.—PART OF A CONVOLUTION OF CHAINS OF MICROCOCCI; FROM AN ARTIFICIAL CULTIVATION STARTED WITH THE SERUM OF A BLISTER OF A RABBIT'S EAR.

A dumb-bell is also called a diplococcus (Billroth). Between the individuals of a dumb-bell there is always noticeable a short pale intervening bridge.

Some species are specially characterised by dividing into a dumb-bell, and each of the elements dividing again transversely into a dumb-bell; a group of four (tetrad or sarcina-form) is thereby produced. Some species are occasionally met with, particularly in products of air-contamination, in which the four individuals are closely pressed against one another, and then each assumes more or less the shape of a cube, a true sarcina (see below). But each of these cubes divides into four small micrococci arranged as a small sarcina, so that a sarcina-within-sarcina-form results.




FIG. 11.—GIANT MICROCOCCI, FROM SAME PUTRID SPUTUM AS IN PREVIOUS FIGURES.

1. Dumb-bells.
2. Division of dumb-bells into sarcina.
3. Incomplete division into sarcina.




FIG. 12.—SARCINA-MICROCOCCUS, FROM AN ARTIFICIAL CULTIVATION.

1. The elements of each sarcina-group of four appears single.
2. The elements incompletely divided into secondary groups.
3. Each element of the previous groups has divided into four small micrococci.

In many instances the individual members resulting from division remain closely adherent without any definite arrangement, and thus form smaller or larger continuous masses, *zooglæa* or colonies, in which the individuals appear embedded in a hyaline gelatinous matrix; the amount of this varies in the different species; in some there is little of the matrix actually visible, the micrococci being in close juxtaposition, in others it is easily recognised, the interstices between the individuals being measurable.

In some of the pigmented species (see below) the interstitial matrix contains the pigment. *Zooglæa* masses always present themselves as uniformly granular, the granules or micrococci being of the same size.

True micrococci never elongate to form rods, although in certain rod-like bacteria the individual elements sometimes assume the shape of spherical elements (see below).

Some species of micrococci form after some days a pellicle on the surface of the nourishing material, although there is also an abundance of these micrococci in the depth of the nourishing

material. This pellicle is composed of zoogloea, and after some time bits of it, or the whole, sink to the bottom if the medium is fluid. Micrococci that thus form pellicles are pre-eminently aërobie (Pasteur), i.e. require a great deal of free oxygen, which they receive from the air to which they are exposed on the surface of the nourishing material. Other species do not require free oxygen (anaërobie, Pasteur), and therefore grow well in the depth and do not form a superficial pellicle. There is a marked distinction in this respect between different species. The micrococci occurring in connexion with disease are anaërobie.

When cultivated in the incubator in suitable fluid nourishing material, they produce after a day or two general turbidity.

Micrococci may be divided, according to their chemical and physiological function, into: (a) septic, (b) zymogenic,¹ (c) chromogenic, and (d) pathogenic micrococci.

(a) The *septic micrococci* are micrococci that occur with other septic bacteria, wherever there is decomposition of organic matter in solids or in fluids. There exists a large number of species of such micrococci, differing from one another in size and mode of growth. They are widely distributed in the air, and contamination by air is often followed by the appearance of micrococci. They also occur in the body of man and animals wherever there is dead tissue, in which they grow well and copiously. Of this kind are the micrococci found in ordinary pus (Ogston), in the normal oral cavity (on the filiform papillæ of the tongue and on the mucous membrane), in the bronchial secretion in ordinary catarrhal exudations (nasal cavity, bronchi, &c.), on the free surface of intestinal and other ulcerations, and in the cavity of the small and large intestine.

(b) *Zymogenic micrococci* are micrococci associated with definite chemical processes. (a) *Micrococcus ureæ*, causing the ammonical fermentation of urine (aërobie, Pasteur), occurs singly, as dumb-bells or chains, and as zoogloea. (β) The micrococcus of the mucoid wine fermentation produces (Pasteur) a peculiar mucoid change in wine and beer, and occurs chiefly in chains. (γ) The micrococcus causing phosphorescence in putrid meat and fish (Pflüger) forms chiefly zoogloea (aërobie).

¹ I adopt this term from Flügge: *Fermente und Mikroparasiten*, Leipzig, 1838.

(c) *Chromogenic micrococci* (Schröter, Cohn). These micrococci are characterised by their power of forming pigment of various colours. They grow well at ordinary temperatures, and occur chiefly as zooglœa; they differ from one another by forming different pigments. The thicker the layer the more marked is the pigment. This is either soluble in water or it is insoluble, and therefore remains limited to the cells and their interstitial substance. The cells are spherical (*Micrococcus prodigiosus*, *chlorinus*, *fulvus*) or slightly elliptical (*M. luteus*, *aurantiacus*, *cyaneus*, *violaceus*). They are all aërobic and produce this pigment only when there is free access of air. They grow best on boiled potato, bread, paste, and boiled-egg albumen. They can be transplanted, and always produce the same pigment. When growing and kept in the depth of a solid nourishing material, i.e. removed from the free surface, they grow as colourless micrococci. They abound in the air—in some localities and at certain seasons



FIG. 13.—OVAL MICROCOCCI WHICH POSSESS A BLUE COLOUR, MICROCOCCUS CYANEUS, SINGLY AND IN DUMB-BELLS.

more than at others. (a) *Micrococcus prodigiosus* is blood-red, the colour is lodged not in the micrococci but in the interstitial substance, and is insoluble in water, soluble in alcohol; it occurs chiefly as zooglœa, in the shape of smaller or larger droplets. The cells are the smallest of all pigment-micrococci. (b) *Micrococcus luteus* is yellowish, and the pigment is insoluble in water. It occurs also in fluid nourishing material, forming a pellicle. I have met with it in the air, and have seen it in fluid pork broth, where it grew very abundantly at a temperature of 32°–38° C. It was found as single cells or dumb-bells, and formed a thick pellicle on the surface, which after some time sank down into the fluid, the pellicle retaining a pale yellow colour. (c) *Micrococcus aurantiacus* grows on boiled-egg albumen, chiefly as zooglœa. The pigment is soluble in water. (d) *Micrococcus cyaneus*, *violaceus*, *chlorinus*, and *fulvus*, produce blue, violet, green, and brown pigment respectively. The first two grow well as zooglœa of elliptical cells on boiled potatoes, the third on boiled-egg albumen, and the last is met with on horses' dung.

Clathrocystis roseo-persicina (Cohn), peach-coloured bacterium, *Bact. rubescens* (Lankester), is an organism of about 0·0025 mm. in diameter, spherical or oval and of a bright red colour. The cells differ from *micrococcus prodigiosus*, not only in their greater size and their intrinsic colour, but also in this—that having formed zoogloea-masses there are gradually developed cavities or cysts therein, which are filled with water, while the coloured cells occupy their periphery. The cysts ultimately break up. Together with this organism occur other pink-coloured organisms described by Cohn as *monades*.

Monas vinosa, spherical cells about 0·002–0·003 mm. in diameter.

Monas Okenii, cylindrical cells, 0·008–0·005 mm. long, 0·005 mm. broad, flagellate.

Rhodomonas rosea, spindle-shaped, 0·004 mm. broad, 0·02–0·03 mm. long, flagellate.

Monas Warmingii, spindle-shaped, 0·008 mm. broad, 0·015–0·020 mm. long, flagellate.

Ascococcus—Billroth first described certain peculiar spherical, oval, or knobbed masses of minute micrococci, which he




FIG. 14.—ASCOCOCCUS BILLROTHI, (AFTER COHN).

found in putrid meat infusion. Each of the masses is enveloped in a resistant firm hyaline capsule of about 0·010 to 0·015 mm. 'thickness. The masses are of various sizes, from 0·02 to 0·07

mm. in diameter, and are composed of small spherical micrococci. Cohn found them also in his (Cohn's) nourishing fluid (see Chapter II. A. 7) where they produce the peculiar smell of cheese. They are capable of changing acid nourishing material into alkaline. Cohn called the organism *ascococcus Billrothi*.

Sarcina ventriculi.—Goodsir was the first to describe in the vomit of some patients, peculiar groups of four cubical cells, with rounded edges, and closely placed against one another. These *sarcinæ ventriculi* are of a greenish or reddish colour. The diameter of the individual cells is about 0.004 mm. They are found in the contents of the stomach of man and brutes in health and disease, where the groups of four cells form smaller and larger aggregations. Occasionally small *sarcinæ* occur on boiled potatoes, egg albumen, and gelatine exposed to the air. These *sarcinæ* are considerably smaller than the *sarcina ventriculi*, and when in large quantities have a yellowish tinge. Like the *sarcina ventriculi* they are in groups of four, and these again occur in larger or smaller aggregations and zooglœa. I have cultivated them successfully through many generations in pork broth, beef broth, mixture of gelatine and broth, at ordinary temperatures and in the incubator; more easily however at ordinary temperatures.

(d) *Pathogenic micrococci*.—Many of these are connected with disease. In the pus of open wounds,¹ and in that of closed abscesses, occur micrococci, singly, in dumb-bells, and in colonies or short chains,² but there are certain acute inflammations, e.g. that produced by subcutaneous injection of turpentine, the pus of which does not contain micrococci or any other organism.³

The secretion of open ulcers, such as occur in ordinary acute inflammations of the skin and mucous membranes, in ulcerations of the throat due to scarlatina, in every ulceration of the intestinal mucous membrane, in the lymph of the vesicles of the skin and mucous membranes of the mouth occurring in various kinds of inflammations, there are almost always present micrococci in dumb-bells, often also in beautiful chains. In the ulcers and abscesses they often form continuous masses, i.e. zooglœa, encroaching upon the tissue of the base of the ulcer. To this category belong the minute micrococci

¹ W. Cheyne, *Path. Transact.* xxx.

² Ogston, "Micrococcus in Acute Abscesses," *Br. Med. Journal*, March 12, 1881.

³ Uskoff, *Virchow's Archiv*, vol. 86, i. p. 150.

(about 0·0005 mm. in diameter) which Klebs described as *microsporon septicum*, found in and around wounds. The spread of purulent inflammation in connective tissues and in parenchymatous organs is often, if micrococci are present in the original focus, associated with a corresponding spreading of the micrococci; these easily grow into all the spaces and crevices of the tissues, but whether this spreading of the micrococci is merely of secondary importance, *i.e.* concomitant




FIG. 15.—FROM THE BASE OF AN ULCER OF THE MUCOUS MEMBRANE OF THE LARYNX IN A CHILD THAT DIED OF ACUTE SCARLATINA.

1. Nuclei and fibres of the tissue.
2. Zoogloea of micrococci.

with or subsequent to the spreading of the inflammation, or whether it is the primary cause as some assume, is not clear, and requires definite experimental proof.

In all cases of diarrhœa the secretions of the bowels swarm with micrococci. In typhoid fever, clumps of micrococci may be found very extensively on the ulcerations of the bowels, and in the mucous membrane surrounding the ulcerations, and may be even traced into the mesenteric glands and the spleen.¹

In dead tissues within the living body, such as occur after embolism, and in the case of various infectious maladies, micrococci may be found in colonies, *i.e.* as zoogloea, in the blood-vessels and in the parts around. The same holds good for the disseminated abscesses and necroses occurring in connexion with surgical pyæmia. In this malady masses of micrococci have been found in many of the affected organs.²

¹ Klein, *Reports of the Medical Officer*, 1876. Letzerich, Sokoloff, Fischel, &c.

² "Report of the Committee of the Pathological Society," *Pathol. Transactions*, vol. xxx.

Wassilieff¹ has shown that these micrococci only occur after the death of the tissue or tissues, that in these they may multiply so as to form extensive colonies, and that therefore the presence of these micrococci is only a secondary phenomenon.




FIG. 16.—CAPILLARY BLOOD-VESSELS OF NECROTIC MASSES FROM THE LIVER OF A MOUSE. THE CAPILLARIES ARE DISTENDED BY, AND FILLED WITH, ZOOGLEA OF MICROCOCCI.

In pneumonia accompanying certain infectious maladies, *e.g.* typhoid fever, tuberculosis, and even in severe catarrhal pneumonia, large masses of micrococci may occur in the air-cells.




FIG. 17.—FROM A PREPARATION OF THE BLOOD OF A CHILD ILL WITH INFANTILE DIARRHŒA.

1. Blood-discs.
2. Dumb-bells of micrococci.

In those cases where lobules and whole lobes become transformed into solid structures—grey hepatisation—masses of micrococci may be found in the air-cells, and even growing into the blood-vessels in which stasis had set in. Such is the case

¹ *Centralblatt f. d. med. Wiss.* No. 52, 1881.

in pleuro-pneumonia of cattle and in the pneumonia of swine fever. Pasteur has cultivated the micrococci in swine fever, and thought that he had reproduced the malady by inoculation. But this is not the case. The micrococci, although very abundantly present in the bowels and in the body,¹ have nothing to do with the malady. Pasteur's inoculations with the cultivated micrococci are quite fallacious; his positive results are no doubt accounted for by accidental air-infection, for this malady is highly infectious, and unless the most rigorous precautions are taken to obviate infection through the air, positive results may be obtained which in reality are due to accidental air-infection.²

Micrococci occur always normally in large quantities in the fluids (saliva and mucus, &c.) of the nasal and oral cavities, pharynx, larynx, and trachea; they are derived no doubt from the atmosphere. On the papillæ filiformes of the tongue they form in some cases large masses.³ Pasteur⁴ has inoculated rabbits with the saliva of a child that suffered from hydrophobia, and having cultivated artificially the micrococci present in this saliva, thought to have discovered that a micrococcus (*microbe spéciale*)⁵ is the cause of hydrophobia. That saliva of the healthy dog and of man inoculated subcutaneously into rabbits sometimes produces death in these animals (Senator) had entirely escaped his notice, and Sternberg⁶ has proved this in an extensive series of experiments. His own saliva proved sometimes fatal to rabbits. They die of what is called septicæmia, and Sternberg thinks it due to the micrococci; but this is not to be considered as proved.

All these micrococci stand therefore in no definite causal relation to the respective maladies, but are probably only of secondary importance.

The following micrococci are considered to stand in an intimate relation to specific diseases:—

1. *Micrococcus variolæ et vacciniæ*.—Chauveau⁷ was the first to prove experimentally that in vaccinia and in variola the active principle is a particulate non-diffusible substance.

¹ Reports of the Medical Officer, 1878, 1879.

² *Ibidem*.

³ Butlin, "Fur of the Tongue," *Proceedings of the Royal Society*, 1880.

⁴ *Comptes Rendus*, xlii.

⁵ It is not quite clear whether this *microbe spéciale* is a dumb-bell micrococcus or a bacterium termo; it is quite possible that it is the latter, viz. a rod constricted in the middle. If so, it would appear identical with the bacterium that produces Davaine's septicæmia in rabbits (see Chapter viii).

⁶ Bulletin, April 30, 1881, National Board of Health, U S A.

⁷ *Comptes Rendus*, 1863.

Burdon Sanderson confirmed and extended this.¹ Cohn² proved that the lymph of vaccinia and variola contains numerous micrococci. Weigert³ showed for human small-pox, Klein⁴ for sheep-pox, that the lymphatics of the skin in the region of the pock are filled with micrococci, and Pohl-Pincus⁵ traced their passage through the epidermis at the point of




FIG. 18.—MICROCOCCLUS IN THE FRESH LYMPH OF HUMAN SMALL-POX.

1. Singly.
2. In dumb-bells.
3. In short chains.

vaccination in the calf. The micrococci are very minute, according to Cohn's estimate 0.0005 mm. and less in diameter, single or in dumb-bells, or in shorter or longer chains, or in small groups. When cultivated on the warm stage, they form very long chains and colonies. In connexion with this it



FIG. 19.—LYMPHATIC VESSEL FROM THE SKIN OF A POCK IN SHEEP-POX

The vessel is filled with micrococci.

must be mentioned, as stated on a former page, that similar micrococci occur also in the fluid contents of vesicles in the skin produced by various non-infective inflammations. To make it sure that the micrococci are the active principle, *i.e.* the *causa morbi*, it would be necessary to artificially cultivate them through several generations, and then, by re-inoculating

¹ Reports on the Intimate Pathology of Contagion.

² Virchow's Archiv, 1872, Keber, Haller, Zürich. ³ Mediz. Centralblatt, 1871.

⁴ Phil. Transact. 1874.

⁵ Vaccination, Berlin, 1882.

them, to reproduce the disease. This essential link in the evidence is, however, still wanting.¹

2. *Micrococcus erysipelatosus*.—The micrococcus is very minute, smaller than that of vaccinia. Lukomsky² showed that, at the margin of an erysipelatosus zone, that is the part where the disease is progressing and marked by the characteristic redness and swelling, the lymphatics of the skin are filled with zooglœa of micrococci, and the injection of these vessels keeps pace with the progress of the erysipelatosus process. Oth³ cultivated these micrococci artificially, and with such cultures produced by inoculation erysipelas in rabbits. Fehleisen⁴ placed this beyond any doubt, inasmuch as he produced successive cultures of these micrococci (derived from the lymphatics of erysipelatosus human skin), and then by re-inoculation produced the disease not only in rabbits but also in man. Fehleisen found the micrococci only in the lymphatics of the affected parts, and these he cultivated artificially for fourteen generations—which it took two months to do—on peptonised meat-extract gelatine, and solid serum. The micrococci form a whitish film on the top of the nourishing material, and when inoculated into the skin (ear) of rabbits, a characteristic erysipelatosus rash makes its appearance after from thirty-six to forty-eight hours, and spreads to the root of the ear, and further on to the head and neck. The animals do not, however, die from it. In the human subject he produced typical erysipelas after inoculation with the pure cultivated micrococcus in fifteen to sixty hours. These inoculations were justifiable because they were undertaken with a view to cure certain tumours. Thus one case of lupus, one case of cancer, one case of sarcoma, were considerably affected, and to the good of the patient.

Fehleisen also in several instances carried out successfully a second inoculation within a few months. The same observer also found that a 3 per cent. solution of carbolic acid and a 1 per cent. solution of corrosive sublimate destroys the vitality of this micrococcus.

3. *Micrococcus diphtheriticus*.—Buhl, Hüter, and Oertel had shown that in diphtheria the membranes include micrococci. Oertel⁵ found this micrococcus in large numbers, not only in

¹ See the prize announcement of the Grocers' Company, London, 1883.

² *Virch. Archiv*, vol. 60.

³ *Archiv f. exp. Path.* Bd. i 1874.

⁴ *Die Aetiologie d. Erysipels*, Berlin, 1883.

⁵ "Experim. Unters. über Diphtherie," *Deutsches Archiv f. klin. Med.* Bd. viii. 1871.

the diphtheritic membranes of the organs of the throat and in their neighbourhood as well as in the surrounding lymphatics, but also in the blood of the general circulation, in the kidney, and in the muscles. The micrococci are about 0·00035—0·001 mm. in diameter, are slightly oval, occur singly or in dumb-




FIG. 20.—PORTION OF A DIPHTHERITIC MEMBRANE.
Numerous micrococci present.

bells or in short chains; they form also continuous masses of zoogloea in the shape of spherical or cylindrical clumps, and as such they penetrate and destroy the surrounding connective and muscular tissues. In severe cases they are found blocking up the capillaries of the glomeruli and the uriniferous tubules of the kidney.

Besides micrococci there occur in the diphtheritic membranes also other (rod-shaped) bacteria, but these are evidently only accessory.¹ Cultivations and inoculations with pure cultivations of this micrococcus are still wanting.

4. *Micrococcus pneumoniae*.—In acute croupous pneumonia there occur in the affected lungs large numbers of micrococci; Klebs, Eberth, Koch, Leyden, and others have seen them, but Friedländer² first pointed out their constant occurrence. According to this observer they are oval, of an oval nail-like shape, about 0·001 mm. long, and occur in the sputum singly, but especially as dumb-bells or diplococci, as chains, and as zoogloea. Ziehl³ found them in very large crowds in the sputum, giving to this in the early stages the peculiar characteristic brownish “prune-juice” tint. But this statement cannot be correct, since this tint may be very pronounced although the sputum contains only a limited number of the micrococci. According to this observer they are very numerous only in the beginning of the illness; after the critical stage they decrease in numbers.

¹ Compare also Klebs, *Archiv f. exp. Path.* iv.; Letzerich, *Virchow's Archiv*, vol. 68; Nassiloff, *ib.* vol. 50; Eberth, *Zur Kenntniss d. bact. Mycosen*, 1872; Wood and Formad, *Rep. of Nat. B. of Health, U.S.A.* 1882.

² *Virchow's Archiv*, vol. 87.

³ *Centralb. f. med. Wiss.* No. 25, 1883.

Griffini and Cambria saw the micrococci also in the blood. Salviali found that their number increases after the third day; on the ninth or tenth day they quite disappear.

G. Giles¹ found them in many cases of pneumonia (in India), both in the sputum and in the blood. Cultivations on boiled potato yielded good crops. These cultivated micrococci injected into the subcutaneous tissue of rabbits produced pneumonia.

Salviali and Zäslein² cultivated the micrococci (derived from the blood) in meat broth, meat extract solution, &c., at 37°-39° C., and obtained good crops of them, with which they produced by inoculation in seven rabbits and six white rats, typical pneumonia yielding the characteristic micrococci. These micrococci stain best in a mixture of Bismarck brown and methyl violet, but they stain also very well in gentian violet.

Quite recently Friedländer and Frobenius³ cultivated the micrococci in gelatine mixtures, and obtained good crops. The micrococci were of the peculiar nail-like shape, and were characterised by a mucinous capsule. Inoculation with the cultivated micrococci into the lungs of dogs was followed occasionally by positive results; in rabbits no result was obtained, and in mice lobar croupous pneumonia and pleurisy invariably followed the inoculation twenty-four to forty-eight hours afterwards. In guinea-pigs the results were not so decisive. About half of the animals escaped, the others died with dyspnoea, the blood, lungs, and pleural exudations containing the same micrococci. From my own observations I cannot accept these statements without qualification, for I find: That even in typical cases of croupous pneumonia of man, the micrococci may be absent or may be only very scarce even between the third and ninth day; that typical sputum of croupous pneumonia does not in many instances produce disease in animals on inoculation; and that the disease produced in rabbits and mice is of the nature of septicæmia, due to a specific septicæmic micrococcus not necessarily always present in the sputum and lungs of human croupous pneumonia.

If the fluids containing these micrococci were heated to about 70° C. they became inefficacious; mice inoculated with them remained healthy. Such inefficient micrococci (*i.e.* first heated to about 70° C.) did not grow any more on gelatine. Friedländer and Frobenius found also that when mice, shut

¹ *Brit. Med. Journal*, July 7, 1883.

² *Centralb. f. med. Wiss.* No. 41, 1883.

³ *Berichte d. physiolog. Gesellschaft in Berlin*, Nov 9, 1883.

up in a chest, were compelled to breathe an atmosphere saturated by means of a spray with the micrococcus pneumoniae, a number of them died from pneumonia and pleurisy, but not till the fourth or fifth day.



FIG. 21.—FROM A SECTION THROUGH THE HUMAN LUNG IN ACUTE CROUPOUS PNEUMONIA.

Three capillary vessels filled with blood are seen surrounding an alveolus which is filled with fibrin and blood-corpuscles, amongst them chains of micrococci.

Magnifying power 700. (Stained with gentian violet.)

Bruylants and Verriers¹ assert that they have successfully cultivated the micrococci of pleuro-pneumonia of cattle.

More recently² T. Poels and Dr. W. Nolen, in Rotterdam, assert that they have ascertained that in pleuro-pneumonia of cattle the pulmonary exudations contain micrococci, which in their morphology and mode of growth in artificial cultures

¹ *Bull. de l'Acad. Belg.* 1880.

² *Centraib. f. d. med. Wiss.* No. 9, 1884.

are identical with the micrococci of human pneumonia. And they further assert that artificial cultures of the micrococci




FIG. 22.—FROM A PREPARATION OF BLOOD OF RABBIT DEAD AFTER INOCULATION WITH SPUTUM OF ACUTE CROUPOUS PNEUMONIA.

Weigert-Koch method, stained with gentian violet. Numbers of blood discs, between them oval micrococci, surrounded by hyaline capsules. Magnifying power 700.

derived either from human pneumonia or from pleuro-pneumonia of cattle, produce in cattle the typical pleuro-pneumonia.




FIG. 23.—FROM A PREPARATION OF PLEURAL EXUDATION OF A MOUSE DEAD AFTER INOCULATION WITH BLOOD OF RABBIT MENTIONED IN PRECEDING FIGURE.

Magnifying power 700.

From my own observations I have reason to doubt the accuracy of these statements.

5. *Micrococcus gonorrhœæ*.—Micrococci have been found in the pus of gonorrhœa. Neisser,¹ and later Bokai and Finkelshtein,² described them as spherical organisms of about 0.008 mm. diameter, generally forming dumb-bells, or sarcina-like colonies of four. Several such groups form a zooglœa. They adhere to the pus-corpuscles and epithelial cells. They stain easily and well in methyl violet and gentian violet. Bockhart³ has succeeded in artificially cultivating these micrococci, and in producing the disease by inoculation with the cultivated organisms.



FIG. 24.—TWO LARGE SCALY EPITHELIAL CELLS OF GONORRHOËAL PUS. The epithelial cells are covered with micrococci, chiefly in dumb-bells, some in sarcina form.

Aufrecht⁴ reports the case of an infant twelve days old who died with suppuration of the umbilical vein and liver. The liver cells and the interlobular tissue were crowded with micrococci (shown in sections by means of a 2 per cent. watery solution of Bismarck brown). These micrococci corresponded in size to the micrococcus gonorrhœa, and he thinks it probable that they were derived from the vagina of the mother; during birth they might have got into the umbilical vein, there caused inflammation, and thence passed into the liver.

6. *Micrococcus endocarditicus*.—Micrococci in the form of zooglœa have been seen in *endocarditis ulcerosa*. They sometimes form plugs in the blood-vessels of the muscular tissue of

¹ *Centralb. f. d. med. Wiss.* No. 28, 1879.

² *Prager med. chir. Presse* May 1880.

³ *Sitzungsberichte der phys.-med. Gesellsch. in Würzburg*, Sept. 1882.

⁴ *Centralb. f. d. med. Wiss.* No. 16, 1883.

the heart (Heiberg,¹ Maier,² Eberth,³ Köster,⁴ Klebs⁵). Heiberg saw the micrococci forming chains in the muscle of the heart, in the detritus of the ulcerations of the endocardium, in the plugs in the vessels of the spleen and kidney.

7. *Micrococcus scarlatinae*.—In *scarlatina* Coze and Feltz⁶ described micrococci as occurring in the blood; as I have mentioned above, I have seen them in the ulcerations of the throat,⁷ and quite recently Pohl-Pincus⁸ described very minute micrococci adhering to the scales of the desquamating epidermis in *scarlatina*. They form small colonies, and stain violet with a saturated solution of methyl violet. Their diameter is very small, only about 0.0005 mm. The same micrococci were noticed by Pohl-Pincus in the throat-discharge.⁹

8. In the so-called *catle plague* (or rinderpest) micrococci have been found in the lymphatic glands by Klebs (1872) and by Semmer in the blood and lymphatic glands (1874 and 1881). In conjunction with Archangelski,¹⁰ Semmer cultivated the micrococci, obtained from the lymphatic glands of a sheep dead of inoculated rinderpest, in beef broth, in meat-extract solution, and in mixture of broth, peptone, and gelatine at 37°-39° C. The micrococci grew very copiously as zooglœa and in chains. With these micrococci (of a first transfer or cultivation) a calf was inoculated, and died after seven days from rinderpest. The cultures when transferred lose gradually their virulence from one generation to the next, but animals (sheep) inoculated with these are protected against further virulent disease. Further, cultures exposed for an hour to a temperature of 46°-47° C. become greatly attenuated in their action, and sheep inoculated with virus thus attenuated are protected against virulent material. Temperatures of -10° to -20° annihilate the activity of rinderpest organisms. The specific nature of these micrococci of rinderpest cannot, however, be considered at all established as in the case of those mentioned above, e.g. micrococci of erysipelas and gonorrhœa.

9. In *puerperal fever* micrococci have been found in the form of zooglœa by Heiberg,¹¹ in all affected organs—endocardium,

¹ Virchow's Archiv, vol 56.

³ Ibid. vol. 57.

⁵ Archiv f. exp. Path. Bd 9

⁷ Report of the Medical Officer of the Privy Council for 1876.

⁸ Centralb. f. d. med. Wiss. No. 36, 1883.

⁹ Seen already by McKendrick, British Med Journal, 1872.

¹⁰ Centralb. f. d. med. Wiss. No. 18, 1883.

² Ibid vol. 62.

⁴ Ibid vol 72.

⁶ Malad. infect. 1872.

¹¹ Leipzig, 1872.

lung, spleen, cornea, in a case of panophthalmitis puerperalis, and in the kidney, forming casts in the uriniferous tubules and emboli in the blood-vessels. Laffler¹ found zooglœa and chains of micrococci in two cases of puerperal fever associated with brain-softening. In both cases emboli, due to micrococci, were found in the surroundings of the softened part of the brain. Emboli of micrococci were also here found in the vessels of the kidney.

10. In *pernicious anæmia* Frankenhäuser² described the occurrence of micrococci (?) in the blood of pregnant women suffering from this anæmia, not uncommon in Zürich. These micrococci were very large, about one-tenth of the broad diameter of a red blood-corpuscle, and some were provided with a flagellum (?). Some were divided in two. In the blood of the liver they occurred in large numbers. Frankenhäuser's description makes it very difficult exactly to understand what he saw. He also states that these micrococci were probably derived from decayed teeth, from which all his patients suffered.

Eppinger³ described micrococci as occurring in *acute yellow atrophy* of the liver.

11. In the *syphilitic mucous patches* of several patients Aufrecht found a micrococcus, forming generally dumb-bells and staining very deeply in fuchsin.⁴ Birch-Hirschfeld⁵ confirmed this.

12. *Micrococcus of acute infectious osteomyelitis*.—Dr. Becker has made, in the laboratory of the Berlin Imperial Sanitary Office, a series of important experiments on the micro-organisms discovered by Schüller and Rosenbach. He collected pus from five cases of acute osteomyelitis in which the abscesses had not been opened, and cultivated the micrococci contained in it on sterilised potatoes, coagulated serum, and gelatine-peptone. In the latter case, the pus was introduced by means of needles into the mass, which was then kept at the temperature of the room during three to five days. After that time, the puncture made by the needles assumed the appearance of white streaks, around which the gelatine liquefied gradually and took an orange-colour. After a few days more, the mass gave out a smell like sour paste, and the microscope revealed the presence

¹ *Breslauer ärstl. Zeitschrift*. 1880.

² *Centralb. f. d. med. Wiss.* No. 4, 1888.

⁴ *Centralblatt f. d. med. Wiss.* No. 13, 1891.

³ *Prager Viertelj.* 1875.

⁵ *Ibid.* No. 44, 1882.

of large numbers of micrococci, having the same appearance as those found in the pus. A small quantity of the mass was mixed with sterilised water and injected into the peritoneal cavity of some animals; they died in a very short time of acute peritonitis. The same fluid injected into the jugular vein caused acute septicæmia and death; but nothing abnormal was found in the bones in either case. Dr. Becker then injected a small quantity of the same fluid into the jugular vein of fifteen rabbits, after having, some days before, fractured or bruised the bone of one of the hind legs. On the day after the injection, weakness and loss of appetite were noticed; but after a short time the symptoms passed away, and the animals seemed to have recovered. At the end of the first week, however, a swelling formed at the seat of the bruise or fracture, the animals lost flesh, and died after a few days. On dissection, large abscesses were found around and in the bones, and in several cases metastatic abscesses had formed in the lungs and kidneys. Numerous colonies of micrococci were discovered in the blood and pus of the animals upon which the experiments were made. (*Brit. Med. Journal*, March 29, 1884.)

13. Koch¹ described various kinds of micrococci intimately connected with certain destructive (pyæmic) processes in mice and rabbits. (a) *Micrococcus of progressive necrosis* in mice. Injecting into the ear of mice—white mice, or better, field mice—putrid fluids, he observed a necrosis of the tissues of the ear (skin, cartilage) starting from the point of inoculation and gradually spreading on to the surrounding parts and killing the animal in about three days. As far as the necrosis reaches, the tissue is crowded with micrococci, chiefly in the form of chains and zooglæa. The individual cells are spherical, of about 0·0005 mm. in diameter. I may mention that I have found a somewhat different micrococcus virulently active on mice. I have inoculated a number of white mice subcutaneously in the tail with a small micrococcus cultivated through several generations, and apparently derived from an artificial cultivation in pork broth, but due to accidental contamination. These micrococci, after having been cultivated in pork broth through several generations, were used in infinitesimal doses for the inoculation of the above mice. In two instances I have seen that the inoculation was followed after two to three days by purulent inflammation at the seat of inoculation, but apparently not spreading beyond it. But as time went on

¹ *Untersuchungen über die Aetiologie d. Wundinfections-Krankheiten*, Leipzig, 1878.

inflammation and abscess in the lungs set in and the animals died after about a week. On making longitudinal sections through the tail, it was found that in most of the lymph-spaces and lymph-vessels of all parts of the cutis and subcutaneous tissue, far away from the seat of inflammation, there were dense crowded masses of the same minute micrococci as were used for inoculation. And these crowds of micrococci could be traced to the seat of inflammation, where they extended amongst the inflammatory products in great masses. The abscesses in the lungs were filled with the same micrococci. Inoculated into the skin of fresh mice, it again produced death by pyæmia. This micrococcus may therefore be called the *micrococcus pyæmiæ* of mice. (b) Micrococcus causing abscesses in rabbits. Putrid blood injected into the




FIG 25.—FROM A SECTION THROUGH THE TAIL OF A MOUSE INOCULATED INTO THE SUBCUTANEOUS TISSUE OF THE TAIL WITH ARTIFICIALLY CULTIVATED MICROCOCCUS.

The part here illustrated is a good distance from the ulceration.

1. A capillary blood-vessel filled with blood-corpuscles.
2. Fat cells.
3. Groups of micrococci filling the lymph-spaces of the connective tissue.

subcutaneous tissue of the rabbit often produces suppurative abscess which, spreading, kills the animal in about twelve days. In the wall of the abscess are found continuous masses of zoogloea of micrococci. The pus is infectious. The micrococci are spherical, and of a very minute size, measuring only about 0.00015 mm. in diameter. (c) Micrococcus causing pyæmia in rabbits. Skin of a mouse was macerated in distilled water for two days, and of this fluid a hypodermic syringe-full was injected under the skin of the back of a rabbit. After two days the animal began to lose flesh and died after 105 hours.

Purulent infiltration spread from the seat of inoculation into the subcutaneous tissue; peritonitis; spleen much enlarged; slight pneumonia. A hypodermic syringe-full of the blood of this animal was injected under the skin of a second rabbit, and this died after forty hours. Post-mortem examination




FIG. 26.—FROM A PYOGENIC MEMBRANE COVERING THE SEROUS COAT OF THE INTESTINE OF A RABBIT DEAD OF PYÆMIA.

1. A large oval nucleus, probably the nucleus of a detached endothelial cell.
2. A pus corpuscle.

The rest of the pyogenic membrane is beset with small micrococci.




FIG. 27.—PYÆMIA OF RABBIT.

Blood of spleen. Between red blood-discs three dumb-bells and two single micrococci are shown. (Gentian violet staining.)

[The micrococci as here represented are somewhat too large.]

showed the same lesions as in the first case. In the blood-vessels of the affected parts were present micrococci, single, as dumb-bells, and in zooglæa; they were spherical, about 0.00025 mm. in diameter. (*d*) Micrococcus causing *septicæmia* in rabbits. An infusion of meat was prepared; this was left to putrefy, and of this fluid a quantity was injected under the skin of the back in two cases. Extensive gangrene with much



FIG. 28.—OVAL MICROCOCCI FROM THE BLOOD-VESSEL OF THE SPLEEN OF A RABBIT, DEAD OF KOCH'S SEPTICÆMIA.

œdematous exudation followed, and death ensued in two days and a half. The blood, the capillaries of the kidney, and the enlarged spleen, contained numerous oval micrococci. Two drops of the œdematous exudation-fluid were injected under the skin of the back of another rabbit. Death followed in twenty-two hours. There was no gangrene here; but œdema

was present, spreading from the seat of the inoculation. Sub-serous hæmorrhages appeared in the intestines; and minute hæmorrhages were also present in the œdematous tissue and in the muscles of the thigh and abdomen. The œdematous fluid, the cutaneous veins, the capillaries in the kidney, especially those of the glomeruli, in the lung, and in the spleen, contained numerous oval micrococci, singly, as dumb-bells, and in zooglœa. The micrococci measured about 0·0008 to 0·001 mm. in their long diameter. These micrococci (taken with the blood) produced in another rabbit and in a mouse the same fatal disease.

14. *Micrococcus bombycis* (*Microzyma bombycis*, Béchamp).—Oval micrococci, of about 0·0015 mm. in length, present in large numbers, singly, and as dumb-bells and chains (straight or curved), in the contents of the alimentary canal and in the gastric fluid of silkworms dead of the "maladie de morts-blancs, *flacherie*."—*Micrococcus ovatus*, *Nosema bombycis*. Present in large numbers in the blood and organs, ova included, of silkworms affected with the disease called "maladie des corpuscles," "pébrine," or *Cornalia's* disease. *Cornalia* first saw them, afterwards Lebert and Nägeli. Pasteur proved definitely that ingestion as well as inoculation of the silkworms with the micrococci produces the disease. The micrococci are comparatively large, 0·003 to 0·004 mm. long, 0·002 mm. broad; they are very bright and occur singly, or in dumb-bells, or in small groups.

CHAPTER VIII.

BACTERIUM (*Microbacterium*, Cohn).

By this name Cohn¹ designates a class of minute schizomycetes, which are slightly elongated and oval, or short and cylindrical, with rounded ends. They divide by fission, like the micrococci, the individuals elongating and becoming constricted in the middle. They are capable of spontaneous locomotion, being possessed of a flagellum at one or both ends, with which they perform active spinning and darting movements (Dallinger). Engelmann has shown that these movements are only possible in the presence of oxygen. Bacteria are found also as dumb-bells, *i.e.* in the act of dividing, and then appear as rods constricted in the middle. Occasionally, after rapid division, several remain connected, thereby forming a short chain. In this state the terminal elements are flagellate. Bacteria, like micrococci, are capable of forming zooglœa, the interstitial gelatinous substance being, as a rule, more copious than in the zooglœa of micrococci. In this state they form pellicles, in which the elements are without flagella; but from the margin of the pellicles one constantly sees elements separating, becoming flagellate, and moving away. In some species the zooglœa is dendritically ramified (*Zooglœa ramigera*, Itzigsohn), as seen on the surface of fluids containing decomposing algæ.

1. *Septic bacteria*.—With Cohn we distinguish two kinds:—*Bacterium termo* and *Bacterium lineola*.

(a) *Bacterium termo*.—The elements are short and cylindrical, about 0.0015 mm. long, a third less in breadth, and appear generally as dumb-bells. They are common in putrefying

¹ *Biologie d. Pflanzen*, ii. (1872), p. 167.

fluids, indeed they form the essential cause or ferment of decomposition, being the true saprogenous ferment (Cohn). They are invested in a thick membrane, and are flagellate. With the end of putrefaction they disappear. They grow well in Cohn's nourishing fluid, and I have found them as constant inhabitants of unfiltered distilled water in the laboratory; so much so that with a drop of this water I am always able to start a copious growth of bacterium termo in pork broth, Agar-Agar, &c. When cultivated in the incubator at 32° to 36° C. in



FIG. 29.—BACTERIUM TERMO FROM AN ARTIFICIAL CULTURE.



FIG. 30.—ZOOGLŒA OF BACTERIUM TERMO.

suitable nourishing material (pork broth, chicken broth,) they produce a uniform turbidity, and after several days an attempt at a pellicle, the whole nourishing fluid becoming thicker. But after from several days to a few weeks the cultures die, a fact which distinguishes them from all other bacteria. Growing in solid Agar-Agar, and peptone mixture, they produce an imperfect liquefaction, numerous gas bubbles appearing in the material.



FIG. 31.—BACTERIUM LINEOLA.

(b) *Bacterium lineola* (*Vibrio lineola*, Ehrenberg, Dujardin), differs from bacterium termo in being longer and thicker. The cells are about 0.003 to 0.005 mm. long, about 0.0015 mm. thick. They occur in well-water and stagnant water, where no distinct putrefaction is going on, and form zoogloea, and pellicles, on the surface of potatoes and various infusions.

2. *Zymogenic bacteria*.—Two kinds are known: *Bacterium lactis* and *Bacterium aceti*.

(a) *Bacterium lactis*.—According to Pasteur, these bacteria

about 0.0015 to 0.003 mm. long, constricted in the centre; they form short chains, or even zoogloea, and they are motile. They produce the lactic acid fermentation, in the course of which lactic sugar is transformed into lactic acid; they are



FIG. 32.—BACTERIUM LACTIS.

anaërobic. Lister,¹ by means of pure cultures, established experimentally their causal relation to the lactic fermentation or souring of milk.

(c) *Bacterium acetii* (*Mycoderma acetii*) is a little smaller than *Bacterium lactis*, being about 0.0015 mm. in length, and often forms chains, and also pellicles, on the surface of the fluid; it is motile. Pasteur maintains that it is the ferment of the acetic acid fermentation. Cohn² found it in enormous masses in beer that had become sour; it forms dumb-bells, seldom chains of four, and sometimes a pellicle on the surface. Pure cultivations have not been made with it, and before deciding whether it is the real cause of the acetic acid fermentation, experiments with such pure cultures, i.e. inoculations of alcoholic fluids with it, are required.

3. *Pigment bacteria*.—Two kinds have been described: *Bacterium xanthinum* and *Bacterium aeruginosum*.

(a) *Bacterium xanthinum*³ is a bacterium, about 0.007 to 0.01 mm. long, motile, single, also in dumb-bells, or short chains. It produces the yellow colour of yellow milk. Its pigment is soluble in water, and insoluble in alcohol or ether. When introduced into boiled milk of neutral reaction, it multiplies with great rapidity; the milk coagulates after twenty-four hours; it is soon teeming with them and turns yellow. The reaction of the yellow milk is at first acid, but soon becomes alkaline, and the alkalinity gradually increases.

(b) *Bacterium aeruginosum*.—In green pus Schroeter discovered a bacterium, *Bacterium aeruginosum*.⁴ The pigment

¹ *Pathological Soc. Transactions*, 1878.

² *Biol. d. Pflanzen*, ii. p. 173.

³ Schroeter, *Biol. d. Pflanzen*, ii. p. 120; *Vibrio synxanthus*, Ehrenberg.

⁴ *Loc. cit.* p. 122.

is greenish, and not lodged in the cells themselves ; it is easily diffusible.

4. *Pathogenic bacteria*.—Three kinds are described ; the bacteria of Koch's septicæmia, of Davaine's septicæmia, and of fowl-cholera.

(a) *Bacterium septicæmiæ* (Koch).—By injecting into rabbits water from the rivulet Pauke, and from putrid mutton, Koch¹ succeeded in producing a rapidly fatal septicæmia, which was characterised by the following appearances :—The blood of all the organs contained very numerous bacteria, the spleen and lymphatic glands were enlarged, and the lungs congested ; but there were no extravasations and no peritonitis. The smallest quantity of this blood inoculated into the skin or cornea of another rabbit produced after an incubation of ten to twelve hours distinct rise of temperature, and death after sixteen to twenty hours. The conditions after death were the same as above. Everywhere the blood contained the bacteria. They



FIG. 33.—BLOOD OF PIGEON, DEAD OF SEPTICÆMIA.
Four blood-discs and four bacteria-termo are shown.

are rods somewhat pointed at both ends, measuring about 0·0014 mm. in length and 0·0006 mm. in breadth. When stained, they show at each end a deeply-tinted granule, the middle part remaining unstained ; for this reason they are easily mistaken for a diplococcus. Generally these rods occur singly ; occasionally they form a chain of two, or more than three.

They have been cultivated successfully in beef broth, blood serum, gelatine, and a mixture of gelatine and broth and peptone. The cultures have the same virulent properties as the original blood.

Mice, pigeons, fowls, and sparrows are also very susceptible to these bacteria ; but guinea-pigs, dogs, and rats resist them successfully.

The microbe found by Pasteur in human saliva, which he

¹ *Mitth. aus d. k. Gesundh.* 1831.

cultivated, and with which he produced septicæmia in rabbits, may perhaps be a bacterium identical with the above, but this is not definitely settled.

(b) *Bacterium of Davaine's septicæmia*.—This is a bacterium which was originally derived by Davaine¹ from putrid ox-blood in the warm season. Injected into rabbits, it produced rapidly fatal septicæmia, of the same nature as in the case just mentioned, the blood teeming with a similar kind of bacterium as in Koch's septicæmia just described. The smallest quantity of the blood is again rapidly fatal in its action. It is distinguished from Koch's septicæmia in the rabbit by this, that Davaine's septicæmia is easily transmissible to guinea-pigs, but not to birds.

Dowdeswell² has shown that when such blood is thoroughly sterilised (i.e. when the bacteria are killed), it has no longer any infective power. Davaine had first shown that the blood of rabbits dead of this form of septicæmia bears an enormous



FIG. 31.—BLOOD OF RABBIT, DEAD OF DAVAINÉ'S SEPTICÆMIA.

amount of dilution without the minutest quantity of it losing its pathogenic properties. Dowdeswell has shown that this is easily explained by the enormous number of bacteria present in every drop of the blood. But it has been shown by Gaffky and Dowdeswell that there is no increase in the virulence of the virus when it is passed through successive animals, as was maintained by Coze and Feltz.³

(c) *Bacterium of fowl-cholera (microbe du choléra des poules)*.—Semmer, Toussaint, and Pasteur⁴ have shown that this

¹ *Bull. d l'Acad. de Méd.* 1872.

² *Proceedings of the Royal Society*, No. 221, 1882.

³ *Strasbourg*, 1866; *Paris*, 1872.

⁴ I place this here as a bacterium, but it is not quite decided, and not quite clear from Pasteur's description, whether the microbe is only a micrococcus dumb bell, or a bacterium termo. Compare also Semmer (*Vergleichende Pathologie*, 1876), Perroncito (*Archiv f. Wiss. u. pract. Thierheilk.*, 1879). Toussaint (*Comptes Rendus*, xci p. 301) considers the disease identical with Davaine's septicæmia. I am inclined to think that Pasteur has not used pure cultivations,

organism is present in large numbers in the blood and organs of fowls dead of this malady, which is chiefly characterised by the following symptoms:—The animals are somnolent, weak in their legs and wings, and they die under symptoms of extreme sopor. On post-mortem examination, hæmorrhage is found in the duodenum. The smallest quantity of the blood is infective. Pasteur successfully cultivated the bacteria in neutral chicken broth, at 25° to 35° C., and with it inoculated the fatal disease. The organism is probably a bacterium termo, very minute and slightly constricted in the middle, so that it appears of the shape of an 8. When cultures of this bacterium¹ are kept for some time (one, two, three or more months), their virulence becomes diminished or attenuated (owing, according to Pasteur, to the action of oxygen), and this diminution of virulence is in direct proportion to the time the culture is kept. The diminution or attenuation shows itself in this—that according to the length of time the culture is kept, the number of animals killed by its inoculation gradually diminishes, and it ultimately ceases to kill at all. Each culture of diminished virulence transmits its attenuation to the next following culture (?). It is possible to obtain cultures of such a low degree of virulence that when inoculated into the skin of a fowl only a local effect is produced, a peculiar infiltration; but the animal survives, and is then protected or “vaccinated” against the more virulent material. But in order to produce this protection, it is necessary that the culture (*vaccine*) should be of the proper strength. If it does not produce a local effect it gives no protection.

In fresh cultures the bacterium is more in the shape of a rod, constricted in the middle; in cultures several days old it looks very much more like a dumb-bell of micrococcus (see note on previous page).

Babes² has found the bacteria in the tissues and blood-vessels of animals dead of the disease, both inoculated and epizootic, in the shape of rods of about 0·0015 to 0·002 mm. in length and about 0·00025 mm. in thickness; the ends always staining more deeply than the middle part.

but had the bacterium of fowl-cholera and an accidental micrococcus together. The latter would predominate as time passed on, so that after a few days it would far outnumber the bacterium; and this is exactly what Pasteur's description suggests. He says that at first the microbe is rod-shaped, and after a few days it becomes a dumb-bell of micrococcus. The gradual attenuation by time of the virulence of Pasteur's cultures of the microbe of fowl-cholera may be due to the presence of this contaminating micrococcus.

¹ *Trans. of the International Med. Congress in London, 1881, vol. i. p. 87.*

² *Archives de Physiologie, July 1863, p. 49.*

Purulent infiltration spread from the seat of inoculation into the subcutaneous tissue; peritonitis; spleen much enlarged; slight pneumonia. A hypodermic syringe-full of the blood of this animal was injected under the skin of a second rabbit, and this died after forty hours. Post-mortem examination




FIG. 26.—FROM A PYOGENIC MEMBRANE COVERING THE SEROUS COAT OF THE INTESTINE OF A RABBIT DEAD OF PYÆMIA.

1. A large oval nucleus, probably the nucleus of a detached endothelial cell.
2. A pus corpuscle.

The rest of the pyogenic membrane is beset with small micrococci.




FIG. 27.—PYÆMIA OF RABBIT.

Blood of spleen. Between red blood-discs three dumb-bells and two single micrococci are shown. (Gentian violet staining.)

[The micrococci as here represented are somewhat too large.]

showed the same lesions as in the first case. In the blood-vessels of the affected parts were present micrococci, single, as dumb-bells, and in zooglæa; they were spherical, about 0.00025 mm. in diameter. (d) *Micrococcus* causing *septicæmia* in rabbits. An infusion of meat was prepared; this was left to putrefy, and of this fluid a quantity was injected under the skin of the back in two cases. Extensive gangrene with much



FIG. 28.—OVAL MICROCOCCI FROM THE BLOOD-VESSEL OF THE SPLEEN OF A RABBIT, DEAD OF KOCH'S SEPTICÆMIA.

œdematous exudation followed, and death ensued in two days and a half. The blood, the capillaries of the kidney, and the enlarged spleen, contained numerous oval micrococci. Two drops of the œdematous exudation-fluid were injected under the skin of the back of another rabbit. Death followed in twenty-two hours. There was no gangrene here; but œdema

was present, spreading from the seat of the inoculation. Sub-serous hæmorrhages appeared in the intestines; and minute hæmorrhages were also present in the œdematous tissue and in the muscles of the thigh and abdomen. The œdematous fluid, the cutaneous veins, the capillaries in the kidney, especially those of the glomeruli, in the lung, and in the spleen, contained numerous oval micrococci, singly, as dumb-bells, and in zooglœa. The micrococci measured about 0·0008 to 0·001 mm. in their long diameter. These micrococci (taken with the blood) produced in another rabbit and in a mouse the same fatal disease.

14. *Micrococcus bombycis* (*Microzymba bombycis*, Béchamp).—Oval micrococci, of about 0·0015 mm. in length, present in large numbers, singly, and as dumb-bells and chains (straight or curved), in the contents of the alimentary canal and in the gastric fluid of silkworms dead of the “maladie de morts-blancs, *flacherie*.”—*Micrococcus ovatus*, *Nosema bombycis*. Present in large numbers in the blood and organs, ova included, of silkworms affected with the disease called “maladie des corpuscles,” “pébrine,” or *Cornalia*’s disease. *Cornalia* first saw them, afterwards Lebert and Nägeli. Pasteur proved definitely that ingestion as well as inoculation of the silkworms with the micrococci produces the disease. The micrococci are comparatively large, 0·003 to 0·004 mm. long, 0·002 mm. broad; they are very bright and occur singly, or in dumb-bells, or in small groups.

CHAPTER VIII.

BACTERIUM (*Microbacterium*, Cohn).

By this name Cohn¹ designates a class of minute schizomycetes, which are slightly elongated and oval, or short and cylindrical, with rounded ends. They divide by fission, like the micrococci, the individuals elongating and becoming constricted in the middle. They are capable of spontaneous locomotion, being possessed of a flagellum at one or both ends, with which they perform active spinning and darting movements (Dallinger). Engelmann has shown that these movements are only possible in the presence of oxygen. Bacteria are found also as dumb-bells, *i.e.* in the act of dividing, and then appear as rods constricted in the middle. Occasionally, after rapid division, several remain connected, thereby forming a short chain. In this state the terminal elements are flagellate. Bacteria, like micrococci, are capable of forming zooglæa, the interstitial gelatinous substance being, as a rule, more copious than in the zooglæa of micrococci. In this state they form pellicles, in which the elements are without flagella; but from the margin of the pellicles one constantly sees elements separating, becoming flagellate, and moving away. In some species the zooglæa is dendritically ramified (*Zooglæa ramigera*, Itzigsohn), as seen on the surface of fluids containing decomposing algæ.

1. *Septic bacteria*.—With Cohn we distinguish two kinds:—*Bacterium termo* and *Bacterium lineola*.

(a) *Bacterium termo*.—The elements are short and cylindrical, about 0.0015 mm. long, a third less in breadth, and appear generally as dumb-bells. They are common in putrefying

¹ *Biologie d. Pflansen*, II. (1872), p. 167.

fluids, indeed they form the essential cause or ferment of decomposition, being the true saprogenous ferment (Cohn). They are invested in a thick membrane, and are flagellate. With the end of putrefaction they disappear. They grow well in Cohn's nourishing fluid, and I have found them as constant inhabitants of unfiltered distilled water in the laboratory; so much so that with a drop of this water I am always able to start a copious growth of bacterium termo in pork broth, Agar-Agar, &c. When cultivated in the incubator at 32° to 36° C. in



FIG. 29.—BACTERIUM TERMO FROM AN ARTIFICIAL CULTURE.



FIG. 30.—ZOOGLŒA OF BACTERIUM TERMO.

suitable nourishing material (pork broth, chicken broth,) they produce a uniform turbidity, and after several days an attempt at a pellicle, the whole nourishing fluid becoming thicker. But after from several days to a few weeks the cultures die, a fact which distinguishes them from all other bacteria. Growing in solid Agar-Agar, and peptone mixture, they produce an imperfect liquefaction, numerous gas bubbles appearing in the material.



FIG. 31.—BACTERIUM LINEOLA.

(b) *Bacterium lineola* (*Vibrio lineola*, Ehrenberg, Dujardin), differs from bacterium termo in being longer and thicker. The cells are about 0.003 to 0.005 mm. long, about 0.0015 mm. thick. They occur in well-water and stagnant water, where no distinct putrefaction is going on, and form zoogloea, and pellicles, on the surface of potatoes and various infusions.

2. *Zymogenic bacteria*.—Two kinds are known: *Bacterium lactis* and *Bacterium aceti*.

(a) *Bacterium lactis*.—According to Pasteur, these bact

about 0.0015 to 0.003 mm. long, constricted in the centre; they form short chains, or even zoogloea, and they are motile. They produce the lactic acid fermentation, in the course of which lactic sugar is transformed into lactic acid; they are



FIG. 32.—BACTERIUM LACTIS.

anaërobic. Lister,¹ by means of pure cultures, established experimentally their causal relation to the lactic fermentation or souring of milk.

(c) *Bacterium acetii* (*Mycoderma acetii*) is a little smaller than *Bacterium lactis*, being about 0.0015 mm. in length, and often forms chains, and also pellicles, on the surface of the fluid; it is motile. Pasteur maintains that it is the ferment of the acetic acid fermentation. Cohn² found it in enormous masses in beer that had become sour; it forms dumb-bells, seldom chains of four, and sometimes a pellicle on the surface. Pure cultivations have not been made with it, and before deciding whether it is the real cause of the acetic acid fermentation, experiments with such pure cultures, i.e. inoculations of alcoholic fluids with it, are required.

3. *Pigment bacteria*.—Two kinds have been described: *Bacterium xanthinum* and *Bacterium aeruginosum*.

(a) *Bacterium xanthinum*³ is a bacterium, about 0.007 to 0.01 mm. long, motile, single, also in dumb-bells, or short chains. It produces the yellow colour of yellow milk. Its pigment is soluble in water, and insoluble in alcohol or ether. When introduced into boiled milk of neutral reaction, it multiplies with great rapidity; the milk coagulates after twenty-four hours; it is soon teeming with them and turns yellow. The reaction of the yellow milk is at first acid, but soon becomes alkaline, and the alkalinity gradually increases.

(b) *Bacterium aeruginosum*.—In green pus Schroeter discovered a bacterium, *Bacterium aeruginosum*.⁴ The pigment

¹ *Pathological Soc. Transactions*, 1878.

² *Biol. d. Pflanzen*, ii. p. 173.

³ Schroeter, *Biol. d. Pflanzen*, ii. p. 120; *Vibrio synxanthus*, Ehrenberg.

⁴ *Loc. cit.* p. 122.

is greenish, and not lodged in the cells themselves ; it is easily diffusible.

4. *Pathogenic bacteria*.—Three kinds are described ; the bacteria of Koch's septicæmia, of Davaine's septicæmia, and of fowl-cholera.

(a) *Bacterium septicæmiæ* (Koch).—By injecting into rabbits water from the rivulet Pauke, and from putrid mutton, Koch¹ succeeded in producing a rapidly fatal septicæmia, which was characterised by the following appearances :—The blood of all the organs contained very numerous bacteria, the spleen and lymphatic glands were enlarged, and the lungs congested ; but there were no extravasations and no peritonitis. The smallest quantity of this blood inoculated into the skin or cornea of another rabbit produced after an incubation of ten to twelve hours distinct rise of temperature, and death after sixteen to twenty hours. The conditions after death were the same as above. Everywhere the blood contained the bacteria. They



FIG. 33.—BLOOD OF PIGEON, DEAD OF SEPTICÆMIA.
Four blood-discs and four bacteria-termo are shown.

are rods somewhat pointed at both ends, measuring about 0·0014 mm. in length and 0·0006 mm. in breadth. When stained, they show at each end a deeply-tinted granule, the middle part remaining unstained ; for this reason they are easily mistaken for a diplococcus. Generally these rods occur singly ; occasionally they form a chain of two, or more than three.

They have been cultivated successfully in beef broth, blood serum, gelatine, and a mixture of gelatine and broth and peptone. The cultures have the same virulent properties as the original blood.

Mice, pigeons, fowls, and sparrows are also very susceptible to these bacteria ; but guinea-pigs, dogs, and rats resist them successfully.

The microbe found by Pasteur in human saliva, which he

¹ *Mitth. aus d. k. Gesundh.* 1851.

described in a former chapter and kept at the ordinary temperature of the room or in the incubator at not more than 22° to 25° C., the spore-formation on the surface occurs only as long as the material remains solid. Anthrax-bacillus as it grows liquefies the gelatine mixture; in consequence of this after some days the superficial layers of the material become fluid, and the bacillar growth, sinking to the bottom of the fluid layer, is thus removed from the surface. The spores which were freely formed while the growth went on on the surface, germinate again into bacilli, but because these have now sunk into the depth, although rapidly multiplying and growing into filaments, they cease to form any spores.¹

Bacilli which are not possessed of the power of locomotion (*i.e.* are without a flagellum) when sown into the depth of a fluid or solid material, if they have no chance, accidental or otherwise, of reaching the surface, do not as a rule form spores; but there are some such bacilli which, although not growing on a free surface, nevertheless form spores, *e.g.* the bacillus butyricus or amylobacter (Prazmowski). Some putrefactive bacilli occurring after death in the abdominal organs (intestine, kidney, spleen, liver), and in fluid exudations within the peritoneal and pleural cavities, show also spore-formation; probably they get their oxygen from the tissues. Anthrax-bacillus, however, never forms spores except it is growing well exposed to the outer air.

The bacilli which are possessed of a flagellum (*e.g.* hay-bacillus, bacillus of putrefaction) generally form a pellicle on the surface, and in this pellicle copious spore-formation goes on.

The spores first formed, when shaken down into the fluid, again germinate into bacilli, and there multiply. The last pellicle formed in a culture, before exhaustion, represents the last crop of spores; and these, owing to the exhaustion of the nourishing fluid, remain as spores, only capable of germinating into bacilli when new nourishing material is added, or when they are transplanted to new nourishing material.

It is a rule that wherever the spores are formed they germinate into bacilli if they have access to nourishing material; but if not, or if the nourishing material is exhausted, they remain as spores. Spore-formation does not take place at low temperatures. Koch found in the case of anthrax-bacillus that a temperature below 12° C. prevents the formation of spores. Pasteur states that in the case of anthrax-bacillus spore-formation does not take place above 40° C.; never for

¹ Report of the Medical Officer of the Local Government Board, 1881.

instance at 42° or 43° C. Koch gives 43° as the upper limit ; but I have found that both in the case of hay-bacillus and anthrax-bacillus the bacilli form spores copiously even at a temperature of 44° C. Moisture is an essential element in the formation of spores.

The spores represent the seeds capable of retaining life and of germinating into bacilli even after what would appear the most damaging influences (that is, damaging to all other kinds of organisms and to the bacilli themselves), such as long lapse of time, drying, heat, cold, chemical reagents, &c. Spores retain the power to germinate into bacilli after the lapse of long periods, and there is no reason to assume that these periods have any limit ; it makes no difference whether they are kept dry or in the mother-liquid.

The temperature of boiling water, while it kills micrococci, bacteria, and bacilli themselves, does not affect the vitality of the spores. Cohn (*loc. cit.*) found spores of hay-bacillus still capable of germination even after boiling ; boiling for half an hour or more killed them. Prazmowski found that the spores of bacillus butyricus (*amylobacter*) are killed by five minutes' boiling. In the case of anthrax-bacillus and hay-bacillus I found that boiling for half an hour does invariably kill them, but ten minutes is not to be relied on. Exposing the spores of anthrax-bacillus to a temperature of 0° to -15° C. for one hour did not kill them. Antiseptics, such as carbolic acid (5-10 per cent.), strong solutions of phenyl-propionic acid and phenyl-acetic acid, corrosive sublimate (1 : 300,000, Koch), although the spores were kept in these fluids for twenty-four hours, did not kill them.

Pure terebene, phenol (10 per cent.), corrosive sublimate (1 per cent.), does not kill the spores of bacillus anthracis.

This great resistance of spores to low and high temperatures, to acids and other substances, is due to this, that the substance of each spore is enveloped in a double sheath : an internal sheath probably of a fatty nature, and an external one probably of cellulose ; both are very bad conductors of heat.

Owing to the fact that spores resist the action of boiling water, if not prolonged for ten minutes, and that the other bacteria (such as micrococcus, bacterium, and bacillus itself) are killed by the temperature of boiling water if kept at this temperature for a few seconds, it is possible to separate the spores of bacilli from the other organisms. All one has to do is to subject the fluid containing these various organisms to the temperature of boiling water for a few seconds. All except

the spores of bacilli will be thereby killed, and thus the fluid becomes free of all other organisms except the spores.

When spores are sown in a nourishing material, fluid or solid, and when this is exposed to a temperature of about 32° to 38° C., the spores after the lapse of a few hours, in some cases six (spores of anthrax-bacillus), in others two to four hours (spores of hay-bacillus), in others more than six hours, are seen to germinate, each spore growing into a bacillus. In the case of solid nourishing material the presence of moisture is essential.

In this germination what one sees is this : the spore increases in thickness, it then loses its dark contour at one pole or at one of the long sides, and at this point a pale projection appears. This projection increases in length and gradually becomes as long as a bacillus, the investment of the spore gradually fading away. This new bacillus soon divides into two, and so on.

The spores are capable of germinating independently of the free access of air.

CHAPTER X.

BACILLUS : NON-PATHOGENIC FORMS.

SEPTIC bacilli.

(a) *Bacillus subtilis* (hay-bacillus).—The elementary rods are of various lengths from 0·002 to 0·006 mm., and are about




FIG. 89.—FROM A CULTURE OF *BACILLUS SUBTILIS* (HAY-BACILLUS).
Various forms between single bacilli and leptothrix.
Magnifying power about 700.

0·002 mm. in thickness. According to Cohn¹ at a temperature of 21° C. division into two requires about one hour and a quarter, at 35° C. only about twenty minutes.

¹ *Loc. cit.*

The bacilli are capable of forming leptothrix filaments. The bacilli when single are possessed of one flagellum, or sometimes of two, one at each end. After division the individual bacilli remain connected, each possessing a flagellum at the free end. Each of them divides again into four, so that a chain of four is formed. But they may separate again or may go on dividing, remaining united, and thus forming a longer or shorter filament. Not all bacilli possess the flagellum, many of them being for a time in a resting state.

The bacilli form a dense resistant pellicle on the surface of the nourishing medium, and in this copious spore-formation




FIG. 49.—FROM A CULTURE OF *BACILLUS SUBTILIS* (HAY-BACILLUS), WITH COPIOUS FORMATION OF SPORES.

1. Mass of spores embedded in hyaline matrix.
2. Bacilli.
3. Single bacilli containing each a spore: the sheath of the bacilli is well seen. Magnifying power about 700.

takes place. If shaken when growing in a fluid, the pellicle falls to the bottom, and soon a new pellicle is formed.

Spore-formation is independent of any deficiency of nourishing material. The spores are oval, bright, of about 0.001 to 0.002 mm. in length, and about 0.0006 to 0.001 mm. in thickness. They do not stain in dyes, and hence form a great contrast to the bacilli.

This bacillus is very common and widely distributed; it occurs in almost every organic substance rich in nitrogenous compounds which is left exposed to the air to decompose. The best material is hay-infusion. An infusion, cold or hot, of hay is made in a beaker or flask; the fluid is filtered, covered with a glass plate, and left to stand in a warm place. After a day or two it swarms with bacillus subtilis, which is also called hay-bacillus, since ordinary hay contains multitudes of its

spores. For this reason even boiling of the fresh infusion for a few minutes does not sterilise it.

The bacillus grows well in every fluid that contains the necessary salts and nitrogenous compounds; thus all kinds of




FIG. 41.—GERMINATION OF SPORES INTO BACILLI.

a. Spores of a small kind.

b. Spores of a larger kind of bacillus subtilis.

Magnifying power about 700.

broth, all kinds of animal fluids (hydrocele, blood-serum, &c.), gelatine, peptone solution, &c., are suitable nourishing media.

The spores of the hay-bacillus are widely distributed in the air, and most contaminations by air are due to its spores.




FIG. 42.—BACILLUS ULNA, IN THE CAPILLARIES OF THE HUMAN LIVER.
POST-MORTEM CHANGE.

1. Liver cells, somewhat swollen.

2. Bacilli.

Magnifying power 300.

(b) *Bacillus ulna*.—By this name¹ Cohn designates certain species of bacilli, stiffer and thicker than those of bacillus

¹ *Loc. cit.* p. 177.

subtilis. The individual elements are about 0·01 mm. long, and 0·002 mm. thick. They are motile just like bacillus subtilis. Although they form chains they do not form proper leptothrix. They occur in putrid fluid. They are very common in the ichor produced by injecting ammonia or other substances producing sloughing and necrosis of the subcutaneous tissue in the guinea-pig.

(c) *Bacillus septicus* occurs in earth, in putrid blood, and in many putrid albuminous fluids. It is non-motile, and is capable of forming leptothrix. The thickness varies from 0·004 to 0·01 mm., and its length depends on the number of elements contained in a row. The shortest are about 0·004 mm. There are various species, differing from one another in the thickness of the elements. They are all anaërobic. The elements, whether in the short rods or in the leptothrix filaments, are cubical or rounded. The rods and filaments are markedly rounded on the ends. It forms spores independently of free access of air. The spores are oval, and differ in thickness according to the thickness of the bacilli they are formed in. The bacillus is found occasionally in the blood-vessels of man and animals after death. In a nourishing fluid, in which micrococcus, bacterium termo, or bacillus subtilis grows, they have no chance of growing, and even when numerous at first they soon disappear.

(d) *Streptothrix* and *Cladothrix*.—Cohn¹ found in a concretion of the human lacrimal canals long, pale, smooth, apparently branched threads, either straight or twisted; they were finer than the threads of leptothrix buccalis; he called them *Streptothrix Foersteri*. They are probably identical morphologically with *Cladothrix dichotoma*. This latter occurs in pond-water containing decomposing organic matter. It consists of long whitish threads fixed on chlorophyll-containing algæ. The threads when fresh appear smooth, pale, occasionally granular, and on staining they are seen to be composed of shorter or longer bacilli just like the leptothrix form of bacillus subtilis; but they are thicker than the bacillus subtilis. Occasionally the ends of the threads are seen not as linear series of bacillar rods, but like bacillus anthracis and the bacillus of blue milk (see below) as chains of torula-like spherical elements. From the threads single motile bacilli are seen to come off. The threads are only apparently branched, since the branches are threads merely stuck on to other threads sideways at an acute angle. A bacillus may be seen to stick to a thread and then to grow

¹ *Beitr. z. Biol. d. Pflanzen*, vol. i. p. 186.

out by continuous divisions into a long chain of bacilli, thus forming, as it were, a side-branch. Some of the threads are wavy and curved ; most of them are, however, straight. Zopf¹



FIG. 43.—STREPTOTHRIX FOERSTERI¹
(AFTER COHN).



FIG. 44.—CLADOTHRIX DICHOTOMA
(AFTER COHN).

claims to have observed that the threads of the cladothrix gave rise to micrococcus, bacterium, bacillus, and spirillum ; and states that each of these is again capable of growing into the threads of the cladothrix. But these observations were not made after exact methods.

(e) *Beggiatoa*.—In stagnant water, particularly in sulphur-containing water, peculiar oscillating colourless threads are met with of the thickness of 0.001 to 0.016 mm. ; they contain highly refractive granules, which Cohn (*Beiträge zur Biol. d. Pfl.* I. 3) has shown to be composed of sulphur. After dissolving these granules it is seen that each thread is septate, being composed of a sheath and transverse septa at regular intervals, by which the threads appear made up of a series of short cylindrical elements. There are a number of species varying from one another in the thickness of the threads.

¹ *Zur Morphologie der Spaltpflanzen*, Leipzig, 1882 ; see also Cienkowski.

Zymogenic bacilli.

Amongst these there is one species definitely known, namely, the *Bacillus butyricus* (*Bacillus amylobacter*,¹ *Clostridium butyricum*, *ferment butyrique*, Pasteur). This bacillus has the same




FIG. 45.—THREADS OF CLADOTRIX DICHOTOMA HIGHLY MAGNIFIED AND STAINED WITH SPILLER'S PURPLE.

1. Threads of bacilli.
2. Torula forms.

The sheath is everywhere well seen.

morphological characters as regards length and thickness of the rods, as regards power to form leptothrix, and as regards motility, as the bacillus subtilis. It is capable of forming

¹ Prazmowski, Leipzig, 1880; van Tieghem, *Bull. de la Soc. botanique*, vol. xxiv. 1877.

zooglycæ, and is anaërobic, since it grows well and forms spores copiously even when not exposed to the air. After the rods have gone on dividing and forming chains and filaments for



FIG. 46.—*CLOSTRIDIUM BUTYRICUM* OR *BACILLUS BUTYRICUS*.
Some of the spindle-shaped forms include an oval spore.

some time, they swell up, become granular and oval with more or less pointed ends, and the formation of oval spores sets in. In this state the oval rods are about 0·002 to 0·003 mm. thick, and the spores are about 0·002 to 0·003 mm. long and 0·001 mm. thick. In solutions of starch, dextrin, and sugar the bacillus forms butyric acid. The fermentation of butyric acid in old milk and ripening cheese is due to this bacillus. Cellulose is decomposed by it, and hence its great importance in the digestive process of herbivorous animals, in whose stomach and intestine it is very common. It is very common also in substances containing starch.

Iodine produces a characteristic blue staining in the protoplasm of the bacillus. In young rods the colour produced by iodine is blue, in older rods it is violet.

E. Kern described (*Biolog. Centralbl.* ii. p. 135) a bacillus under the name of *dispora caucasica*, which he found in the Caucasus, and which is used as ferment to produce from cow's milk a peculiar drink called "kephir" or "hpyppö." The bacillus is similar to the *bacillus subtilis*, but is distinguished from it and all other bacilli by this, that every bacillus forms two spores, one at each end, hence the name *dispora*. But after recent investigations it appears that this bacillus is accidental, the fermentation being produced by *saccharomyces mycoderma* (see Chapter XIV.).

Pigment bacilli.

(a) *Bacillus ruber*.¹—This appears as minute rods, isolated or

¹ Cohn, Frank, *Beitr. z. Biol. d. Pflanzen*, vol. iii. p. 181.

in twos and fours, and motile. It was found on boiled rice by Frank. Its colour is red, and contained in the bacilli themselves.

(b) *Bacillus erythrosporus*.—Motile isolated rods and leptothrix. It was found in meat extract solutions and on decomposing albumen, and forms pellicles. In the rods are found oval spores.¹

(c) *Bacillus syncyanus* (Neelsen) causes the blue colour of milk after the milk has become acid; it grows well in ammonium lactate. It consists of motile rods, single or in short chains. Neelsen² saw the bacilli assuming a torula-form, the individual cells being hourglass-shaped or oval, or even spherical (compare *Bacillus anthracis*, Chapter XI.). The rods appear also as a non-motile variety, and are then found to



FIG. 47. *BACILLUS SYNCYANUS* (AFTER NEELSEN).

1. Typical bacilli, motile.
2. Non-motile rods invested in a gelatinous capsule.
- 3 and 4. Bacilli in which spore-formation is going on.
5. Torula-form of the bacillus.

be invested with a thick hyaline gelatinous envelope. The rods form bright oval spores, being at the same time swollen up and ovoid. In Cohn's nourishing fluid they are capable of forming leptothrix, in which occur here and there huge spherical or oval swellings, which, according to Neelsen, probably represent gonidia.

¹ Cohn and Mifet, *Beitr. z. Biol. d. Pflanzen*, vol. iii. p. 128.

² *Beitr. z. Biol. d. Pflanzen*, vol. iii. p. 187.

CHAPTER XI.

BACILLUS : PATHOGENIC FORMS.

PATHOGENIC bacilli.¹

(a) *Bacillus of septicæmia of mice* (Koch).—By inoculating ordinary house mice with minute quantities of putrid fluids




FIG. 49.—FROM A SECTION THROUGH THE LUNG OF A MOUSE DEAD OF KOCH'S SEPTICÆMIA.

1. Small vessel filled with blood ; the white blood-corpuscles are filled with very minute bacilli.
2. Interalveolar tissue : in it a white corpuscle filled with the bacilli. Magnifying power 700.
3. A white blood-corpuscle more highly magnified, 1000.
(Stained with magenta.)

Koch found that occasionally one or another of these animals showed signs of conjunctivitis and sopor, and finally death followed in about forty to sixty hours. In these cases slight

¹ What has been said of the micrococci associated with open wounds and abscesses applies also to bacilli ; *i.e.* there are often bacilli present in the secretions of open wounds, and in the tissue of the base of ulcers ; and as the inflammation spreads, so also do the bacilli gradually invade the surrounding tissues. To mention one series of cases only, in ulcerations and in inflammations

œdema is found at the seat of inoculation; the spleen is large; in the œdematous tissue and in the blood-vessels, large and small, numbers of minute bacilli are found, chiefly contained in the white blood-corpuscles, but also free. They are very minute, about 0·0008 to 0·001 mm. long, to 0·0001 to 0·0003 mm. thick, isolated or in couples, or in chains of four or more. The smallest quantity of this blood invariably kills, with the same symptoms, house mice and sparrows, but not field mice. Rabbits inoculated with these bacilli in the skin of the ear or the cornea show only a local inflammation, and the tissues presently contain numerous bacilli of the same kind. Such animals,




FIG. 49.—FROM A SECTION THROUGH THE SMALL INTESTINE OF A MOUSE DEAD OF SEPTICÆMIA.

The figure represents a section through a small vein in the submucous tissue, filled with blood. At 1, there is a homogeneous substance and in it numerous bacilli, but these bacilli are much larger than the bacilli of Koch's septicæmia in the mouse.

Magnifying power about 700. (Stained with methylene blue and vesuvin.)

after the local effect has passed off, are protected against any further attack by the same bacilli. Koch cultivated these bacilli artificially on mixtures of aqueous humour and gelatine, of gelatine and peptone (1 per cent.), salt (0·6 per cent. NaCl), and sodium phosphate in sufficient quantity to produce a just alkaline reaction. The bacilli grow well on this mixture, and by repeated and rapid division form peculiar branched series.

(b) *Bacillus of septicæmia of man.*—In several cases of human septicæmia I have found in the blood-vessels of the swollen lymphatic glands large numbers of minute bacilli,

of the mucous membrane of the stomach and intestine, large numbers of bacilli are occasionally found on the surface of the inflamed parts, and gradually invading the inflamed tissue. Von Recklinghausen (*Virchow's Archiv*, vol. xxx.), von Wahl (*ibidem*, vol. xli.), saw minute pustular nodules in the inflamed gastric mucous membrane which were full of bacilli. Whether the presence and growth of these bacilli was the primary cause or only a concomitant symptom (due, for example, to the loss of active vitality of the tissue) remains to be proved.

slightly thicker than those just mentioned. They form continuous masses, both in the capillaries and in the minute veins, amounting in some cases to veritable emboli. They occur isolated or in short chains, their length about 0·001 to 0·0025 mm., their thickness about 0·0003 to 0·0005 mm. Arloing




FIG. 50.—FROM A SECTION THROUGH A LYMPHATIC GLAND OF MAN DEAD OF SEPTICÆMIA.

1. A blood-vessel which at one place is distended by and filled with minute bacilli.
 2. Lymph-corpuscles.
 3. Degenerated lymph-corpuscles.
- Magnifying power 700. (Stained with gentian violet.)

and Chauveau (mentioned in the *British Medical Journal*, Jan. 12, 1884) found in gangrenous septicæmia around wounds short bacilli, some containing one or two spores, which they consider as the true cause of the gangrene. They are destroyed when fresh by a temperature varying between 90° and 100° C. ; after drying, a temperature of 120° C. is required.

(c) *Bacillus of typhoid fever of man*.—Klebs¹ described in the inflamed Peyer's glands, in the mesenteric glands, larynx, and lungs of patients dead of typhoid fever certain bacilli, which are about 0·0002 mm. thick and of various lengths, forming filaments up to 0·05 mm. long. These bacilli form spores. Eberth² found in about 50 per cent. of cases of patients dead of typhoid fever, in the mesenteric glands and spleen, peculiar short bacilli, rounded at their ends, and occasionally slightly constricted in the middle; some of them




FIG. 51.—FROM A SECTION THROUGH THE MESENTERIC GLAND OF A PERSON WHO DIED OF TYPHOID FEVER.

1. Capillary blood-vessel filled with blood-corpuscles.
2. Large lymph-cell.
3. Nuclei.
4. Bacilli.

Magnifying power 700.

contained spores. The bacilli stain very freely with methyl-violet. It is, however, doubtful whether these bacilli can be considered as necessarily and intimately connected with typhoid fever, seeing that they are not constant, and only occur in the mesenteric glands and spleen, *i.e.* in localities into which an immigration of putrefactive bacilli from the bowels may easily take place; especially when we remember that in cases of typhoid fever that end fatally there constantly occur severe sloughing and necrosis of the mucous membrane of the Peyer's glands. The bowels in typhoid fever always contain innumerable masses of micrococci in colonies; and these micrococci are found not only in the tissue of the intestinal

¹ *Archiv f. exp. Path.* vol. xii.

² *Virchow's Archiv*, vols. lxxxiii., lxxxvii. See also Koch, *Mittheil. a. d. k. Gesundheitsamte*, i. 1881; and Gaffky, *ibid.* 1882.

mucous membrane but also in the mesenteric glands and spleen.¹



FIG. 52.—FROM A SECTION THROUGH THE KIDNEY OF A CASE THAT DIED AFTER MEAT-POISONING AT WELBECK.

The figure represents part of a glomerulus of a Malpighian corpuscle, in which some of the capillary blood-vessels are filled with the bacilli. Magnifying power 700.

1. Capsule of Malpighian corpuscle.
2. Capillaries filled with bacilli.
3. Capillaries empty.
4. Bacilli contained between capillaries.

(d) *Bacillus of choleraic diarrhoea from meat-poisoning.*—In July, 1880,² there occurred in Welbeck, Notts, an extensive outbreak of diarrhoea among over seventy-two persons who had

¹ Klein, *Reports of the Medical Officer of the Privy Council*, 1875.

² Report by Dr. Ballard in the *Reports of the Medical Officer of the Local Government Board*, 1880.

partaken of beef and ham sandwiches sold at Welbeck on the occasion of a sale of timber and machinery on the estate of the Duke of Portland. The infection showed itself after an incubation-period varying from twelve hours or less to forty-eight hours or more. The first symptoms were a sudden feeling of languor, nausea, griping in the abdomen, in some cases giddiness and fainting, and pain in the trunk. Then followed pain in the abdomen, diarrhoea, and vomiting, the diarrhoea being most constant. Four cases ended fatally. On *post-mortem* examination enteritis and pneumonia were most prominent. Part of the kidney was examined in microscopic sections, and it was found that many of the tubuli uriniferi contained hyaline casts; that the capillaries of the glomeruli of the Malpighian corpuscles, and the afferent arterioles, contained numbers of bacilli, some of the capillaries being distended by and plugged with masses of bacilli densely aggregated. In February, 1881, a similar but less extensive



FIG. 58.—ISOLATED BACILLI IN A SMALL ARTERY OF THE SAME KIDNEY AS IN PRECEDING FIGURE.

Some bacilli contain spores.

outbreak occurred at Nottingham, among fifteen persons that had partaken of certain baked pork. The symptoms were similar to those in the Welbeck outbreak. One case ended fatally. *Post-mortem*: bloody exudation in pericardium, intense pneumonia, mesenteric glands enlarged, enteritis, Peyer's glands enlarged. Bacilli similar to those of the above case were found in the blood, in the pericardial exudation, in the juice and in the bloody fluid filling the alveolar cavities of the inflamed lung, in the vessels of the kidney, in the submucosa of the inflamed Peyer's glands of the small intestine, in the blood-vessels of the spleen and around them.

The bacilli vary in length between 0·003 and 0·009 mm.; their thickness is about 0·0013 mm. They are rounded at their extremities, single or in chains of two, and some contain a bright oval spore, situated in the centre or at one end, and about 0·001 mm. thick. This was the case with the bacilli in the glomeruli of the kidney of the Welbeck case. The

bacilli containing spores were thicker than those without them.

Experiments by feeding and inoculation made on dogs and cats, rabbits, guinea-pigs, and mice, with the ham that had done the mischief in the Welbeck case produced positive results. In all cases we found pneumonia and hæmorrhage in the liver, peritonitis in some, spleen enlarged in most. The bacilli found in this ham were cultivated in the incubator in white of egg, and after two days' cultivation four white rats, and several guinea-pigs and white mice were inoculated, and they became ill after twenty-four hours; they were quiet, did not feed well, and were more or less soporous. When killed the spleen was found enlarged, and in the lungs were found hæmorrhage and hyperæmia, and in some cases extensive pneumonia.

Blood, pericardial exudation, and lung juice from the fatal Nottingham case inoculated into ten animals (guinea-pigs and white mice) produced fatal results in six, the other four were killed; but in all there was severe pneumonia, in eight out of the ten there was peritonitis, in four also pleuritis, and in two in addition enlargement of the liver and spleen. Bacilli were found in the blood and exudations of these animals. On cultivating blood and lung juice from the above case, a crop of bacilli was produced, which on inoculation proved very poisonous in the same way as in the previous cases.¹

(e) *Bacillus malarie*.—Klebs and Tommasi-Crudeli² described a bacillus occurring in the soil of the Roman Campagna, which they cultivated on gelatine. The rods are about 0·002 to 0·007 mm. long; they grow in cultures into long leptothrix filaments composed of short joints. The rods form spores either in the centre or at their ends. They grow well also in other media, e.g. albumen, urine, and glue. They require oxygen for their growth, and are therefore aerobic. According to Marchiafava,³ they occur also in the blood of patients suffering from malaria. Inoculations of rabbits with the cultivated or original bacilli produced a febrile disorder, which Klebs and Tommasi-Crudeli consider analogous to the human intermittent fever; but experiments made by Sternberg with material derived from the soil of malarious localities in America did not bear this out. The febrile disorder had nothing of the character of human intermittent fever, and

¹ Compare also Huber, *Archiv f. klin. Med.* xxv.

² *Archiv f. exp. Path.* vol. xi.

³ *Archiv f. exp. Path.* vol. xiii.

besides, could be produced by other bacilli than those of malarious soil.

(f) *Bacillus of ulcerative stomatitis in the calf.*—In the *Lancet* of May, 1883, A. Lingard and E. Batt described peculiar bacilli in ulcerations occurring on the tongue and buccal mucous membrane of the calf. "The typical ulcer in advanced



FIG. 54.—FROM A SECTION THROUGH NECROSSED AND ADJOINING INFLAMED PARTS OF THE EAR OF A RABBIT, INOCULATED WITH MATTER TAKEN FROM ULCERATIVE STOMATITIS OF THE CALF.

1. Necrosed part.
2. Inflamed tissue.
3. Bundles of bacilli.

Magnifying power 700. (Stained with magenta.)

cases consists of a sore with free overhanging edges. On section through the sore the tongue is found necrosed to a considerable depth." "Whenever the sore touches any other part of the mouth or cheek, the disease is communicated and rapidly spreads. In some cases similar necrotic changes had taken place in the lung. The line of junction of the necrotic with the healthy tissues was found to be occupied by a dense mass

of bacilli having the appearance of a dense phalanx advancing upon the healthy tissues. The disease has been proved capable of transmission (to the rabbit and mouse) by injection of the bacilli in question, which are equally numerous and virulent after passing through several generations by inoculation."

The disease often ends fatally in calves.



FIG. 55.—FROM A SECTION THROUGH TONGUE OF CALF, ULCERATIVE STOMATITIS.

1. Muscular fibres.
2. Inflamed tissue.
3. Bundles of the bacilli.

Magnifying power 700. (Stained with magenta.)

The best method of staining the bacilli was found to be this: The sections, both those prepared from the ulcerations of the calf's tongue and from the inoculated tissues of the rabbit, are immersed in a mixture of magenta and methyl-blue, then washed in spirit, and after clarifying in clove-oil are mounted in Canada balsam solution. The bacilli are stained deep pink, the inflamed tissue blue. The bacilli appear as thin rods in rows, thus forming a leptothrix-like growth. In

some of the long filaments the individual bacilli are not well shown. The filaments are either straight or more or less curved. The length of the single bacilli varies from 0·004 mm. or less to 0·008 mm. or more; the thickness is about 0·001 mm. Many of them contain spores. In the ear of the rabbit they invade the connective tissue as well as the cartilage over the whole extent of the ulceration and its neighbourhood. Mr.




FIG. 56—FROM A SECTION THROUGH THE CARTILAGE OF RABBIT'S EAR IN WHICH ULCERATION HAD BEEN PRODUCED BY INOCULATION WITH NECROSSED MATTER OF CALF'S TONGUE.

1. Cartilage capsules.
2. Bundles of good bacilli.
3. Bundles of degenerating bacilli.

Magnifying power 700. (Stained with magenta.)

Lingard found the same bacilli, having the same arrangement, in a case of noma in the human subject.

(g) *Bacillus of glanders*.—In 1882 Schütz and Löffler¹ ascertained the occurrence of peculiar bacilli in the nodules of the nasal mucous membrane and internal organs, such as the lung, spleen, and liver, of horses dead or dying from glanders.

¹ *Deutsche med. Wochenschrift*, 52, 1882.

These bacilli are very minute, being of about the size of tubercle-bacilli (see below), and are brought out by staining the tissues with a concentrated watery solution of methylene-blue, and after this washing with very dilute acetic acid. They succeeded in artificially cultivating the bacilli in the incubator at 38° C., on solid sterilised serum of horses' and sheep's blood, using for the purpose particles (see below, under "Tubercle-bacilli") of nodules of the lung and spleen of a horse dead of glanders. After two days (*i.e.* on the third day) there appeared on the surface of the inoculated material the first traces of the growth in the form of minute transparent droplets which consisted entirely of the characteristic bacilli. Cultivating these through several generations or transferences,




FIG. 57.—PUS OF A PULMONARY ABSCESS IN A HORSE DEAD OF GLANDERS.

1. The nuclei of pus cells.
2. The glanders-bacilli.

Magnifying power 700. (The preparation had been stained with methylene-blue.)

and then inoculating with them a horse, rabbits, guinea-pigs, and mice, positive results were obtained, especially in the guinea-pigs, which appear very susceptible to the disease. On the site of the subcutaneous inoculation appears an ulcer with indurated base, speedily enlarging and spreading; other ulcers follow in the neighbourhood, the neighbouring lymphatic glands become swollen, and the general infection follows, in the form of nodules and ulcers on the nasal septum, and nodules in the internal organs. In the guinea-pig a characteristic tumour of the testis, ovary, and vulva is often observed. In all these cases the diseased tissues and organs contained the characteristic bacilli.¹

¹ Other writers on the bacilli of glanders are Drs. Bouchard, Capitan, Charvin, in the *Revue médicale française*, Dec. 30, 1882; N. P. Wassilieff, in *Deutsche med. Woch.* 11, 1893, observed the bacilli in human glanders.

Bacillus of swine plague.—In a report to the medical officer of the Local Government Board for 1877-1878, I have shown that in this acute infectious disease the affected organs contain a form of bacterium in morphological respects identical with *Bacillus subtilis*, i. e. consisting of longer or shorter motile rods, capable of forming spores; further, that artificial cultures of these bacilli cause the disease in pigs after inoculation; and lastly, that mice and rabbits become affected with this disease after inoculation with material directly derived from the diseased organs of the pig or with artificial cultures. Last year Pasteur claimed to have cultivated from the blood of the pig affected with the disease a microbe which is not a bacillus, but a dumb-bell micrococcus. He states that he has produced




FIG. 58.—FROM A SECTION THROUGH THE INFLAMED INGUINAL LYMPH-GLAND OF A PIG DEAD OF SWINE PLAGUE.

1. A capillary blood-vessel filled with bacilli.
2. Reticulum of adenoid tissue.
3. A lymph-cell.

Magnifying power 700. (Stained with Spiller's purple.)

with these cultures fatal illness in pigeons and rabbits, and has also caused the plague in swine. I have been able to show by new experiments that Pasteur is wrong in all these points. First, I have proved that pigeons are altogether insusceptible to the disease, since inoculations with material directly derived from the diseased organs of the pig dead of swine plague, material which is well known to produce the disease in the pig, mouse, and rabbit, are altogether harmless to pigeons; and similarly cultivations of the true bacteria of swine plague do not affect pigeons in the least. According to Pasteur's statement the pigeons inoculated with his cultures of the dumb-bell micrococcus died with symptoms and with anatomical lesions almost identical with those of the form of

septicæmia known as fowl-cholera; and the conclusion is therefore forced upon us that Pasteur's cultures were contaminated with, or contained solely, the organism of this septicæmia. Similarly his rabbits probably died from the same disease, since these animals are exceedingly susceptible to septicæmia.

On examining the diseased tissues of pigs dead of swine plague by the modern methods of anilin staining, I ascertained that all the diseased organs (lungs, intestines, inguinal and bronchial lymph-glands) contain the characteristic bacilli, mostly filling and plugging minute blood-vessels. So do the diseased organs of mice and rabbits (spleen, liver, lung) dead of the disease.

Artificial cultivations made in broth and hydrocele fluid from diseased organs of the pig, mouse, and rabbit, after an



FIG. 59.—FROM A PREPARATION OF BRONCHIAL MUCUS OF A PIG DEAD OF SWINE PLAGUE.

1. Detached epithelial cells of alveoli.
2. Bacilli.
3. Micrococci.

Magnifying power 700. (Stained with Spiller's purple.)

incubation of twenty-four hours at temperatures ranging between 30° and 42° C. contain the above rods, which crowd the nourishing fluids, all being rather short, about 0.002 to 0.003 mm. long, and all possessed of the power of active locomotion, such as is known to be possessed by the septic bacterium *termo* and *bacillus subtilis*. During the following days of incubation, while the rods multiply, many of them lose their motility, grow longer, up to 0.005 mm. and more, and in some of the longer samples bright spores make their appearance, one spore at one or both ends or sometimes in the centre.

From these cultivations new cultivations may be made and carried on through successive generations, all cultures behaving




FIG. 60.—FROM A SECTION THROUGH THE KIDNEY OF RABBIT DEAD OF SWINE PLAGUE, SHOWING A MALPIGHIAN CORPUSCLE, THE CAPILLARIES OF THE GLOMERULUS BEING TRANSFORMED INTO HYALINE IMPERMEABLE CYLINDERS.

1. Bacilli.

Magnifying power 500. (Stained with Spiller's purple.)



FIG. 61.—BLOOD OF FRESH SPLEEN OF A MOUSE THAT DIED OF SWINE PLAGUE.

1. Blood discs.
2. A large nucleus.
3. Groups of minute bacilli.
4. Long bacilli.
5. Dumb-bells of bacilli.

Magnifying power 700. (Stained with gentian violet.)

in the same manner; and in all of them the rods only are present, and show exactly the same changes as in the parent culture.

The smallest droplet of any of these cultivations produces the disease in pigs, mice, and rabbits. The mice and rabbits



FIG. 62.—FROM A SECTION THROUGH A NECROTIC PATCH OF THE LIVER OF A MOUSE DEAD OF SWINE PLAGUE.

1. Tracts of liver cells shrunk.
 2. Capillary blood-vessels filled with very small micrococci, amongst which are seen the bacilli.
 3. Bacilli only.
- Magnifying power 700. (Stained with Spiller's purple and magenta.)

die with exactly the same appearances and with the same anatomical lesions as when they are inoculated with material directly taken from the diseased organs of a pig dead of swine



FIG. 63.—BACILLI OF SWINE PLAGUE, FROM AN ARTIFICIAL CULTURE, AFTER FORTY-EIGHT HOURS' INCUBATION.

Magnifying power 700. (Dried and stained with Spiller's purple.)



FIG. 64.—BACILLI OF SWINE PLAGUE, FROM AN ARTIFICIAL CULTURE, DURING SIXTH DAY OF INCUBATION.

- 1 and 2. Bacilli.
3. Bacilli in which spores have been formed. Magnifying power 700. (Fresh specimen.)

plague. Those animals generally die on the fifth, sixth, or seventh day, and on *post-mortem* examination show a characteristic swelling of the spleen, a characteristic disease of the

of the blood (containing not one of these of the liver tissue),
 and that the bacilli are not in the blood.

It is to be noted, however, that not all inoculations made
 in this manner are successful. In such animals always result
 in the formation of a large number of the characteristic bacilli as do
 the animals with the long and terminal bacilli of pigs lead
 to the formation of the blood of the pig, the bacilli
 of the blood of the pig, and as a rule from the blood of
 the pig, the bacilli are taken from the blood of rabbits
 and the blood of the pig.

It is to be noted, however, that pigs inoculated with
 the blood of these rods, started from the pig, mouse,
 and rabbit, of the plague, as with the diseased organs of a
 mouse, rabbit, or guinea-pig, a modification of the disease, which
 after a few weeks passed off completely. I have had pigs
 inoculated with the blood of the first time with artificial
 plague, and the second time with diseased organs of mouse and
 rabbit, and the third time the pig suffered from a mild form of
 the disease. The fourth time inoculated a third time with the
 blood of the diseased mouse, and of a pig lead of the plague;
 the pig died after a few days' suffering of the disease,
 but after a few days' rest they completely recovered. If
 normal and experimental inoculated pigs are inoculated with
 matter from the diseased fresh tissue of a pig lead of the
 plague, they are able to form a variant form of the disease.
 But in the above case they were protected by previous in-
 noculation, not altogether against a new attack but against a
 fatal attack.

Bacilli of Leprosy — Armand Hansen first ascertained the
 existence of large numbers of minute bacilli in the peculiar
 large leprosy-cells of Virchow, which occur in the nodules of
 leprosy patients. Nisser confirmed this, and considerably
 extended our knowledge of the bacilli, showing that they can
 be readily stained pink with carmalum or with Ehrlich's acid
 solution of eosin-haematoxylin. The bacilli are fine rods
 0.001 to 0.006 mm. long and less than 0.001 mm. thick.
 They are pointed at their ends, and always occur in masses
 within the large leprosy-cells of the leprosy tubercles of the
 skin and internal organs. But they are also present in the
 interstitial tissue of the nervous branches in the anæsthetic
 variety of the disease.¹ Some bacilli are motile, others not;

¹ *Virchow's Archiv*, vol. lxxix; and *Quart. Journ. of Micro. Sci.* 1880.

² *Comptes rend. Acad. Médicale*, 1881, Nos. 178, 179, and *Bates
 Archives of Pathology*, July, 1883.

some possess bright oval spores, and others are more or less beaded, owing to local collections of the protoplasm within their sheath. Neisser and Armauer Hansen have cultivated




FIG. 65.—FROM A SECTION THROUGH THE LARYNX OF A PATIENT DEAD OF LEPROSY.

Huge cells in fibrous connective tissue; the cells are filled with the leprosy bacilli.

Magnifying power 600. (Stained with magenta and vesuvin.)

them artificially in blood-serum and in solutions of meat-extract. Neisser has also shown that the characteristic leprosy-cells are only wandering cells modified by the growth and



FIG. 66.—BACILLI OF THE SAME PREPARATION AS IN PRECEDING FIGURE.

More highly magnified, 1000.

multiplication in them of the bacilli. In the blood the bacilli do not occur, but they spread probably only by way of the lymphatics.

Inoculation experiments on domestic animals and monkeys have hitherto failed.¹ Damsch² maintains, however, that he was able, by inoculation with leprous tissue into the peritoneal cavity and into the skin, to produce in cats a distinct increase and sprouting of the bacilli. Preparations of leprous nodules of the larynx and skin made by my friend, Mr. A. Lingard, and stained with Weigert's solution of magenta and vesuvin, showed the leprosy-bacilli completely filling all the cells,



FIG. 67.—CELLS OF THE LEPROSY NODULES OF MAN, FILLED WITH THE LEPROSY BACILLI (AFTER NEISSER).

small and large, spherical and spindle-shaped, contained between the connective-tissue bundles.

In a section through the liver of a bird (*Rhea*) that died in the Zoological Gardens in London, prepared by Dr. Gibbes after his method of staining for tubercle-bacilli, there were seen innumerable aggregations of larger and smaller pink masses (visible to the unaided eye as dots of the size of a pin's point to that of a pin's head or millet seed, and larger). Under



FIG. 68.—FROM AN ARTIFICIAL CULTURE OF BACILLUS OF LEPROSY (AFTER NEISSER).

the microscope these pink masses were seen to be composed of cells of various sizes, each filled with an enormous number of what appeared under a high power very short bacilli, much shorter than tubercle-bacilli. But they gave the same reaction as tubercle-bacilli. Here and there isolated cells of various sizes could be seen filled with the bacilli. In the large cells the cell-outline was becoming indistinct, and in some the cell-substance was seen to break down, whereby the

¹ Köbner, *Virchow's Archiv*, vol. lxxxviii. ; Hansen, *ibidem*, vol. xc.

² *Virchow's Archiv*, vol. xcii.

bacilli became free. In these respects, in the size, distribution, and character of the bacilli, there exists a remarkable similarity between the nodules in leprosy and the nodules just mentioned.




FIG. 69.—FROM A SECTION THROUGH A NODULE OF THE LIVER OF RHEA.

1. Cells of various sizes filled with minute bacilli ; owing to the smallness of the bacilli and to their being crowded in the cells and owing to the comparatively low magnifying power (300) the bacilli appear like dots.

(Stained with fuchsin and methyl-blue.)



FIG. 70.—TWO CELLS OF THE LEPROSY (?) NODULES IN THE LIVER OF A BIRD (RHEA).

The cell substance is crowded with minute bacilli, similar to leprosy-bacilli. Magnifying power 700. (Stained with magenta.)

(j) *Bacillus of malignant œdema* (Koch), *vibron septique* (Pasteur). By inoculating mice, rabbits, and especially guinea-pigs subcutaneously with a comparatively large quantity of earth, or of putrid fluid, one occasionally produces death in

twenty-four to forty-eight hours. This form of septicæmia is also called "Pasteur's septicæmia," and is of course distinct and different from Davaine's septicæmia.¹ At the seat of the inoculation and spreading from it into the subcutaneous tissue of adjoining parts there is much discoloration and occasionally hæmorrhage; a turbid offensively-smelling ichor fills the spaces of the subcutaneous tissue, and in it are found large numbers of bacilli, some motile, others not. The lungs are hyperæmic and have small hæmorrhagic spots. The spleen is




FIG. 71.—BLOOD OF A GUINEA-PIG DEAD OF KOCH'S MALIGNANT ŒDEMA.

1. Red blood discs.
2. White corpuscles.
3. Single bacilli.
4. Chain of long bacilli.
5. Leptothrix.

Magnifying power 700. (Stained with gentian violet.)

invariably enlarged and hæmorrhagic spots are often noticed on the peritoneum of the abdominal organs, and there is some peritoneal exudation. The blood of the spleen, of the liver, lung, and intestine, the serous coating of the abdominal organs,

¹ Rosenberger maintains (*Centralblatt f. d. med. Wiss.* 4, 1883) that the blood and exudation-fluids of rabbits dead of Davaine's or Pasteur's septicæmia can be effectually sterilised by heat without losing their specific action, reproducing on injection into fresh animals the disease with the recurrence of the organisms characteristic of the disease. Dowdeswell, however, states (*Proceedings of the Royal Society*, 221, 1882) that this is not the case, for on really effectual sterilisation by heat the organisms are killed, and the fluids become innocuous.

and the peritoneal exudation, contain the same bacilli as the subcutaneous exudation. Many of them include spores. By injecting the bacillus into the peritoneal cavity of guinea-pigs death is produced rapidly, especially after passing it through two generations, so rapidly indeed that the animals often die within sixteen hours (Burdon Sanderson and Klein). In all these instances a viscid transparent slightly but spontaneously coagulable exudation, poor in white and red corpuscles, is found in the peritoneal cavity, and the peritoneum of all parts is highly inflamed. Bacilli are present in it in enormous numbers, many of them containing spores. The blood of the heart does not contain bacilli immediately after death, but has them some hours after.

The bacilli in question are about 0·003 to 0·005 mm. long and a little over 0·001 mm. thick ; they are rounded at their ends ; they form chains of two and more, and these chains are straight or broken. They also form leptothrix, straight, or more commonly curved. The bacilli have been artificially cultivated by Pasteur¹ in blood-serum and in neutral solution of Liebig's meat extract. Gaffky² grew them on potatoes at 38° C. The artificial culture is capable of producing the malignant œdema, but it is always necessary to inject more than minimal quantities. The bacilli grown in fluids outside and inside the body form spores without free supply of air, and are therefore anaërobic (Pasteur).

In human fœcal matter there are always present innumerable masses of bacteria—micrococci, single and in dumb-bells, and in clumps of zoogloea, bacterium termo, and various species of bacilli, varying in thickness, length, and in motility, some being motile, others not. It has been shown by Bienstock (*Centralbl. f. med. Wiss.* 1888, p. 949) that a bacillus can be cultivated from normal human fœces which in many respects resembles the bacillus of malignant œdema ; it produces death in mice, but without the symptoms of malignant œdema.

Professor Rossbach has maintained (*Centralblatt f. d. med. Wiss.* 5, 1882) that when a solution of papayotin (the juice of *Carica papaya*) is injected into the veins of a rabbit, the animal dies, and shortly after death—even so short a time as fifty minutes after the injection—there are found in the blood large numbers of bacteria. Dowdeswell, however, states (*Practitioner*, May, 1883) that solutions of papayotin contain as a rule the spores of a motile bacillus which in all respects resembles *Bacillus subtilis* ; in artificial cultures in 10 per cent solutions of papayotin, in blood-serum, and in broth, these spores develop into bacilli which form leptothrix filaments, and in them spores soon make their appearance. Filtered papayotin solutions, when injected into the blood of rabbits, kill like unfiltered ones, but neither during life nor after the death of the animals could any organisms be detected in the blood. It appears to follow from these experiments, that papayotin solutions contain spores, and that these spores are those of a *Bacillus subtilis* which does not possess any specific pathogenic properties.

(k) *Bacillus of symptomatic anthrax* (Ger. *Rauschbrand* ; Fr. *charbon symptomatique*, Arloing, Cornevin, and Thomas,

¹ *Bull. de l'Acad.* 1877.

² *Mittheil. a. d. k. Gesundh.* 1880.

Bull. de l'Acad., 1881 ; Eng. *black leg, quarter-evil*). This disease, which is not uncommon in cattle, generally ends fatally and is very infectious. It is characterised by hæmorrhagic effusion (or "tumour") in the subcutaneous and inter-muscular tissues of one or other, or both, anterior or posterior extremities, in consequence of which the movements of the animal so affected become greatly impeded. The animals generally die in the course of the second or third day after infection. The subcutaneous tumour contains numerous bacilli, as do the abdominal and thoracic viscera.

The bacilli are about the size of those of malignant anthrax or a little thicker ; they are rounded at their ends and often include at one end a bright oval spore ; this is also present in the bacilli of the parenchymatous organs (as will be shown below, this never occurs in the bacillus of malignant anthrax). The bacilli are either single or form short chains. Some of the bacilli are motile.

Inoculations with them into the subcutaneous tissue of guinea-pigs, rabbits, sheep, and calves always prove fatal, the same subcutaneous hæmorrhagic effusions being produced.




FIG. 72.—BLOOD OF A GUINEA-PIG DEAD OF SYMPTOMATIC ANTHRAX.

Blood-corpuscles and between them several bacilli.
Magnifying power 700. (Stained with Spiller's purple.)

Injections of small quantities of bacillus-containing material into the veins produces only a slight febrile disorder ; large doses produce death. Animals in which by intravenous injection of small doses slight illness has been produced are afterwards protected against the fatal dose. But minimal doses injected subcutaneously also produce only a slight

transitory swelling, and the animal so treated is afterwards protected against the fatal dose (Arloing, Cornevin, and Thomas). The spores of the bacilli when heated up to 85° C. for six hours lose their virulence (Arloing, Cornevin, and Thomas).

(*l*) *Bacillus anthracis*.—Pollender,¹ Brauell,² Davaine,³ and then Bollinger⁴ recognised in the blood of animals dead of malignant anthrax the presence of stiff short and long rods, which Davaine called *bactéridie du charbon*. They were identified by Cohn⁵ as bacilli in morphological respects similar to bacillus subtilis, except that the bacilli anthracis are non-motile.

Koch⁶ showed the ubiquitous distribution of these bacilli in the blood of the organs, and especially of the spleen. He




FIG. 73.—HEART'S BLOOD OF A MOUSE DEAD OF ANTHRAX.

1. Blood-discs.
2. White blood-corpuscle.
3. Bacilli anthracis.

Magnifying power 700. (Fresh specimen.)

succeeded in cultivating these bacilli artificially, taking a bit of spleen of a mouse (which animal is very susceptible to fatal anthrax), and watching the growth of the bacilli in a microscopic specimen. He saw that the rods multiply by division, and that they grow into long, homogeneous-looking, straight or twisted filaments in which after some time, and with free access of air, bright oval spores make their appearance, while the filaments become homogeneous and swollen. These spores become free, and when artificially cultivated or injected into a rodent animal, germinate into the characteristic bacilli;

¹ *Viertelj. f. gericht. Med.* 1855.

³ *Comptes Rendus*, lvii. 1863.

⁵ *Beitr. s. Biol. d. Pflanzen*, vol. ii.

² *Virchow's Archiv*, vol. 14, 1858.

⁴ *Med. Centralblatt*, June, 1872.

⁶ *Ibid.* vol. iii.

these elongate and divide, and in artificial cultures again grow into the long leptothrix filaments, which again form spores. Koch¹ saw in preparations of aqueous humour kept at 35° C. in the incubator the spores germinating after three to four hours. The single bacilli as they present themselves in the blood



FIG. 74.—PART OF A BLOOD CLOT FROM THE HEART OF A MOUSE DEAD OF ANTHRAX. Magnifying power 500. (Stained with Spiller's purple.)

measure between 0·005 and 0·02 mm. in length, and 0·001 to 0·0012 in thickness; they are truncated.² The spores produced by growing the bacilli with free access of air are about 0·001 mm. thick, and about 0·002 to 0·003 mm. long. They are not stained by dyes and differ herein from the bacilli.

In the human subject malignant anthrax occurs as "wool sorter's disease": for the ætiology and pathology of this malady see Spears (*Report of the Medical Officer of the Local Government Board*, 1881 and 1882) and Greenfield (*ibid.* 1881).

All rodents and herbivorous animals are susceptible to anthrax; rats are, however, infected with difficulty, pigs are very insusceptible, and so are dogs and cats. Infection of animals can be produced by inoculation into the skin and subcutaneous tissue, intravascular injections, and by inhalation of spores (Buchner, *Untersuchungen über niedere Pilze*, by Prof. v. Nägeli, 1862, p. 178). In wool sorter's disease the usual mode of infection is by inhalation of spores adhering to the wool of the fleeces of animals (sheep, goats) dead of anthrax. As in rodents infected with anthrax, so also in man, the blood-vessels of all organs contain the bacilli, and extravasations of the infected blood are frequent in many parts of the body. The presence of bacilli in the extravasations into the mucous membrane of the trachea and bronchi does not necessarily mean that these parts represent the points of entrance of the bacilli into the system, as Greenfield seems to regard as self-evident (*Reports of the Medical Officer of the Local Government Board*, 1881). As a matter of fact I find in every lung of mouse, rabbit, and guinea-pig, dead after subcutaneous inoculation with anthrax, bacilli antracis in the alveolar cavities and in the smaller and larger bronchi. Ingestion of bacillar material is sometimes followed by anthrax, but in

¹ *Beitr. z. Biol. d. Pflanzen*, vol. ii. part ii. p. 288.

² It is generally assumed that the bacilli are the same in all animals affected with splenic fever, but this is most undoubtedly not the case, as has been already pointed out by Huber (*Deutsche med. Woch.* 1881); the bacilli of the guinea-pig are thicker than those of the mouse or sheep, and these again are thicker than those in the rabbit.

these cases abrasions in the mucous membrane of the mouth, pharynx, or gut, may have been the real place of entrance. Mice fed with anthrax material do not become infected (Klein, *ibid* 1881). But the reported cases of intestinal mycosis (see for the literature of this subject, Koch, "Ätiologie d. Milzbrandes," *Mittheil. a. d. k. Gesundheitsamte*, 1881,) seem nevertheless to indicate that such a mode of infection, namely, by the alimentary canal, is not excluded. Compare also Falk, *Virchow's Archiv*, vol. xciii.

Rodents inoculated with the bacillus of the blood or spleen of an animal dead of anthrax, or with the bacillus or spores of an artificial culture, die generally within forty-eight hours ; in some instances in twenty-four to thirty hours, in other exceptional instances after forty-eight to sixty hours. The




FIG. 75.—FROM A PREPARATION OF HEART'S BLOOD OF A GUINEA-PIG DEAD OF ANTHRAX.

1. Red blood-discs.
2. White corpuscle.
3. Bacilli anthracis.

Magnifying power 700. (Stained with Spiller's purple.)

blood in all instances contains the bacilli, the spleen is large and full of bacilli, and so are the blood-vessels of most other organs, the exudations, and the urine. In the placenta of a pregnant guinea-pig dead in consequence of inoculated anthrax, I have seen that the bacilli keep strictly as a rule within the maternal blood-vessels, and are wholly absent in the blood of the vessels of the fœtus. Subcutaneous inoculation or injection into the cutis of the minutest quantity of bacillus-containing material (blood or artificial culture) invariably produces death. Subcutaneous injection of bacillus-containing material in the guinea-pig almost always produces a characteristic œdema, spreading sometimes over a large area. The œdematous fluid is clear and contains only a few bacilli.

Archangelski (*Centralblatt f. d. med. Wiss.*, 1888, p. 257) claims to have ascertained that if an animal be inoculated with anthrax, many hours before the bacilli appear in the blood, there are present numbers of spores. Just before death they all become changed into the bacilli. He further maintains that those spores taken from the blood can be shown to multiply by division, and without changing into bacilli, by cultivating them artificially with exclusion of oxygen. I have shown, however (*Reports of the Medical Officer of the Local Government Board for 1883*), that none of these assertions are borne out by actual observation, and that they are erroneous.

Any fluid containing proteid material is a suitable nutrient medium for the bacilli; they grow abundantly at all temperatures between 15° and 43° C., best between 25° and 40° C. They elongate and divide rapidly, and the bacilli grow out into long curved and peculiarly twisted filaments which often form bundles, the individual filaments being twisted round one another like the strands of a cable.

The bacillus anthracis grows best in neutral fluids, but to a limited extent also in acid or alkaline fluids containing proteid




FIG. 76.—FROM AN ARTIFICIAL CULTURE OF BACILLUS ANTHRACIS.
Convolutions of threads, each composed of bacilli.
Magnifying power 300. (Stained with Spiller's purple.)

material. When growing in neutral nourishing fluids, it forms on the bottom of the fluid characteristic fluffy whitish masses, which are convolutions of the characteristic filaments. These appear homogeneous in the fresh state, their ends being slightly

through white mice¹ it loses this virulence for sheep and cattle. The blood of white mice dead of anthrax does not kill sheep; it produces only a transitory illness and the animals are, for a time at least, protected against virulent




FIG. 82.—FROM A SECTION THROUGH THE KIDNEY OF A RABBIT DEAD OF ANTHRAX.

The capillaries of the cortex are naturally injected with the *Bacillus anthracis*.

1. A glomerulus.

2. Capillaries surrounding the convoluted uriniferous tubules not shown here.
Magnifying power 450. (Spiller's purple.)

anthrax. The blood of guinea-pigs dead of anthrax produces illness, sometimes death, in cattle, but as a rule does not kill (Sanderson and Duguid), and the blood of the biscachia of

¹ Klein, *Reports of the Medical Officer of the Local Government Board*, 188

South America does not kill cattle, while it gives them a transitory illness, and after this immunity for a time.¹ Again, Pasteur's "vaccine," which as a rule (but not without exception) does not kill sheep or cattle, is fatal to rodents.² From all this it follows that as regards virulence the bacilli anthracis differ in the different species of animals, and in them acquire different qualities. A culture that does not kill mice, such as an artificial culture of blood-bacillus after one or two weeks' incubation at 20°—35° C., or a culture that for other reasons, as when attenuated by heat or antiseptics, does not produce fatal anthrax in guinea-pigs, fails to give to these animals any immunity whatever. Rodents, so far as my experience goes, either die of inoculation with anthrax-bacilli or they do not die; but they cannot be provided with immunity by any attenuated virus.

Koch³ maintains that in neutral chicken-broth the bacilli growing at 42° C. lose their virulence in thirty days, and at 43° C. in six days, first for rabbits, then for guinea-pigs, and lastly for mice. I am quite sure from my own observations, that these results are not uniformly obtained, since I have seen anthrax-bacilli very virulent both for rabbits and guinea-pigs even after growing for thirty-six days at 42·5 C.

Bacillus anthracis is capable, as we have seen, of growing well outside the body, and, when well supplied with oxygen from the air, of forming spores which represent the permanent seeds. Thus if animals, such as sheep and cattle, die of anthrax in a field, the bacilli of the effusions from such animals (*e.g.* urine, blood, effluvia from the mouth and nostrils) always contain numbers of the bacilli, and these will be able to grow indefinitely on the surface of the soil, there being always present a large amount of suitable nourishing material, like vegetable and animal decaying matter, and as free access of air is always insured they will eventually form spores. Such soils, owing to the presence of these spores, will remain a permanent source of infection to sheep and cattle sojourning on them (Koch).

(*m*) *Bacillus tuberculosis* (Koch).—In all cases of tuberculosis in man, cattle (Perlsucht) and monkeys, of tuberculosis

¹ Roy, *Nature*, December, 1883.

² Klein, *Reports of the Medical Officer of the Local Government Board*, 1882. Similar results have been obtained by Gaffky (*Mittheil. a. d. k. Gesundheitsamte*, 1882).

³ *Ueber d. Milzbrandimpfung*, 1882.

Spores of bacillus anthracis stand heating to 100° C. in the dry state for over an hour without being killed; in the moist state, *e.g.* exposed to steam at 100° C., they are killed after 15 minutes' exposure (Koch).

When up, it will be noticed on incubation that the fluid remains acid. All the growth, in the shape of the fluffy whitish mass, takes place at the bottom.

After a few days' incubation, no matter what the temperature is, many of the bacilli and their leptothrix-filaments show signs of degeneration, consisting in the granular disintegration and absorption of the protoplasmic contents of the



78.—FROM A PREPARATION OF THE BLOOD OF SPLEEN OF A GUINEA-PIG DEAD OF ANTHRAX.

1. White blood-corpuscle.
2. Red blood-discs, shrunken.
3. Chains of bacillus anthracis.
4. Degenerating bacilli, the sheath only being preserved.

(Magnifying power 700. (The preparation has been stained with gentian-violet.)

li and their filaments, at first only here and there, but multiplied by over longer pieces. Such bacilli and leptothrix-filaments appear in such places as if empty. This is also evident in the bacilli of the blood and spleen of an animal infected with anthrax even at the point of death or soon after death, if the number of bacilli is great.

Another form of degeneration consists in the filaments of bacilli becoming much curled and swollen, and finally degenerated into an amorphous debris.

As long as the bacilli grow in the depth of a fluid they

never form spores, but when grown on the surface with free access of air, or on solid media (*e.g.* serum gelatine, gelatine broth, Agar-Agar, potato, &c.), the bacilli, having developed into filaments, proceed to form spores. But they may form spores even in fluid media if by some accident, either by




FIG. 79.—FROM AN ARTIFICIAL CULTURE OF *BACILLUS ANTHRACIS* IN BROTH AFTER MANY DAYS INCUBATION.

The threads are swollen and curled up, and in many places the protoplasm has disappeared, leaving the sheath and septa distinct. Magnifying power 700. (Stained with Spiller's purple.)


sticking to the glass vessel containing the fluid or by means of a cotton-wool fibre, some of the bacilli remain on the surface of the fluid. This formation of spores is not due to exhaustion of the nourishing medium, as is maintained by Buchner—it has, in fact, nothing to do with it—but represents the last stage in the life-history of the bacilli, provided they have an ample supply of oxygen. If this latter condition is not fulfilled, as when they are grown at the bottom of a fluid, the bacilli gradually degenerate as mentioned above.

Spore-formation occurs, *cæteris paribus*, at all temperatures between 18° and 45° C. Koch found 15° C. the lower limit. Pasteur states that in a nutrient medium exposed to a temperature of 42° to 43° C. the bacilli are not capable of forming spores; but this is not correct, for when the bacilli are growing on the surface of the nutrient medium, they form spores even at a temperature of 44° to 45° C., as I have conclusively shown by growing them on Agar-Agar and peptone mixture. The spore-formation consists in the appearance of a bright glistening spherical body in the protoplasm of an elementary mass or cell; this body gradually enlarges till it reaches its full size, becoming at the same time oval. The bacilli at these points are thicker than where no spore-

tubercle-bacilli have been shown to exist. They are most
 aersously found in the caseous masses in the lung found in



g. 84.—FROM A PREPARATION OF CASEOUS MATTER FROM PULMONARY
 DEPOSITS IN BOVINE TUBERCULOSIS, STAINED AS IN PRECEDING FIGURE.
 Magnifying power 700.



g. 85.—FROM A SECTION THROUGH TUBERCULOUS DEPOSITS IN THE LUNG
 OF A COW.
 Two giant-cells and two small cells containing tubercle-bacilli.
 Magnifying power 700.

ine tuberculosis. Here Koch found them not only scattered
 ough the caseous masses, but also in the well-known giant-
 ls ; in some cases they form a more or less regular zone in

affects greater and greater numbers, and when the fluid is exhausted for the formation of new bacilli, it necessarily follows that the whole growth gradually becomes involved in the process of degeneration, the whole mass becoming smaller, and finally only débris is left. Such cultures, namely, those in which the degeneration involves the whole mass of the bacilli, are quite innocuous when inoculated into animals, or into fresh nourishing media. But as long as there are any good protoplasmic elements of the bacilli left, the culture is virulent to rodents, with the exception of mice, as will be stated presently; and it is capable, when transferred to new suitable nourishing media, of starting new cultures that prove virulent to all rodents and sheep.

The same holds good of the bacilli in the blood and organs of an animal dead of anthrax, provided the animal be not opened, and its organs, exudations, or urine be not exposed to the free air; for the bacilli not exposed to the air gradually degenerate, and the blood and organs of such an animal, although at first deadly poison to other susceptible animals, become at length quite innocuous. Systematic observation has shown me that small animals, such as mice and guinea-pigs, when kept unopened or buried in earth, become quite innocuous after five to eight days, the anthrax-bacilli having by this time, by degeneration, altogether disappeared from the blood, spleen, and other organs. Pasteur's statement that in animals dead of anthrax and buried, the bacilli form spores, that these spores are taken up by earthworms and carried to the surface of the soil, where they are deposited with their castings and thus are capable of infecting animals grazing or sojourning on this soil, is not borne out by the above observations. And further, Koch has proved¹ by direct experiment that spores of anthrax-bacilli, when mixed with earth in which worms are present, are not taken up by these creatures.

Drying bacilli of the blood or of a culture in a thin layer invariably kills them, but the spores remain unaffected.

The bacilli of the blood of a rodent dead of anthrax are always thinner than the bacilli cultivated in a neutral fluid medium like pork-broth.

Cultivation of the blood-bacilli at temperatures varying between 20° and 40° C. in any suitable nourishing material, solid or fluid, however many transferences (new cultivations, or so-called new generations) be made, always yields a crop of virulent bacilli. It is absolutely incorrect to say, as Buchner²

¹ *Mittheil. a. d. k. Gesundheitsamte*, 1881.

² *Ueber d. Erzeug. des Milzbrandes*, Munich, 1860.

then caseating in the centre ; enlargement and subsequent caseation of the bronchial glands.

Owing to the fact that the tubercle-bacilli require for their growth high temperatures (38°—40° C.), it is evident that, unlike some other pathogenic organisms, they do not thrive in the outside world in temperate climates.

Inoculation with the pure bacilli into the anterior chamber of the eye of rabbits and guinea-pigs produces the characteristic tuberculosis described by Cohnheim and Salomonsen. After an incubation of from two to three weeks there appears on the iris a crop of minute grey tubercles enlarging and undergoing caseous degeneration. Later on general tuberculosis of the eyeball and other organs follows. So that Cohnheim's assertion, that only tuberculous matter implanted into the anterior chamber of the eye can produce this outbreak of a crop of tubercles on the iris, is, by Koch's observations, strengthened in the highest degree ; the tubercle-bacilli present in, and characteristic of, true tubercles are thus manifestly connected with the real cause of the morbid growth. A large number of pathologists have, since the publication of Koch's paper, devoted themselves to various parts of this question of the relationship of the tubercle-bacilli to the tuberculous process, and have, with few exceptions, verified Koch's observations. The chief opposition, leaving out of account those who, either from imperfect technical skill in the manipulation and staining of the bacilli, or by reason of the inadequate number of their observations, have denied Koch's statements, comes mainly from observers who, like Toussaint, Klebs, and Schüller, maintain that tuberculosis is due to a micro-organism which is a micrococcus and not a bacillus, or who, like Schottelius and others, do not admit that human and bovine tuberculosis are the same, and are, therefore, not interchangeable, which they ought to be if in both the same bacillus occurs, and if this bacillus is the *vera causa morbi*. But there can be no doubt that a vast number of competent observers have fully verified Koch's dictum, that the tubercle-bacilli are specific and different from other bacilli, except those of leprosy, as regards their chemical nature (compare their behaviour to nitric acid) ; and that wherever they are present in the sputum we have to deal with real tuberculosis, wherever after repeated examinations they are found to be absent there is no tuberculosis. This has by this time, although not much more than a couple of years has elapsed since Koch's first publication, become in the hands of all competent workers a matter of daily practical application, especially as regards the examination for bacilli of the sputum of patients suspected of tuberculosis.

they contain good bacilli at all.¹ But fresh cultures made of such bacilli invariably produce a growth which is fatal to all rodents during the first or second week.

Pasteur has stated that blood-bacilli which have become attenuated in virulence by exposure to 42° to 43° C. for twenty days are capable of starting new cultures of attenuated virus. This I question, for I find that such a culture starts new cultures of virulent bacilli; in the same way the bacilli of a culture that is only "vaccine" for sheep, when it is inoculated into a guinea-pig kills it with anthrax, and then yields bacilli that are fatal to sheep.

Blood-bacilli exposed to a temperature of 55° C. or to a solution of $\frac{1}{2}$ to 1 per cent. of carbolic acid, lose their virulence (Toussaint). Chauveau found that exposure to a temperature of 52° C. for fifteen minutes, or of 50° C. for twenty minutes, destroys the virulence of the blood-bacilli. Pasteur² ascertained that by cultivating blood-bacilli in chicken-broth at 42°—43° C. they lose their virulence after twenty days' cultivation, not as Pasteur thinks owing to the action of oxygen, but owing to the high temperature; and when such bacilli are injected into sheep and cattle they do not kill though they induce sometimes a slight illness. After this illness has passed off, the animals are protected against virulent anthrax. But with reference to this "vaccination," it must be borne in mind that twenty days' cultivation of blood-bacilli at 42°—43° C. does not always yield attenuated virus,³ and also that sheep and cattle not killed by inoculation of attenuated virus produced by Pasteur's method⁴ or by other means (see below), although they are protected against virulent anthrax, remain so only for a limited time, probably about nine months.

In all these experiments with the anthrax-bacillus it is necessary to bear in mind that by passing the bacilli through different species of animals they become endowed with different qualities, and that bacilli which are fatal to some are not fatal to all animals. While, for instance, the blood-bacillus of sheep or cattle dead of anthrax invariably produce death when inoculated into sheep or cattle, after passing

¹ Klein, *Reports of the Medical Officer of the Local Government Board*, 1881

² *Comptes Rendus*, 1881; *Transactions of the International Medical Congress* in London, 1881, vol. i.

³ Klein, *Reports of the Medical Officer of the Local Government Board*, 1882

⁴ Pasteur thinks that such cultures remain free of spores because of the temperature of 42°—48° C.; but this is not so, as has been pointed out above; statement only holds good so long as the bacilli are prevented from growing on the surface.

ugh white mice¹ it loses this virulence for sheep and
 2. The blood of white mice dead of anthrax does not
 sheep; it produces only a transitory illness and the
 als are, for a time at least, protected against virulent



. 82.—FROM A SECTION THROUGH THE KIDNEY OF A RABBIT DEAD OF ANTHRAX.
 capillaries of the cortex are naturally injected with the Bacillus anthracis.
 glomerulus.
 capillaries surrounding the convoluted uriniferous tubules not shown here.
 Magnifying power 450. (Spiller's purple.)

rax. The blood of guinea-pigs dead of anthrax produces
 ss, sometimes death, in cattle, but as a rule does not kill
 erson and Duguid), and the blood of the biscachia of
 ein, *Reports of the Medical Officer of the Local Government Board, 1882.*

South America does not kill cattle, while it gives them a transitory illness, and after this immunity for a time.¹ Again, Pasteur's "vaccine," which as a rule (but not without exception) does not kill sheep or cattle, is fatal to rodents.² From all this it follows that as regards virulence the bacilli anthracis differ in the different species of animals, and in them acquire different qualities. A culture that does not kill mice, such as an artificial culture of blood-bacillus after one or two weeks' incubation at 20°—35° C., or a culture that for other reasons as when attenuated by heat or antiseptics, does not produce fatal anthrax in guinea-pigs, fails to give to these animals any immunity whatever. Rodents, so far as my experience goes, either die of inoculation with anthrax-bacilli or they do not die; but they cannot be provided with immunity by any attenuated virus.

Koch³ maintains that in neutral chicken-broth the bacilli growing at 42° C. lose their virulence in thirty days, and at 43° C. in six days, first for rabbits, then for guinea-pigs, and lastly for mice. I am quite sure from my own observations, that these results are not uniformly obtained, since I have seen anthrax-bacilli very virulent both for rabbits and guinea-pigs even after growing for thirty-six days at 42°·5 C.

Bacillus anthracis is capable, as we have seen, of growing well outside the body, and, when well supplied with oxygen from the air, of forming spores which represent the permanent seeds. Thus if animals, such as sheep and cattle, die of anthrax in a field, the bacilli of the effusions from such animals (*e.g.* urine, blood, effluvia from the mouth and nostrils) always contain numbers of the bacilli, and these will be able to grow indefinitely on the surface of the soil, there being always present a large amount of suitable nourishing material, like vegetable and animal decaying matter, and as free access of air is always insured they will eventually form spores. Such soils, owing to the presence of these spores, will remain a permanent source of infection to sheep and cattle sojourning on them (Koch).

(*m*) *Bacillus tuberculosis* (Koch).—In all cases of tuberculosis in man, cattle (Perlsucht) and monkeys, of tuberculosis

¹ Roy, *Nature*, December, 1888.

² Klein, *Reports of the Medical Officer of the Local Government Board*, 1882. Similar results have been obtained by Gafky (*Mittheil. a. d. k. Gesundheitsamt*, 1882).

³ *Ueber d. Milzbrandimpfung*, 1882.

Spores of bacillus anthracis stand heating to 100° C. in the dry state for over an hour without being killed; in the moist state, *e.g.* exposed to steam at 100° C., they are killed after 15 minutes' exposure (Koch).

artificially produced (by inoculation with human or bovine tuberculous matter) in cats, guinea-pigs, rabbits, and rats, and in spontaneous tuberculosis in birds (hens), Koch¹ found in the fresh state, and particularly after staining with methylene-blue and vesuvin, peculiar fine bacilli, some with bright oval spores, some without, some smooth and homogeneous-looking, others more of a beaded appearance. One cubic centimetre of a concentrated alcoholic solution of methylene-blue is mixed with 200 .ccm. of distilled water ; to this are added two ccm. of a ten per cent. solution of caustic potash. In this solution the fresh or hardened sections or particles of tubercles are kept for half an hour if heated up to 40° C., or



FIG. 68.—FROM A PREPARATION OF HUMAN TUBERCULOUS SPUTUM, STAINED AFTER THE EHRlich-WEIGERT METHOD.

The nuclei are stained blue, the tubercle-bacilli pink. Magnifying power 700.

for twenty-four hours if not heated. After this the preparation is stained for two minutes in a filtered concentrated watery solution of vesuvin, then washed in distilled water. On examination with a $\frac{1}{2}$ oil-immersion lens and Abbé's condenser, it will be found that all the elements are stained brown with vesuvin except the bacilli, which are blue. A still more successful and more delicate reaction is shown by the

¹ *Berliner klin. Wochenschrift*, xv. 1882.

bacilli if the preparation is stained after Ehrlich's method. About 5 ccm. of pure anilin (anilin oil) are well mixed with 100 ccm. of distilled water and filtered; to this is added a saturated alcoholic solution of fuchsin, and with this the preparation is stained for a quarter to half an hour. It is then washed for a few seconds in a mixture of one part of nitric acid and two parts of water, and then is well washed in distilled water. The preparation when now examined shows no trace of colour except in the tubercle-bacilli, which retain the red colour of the fuchsin. The tissue may now be stained either with vesuvin or methylene-blue, which make the groundwork brown or blue, but the bacilli remain red. This reaction after washing with nitric acid is exceedingly delicate, and is perfectly characteristic and trustworthy, as a putrefactive organisms become discoloured by the washing with nitric acid, the tubercle-bacilli only retaining the colour. There are other methods which are very good; those of Weigert and of Gibbes¹ are very quick and trustworthy in the action.

Weigert has devised a staining fluid which gives very beautiful results and is very useful for staining sections, fresh or hardened; it is as follows:—Take of a two per cent. watery solution of gentian-violet 12 ccm., and of a saturated watery solution of anilin oil 100 ccm. Mix. This is used like an ordinary staining-fluid for the first stain. For the second or contrast stain the following solution is used:—

Bismarck brown	1 gramme.
Spiritus vini rectificati (sp. gr. 830)	10 ccm.
Distilled water	100 ccm.

The sections remain in a few drops of this solution for fifteen minutes. This method yields the finest specimens of tubercle bacilli in sections through tuberculous growths that I have seen; unfortunately the colour of the bacilli is very liable to fade.

In the case of tuberculous sputum, or similar matter, a small droplet or particle is spread out in a thin layer on a cover-glass, well dried by passing it over the gas-flame of a Bunsen burner, and then stained in the way described in Chapter I. Sections of tubercles, fresh or hardened, are stained without first drying.

In all cases of human tuberculosis, particularly in the sputum, in caseating scrofulous glands, in bovine tubercles, and artificially-induced tubercles and caseating glands of rodent

¹ *Lancet*, August 5, 1883.

the tubercle-bacilli have been shown to exist. They are most numerous found in the caseous masses in the lung found in



FIG. 84.—FROM A PREPARATION OF CASEOUS MATTER FROM PULMONARY DEPOSITS IN BOVINE TUBERCULOSIS, STAINED AS IN PRECEDING FIGURE.

Magnifying power 700.




FIG. 85.—FROM A SECTION THROUGH TUBERCULOUS DEPOSITS IN THE LUNG OF A COW.

Two giant-cells and two small cells containing tubercle-bacilli.
Magnifying power 700.

bovine tuberculosis. Here Koch found them not only scattered through the caseous masses, but also in the well-known giant-cells; in some cases they form a more or less regular zone i

the peripheral portion of the cell. But according to Koch the bacilli by and by disappear again from the giant-cells.

The bacilli do not show any motility and often include spores; they are thus capable of forming spores within the body. Owing to these spores, human phthisical sputum retains its virulence even after drying for considerable periods. Koch cultivated the bacilli artificially, *i.e.* outside the body, and by carrying on the cultivation for several successive transmissions succeeded in isolating and clearing them from the tuberculous tissue. These pure bacilli, no matter how many times they have been transferred, no matter how far removed from their original breeding-ground, always produced the characteristic disease when inoculated into suitable animals. The cultivation succeeded equally with material derived from human tubercles, from bovine tubercles, and from the artificially-induced tuberculosis of guinea-pigs. The bacilli grow well at a temperature varying between 37° and 39° C. in solid serum, Agar-Agar peptone mixture, and solidified hydrocele fluid.¹ (See Chapter II.) An incision is made into a tubercle with clean (overheated) scissors, and a particle of a tubercle is taken up with the point of a clean (overheated) needle and deposited on the top of one of these sterile solid media kept in a test-tube plugged with sterile cotton-wool. After keeping it for ten days to a fortnight in the incubator at 37°—39° C. the first traces of growth make their appearance in the shape of small dry whitish scales, which gradually increase in size until they coalesce. These scales are made up of the typical tubercle-bacilli lying closely side by side; some of the bacilli are longer, others shorter, and many of them have spores. New cultures may be established from these bacilli. Inoculation with them or with further cultivations into the subcutaneous tissue, peritoneal or pleural cavity of guinea-pigs and rabbits, produces after three, four, or more weeks, the typical lesions characteristic of artificial tuberculosis; namely, swollen lymphatic glands near the seat of inoculation, with subsequent caseation and ulceration; enlargement of the spleen due to numerous whitish tubercles, the larger ones caseous; enlargement of the liver, which is mottled by the presence of uniformly distributed whitish points and streaks, which by and by become confluent and caseous; tuberculosis of the peritoneum; isolated tubercles in the lungs, at first grey and transparent,

¹ Solidified hydrocele fluid has been successfully used for the cultivation of the tubercle-bacilli, not by Koch, but by my friend Mr Makins of St. Thomas's Hospital.

then caseating in the centre; enlargement and subsequent caseation of the bronchial glands.

Owing to the fact that the tubercle-bacilli require for their growth high temperatures (38°—40° C.), it is evident that, unlike some other pathogenic organisms, they do not thrive in the outside world in temperate climates.

Inoculation with the pure bacilli into the anterior chamber of the eye of rabbits and guinea-pigs produces the characteristic tuberculosis described by Cohnheim and Salomonsen. After an incubation of from two to three weeks there appears on the iris a crop of minute grey tubercles enlarging and undergoing caseous degeneration. Later on general tuberculosis of the eyeball and other organs follows. So that Cohnheim's assertion, that only tuberculous matter implanted into the anterior chamber of the eye can produce this outbreak of a crop of tubercles on the iris, is, by Koch's observations, strengthened in the highest degree; the tubercle-bacilli present in, and characteristic of, true tubercles are thus manifestly connected with the real cause of the morbid growth. A large number of pathologists have, since the publication of Koch's paper, devoted themselves to various parts of this question of the relationship of the tubercle-bacilli to the tuberculous process, and have, with few exceptions, verified Koch's observations. The chief opposition, leaving out of account those who, either from imperfect technical skill in the manipulation and staining of the bacilli, or by reason of the inadequate number of their observations, have denied Koch's statements, comes mainly from observers who, like Toussaint, Klebs, and Schuller, maintain that tuberculosis is due to a micro-organism which is a micrococcus and not a bacillus, or who, like Schottelius and others, do not admit that human and bovine tuberculosis are the same, and are, therefore, not interchangeable, which they ought to be if in both the same bacillus occurs, and if this bacillus is the *vera causa morbi*. But there can be no doubt that a vast number of competent observers have fully verified Koch's dictum, that the tubercle-bacilli are specific and different from other bacilli, except those of leprosy, as regards their chemical nature (compare their behaviour to nitric acid); and that wherever they are present in the sputum we have to deal with real tuberculosis, wherever after repeated examinations they are found to be absent there is no tuberculosis. This has by this time, although not much more than a couple of years has elapsed since Koch's first publication, become in the hands of all competent workers a matter of daily practical application, especially as regards the examination for bacilli of the sputum of patients suspected of tuberculosis.

The other equally important part of Koch's discovery, namely, the artificial cultivation of the tubercle-bacilli and the production with them of tuberculosis, has also been verified by Weichselbaum.¹ Weichselbaum also ascertained² that in acute miliary tuberculosis of man the blood contains the bacilli.

An important series of observations was published by Mr. Watson Cheyne in the *Practitioner* for April 1883, in which he proved, (1) That the organs of rabbits and guinea-




FIG. 86.—FROM A SECTION THROUGH A TUBERCLE OF THE LUNG FROM A CASE OF ACUTE MILIARY TUBERCULOSIS IN A CHILD.

Several alveoli are seen filled with debris; in the centre of this are numerous nuclei, and amongst them the tubercle-bacilli. Magnifying power about 350.

pigs suffering from the tuberculosis induced by Toussaint's cultivations from the blood of tuberculous animals, which cultivations Toussaint considered to be those of micrococci, turned out, on careful microscopic examination and suitable staining, to contain the typical tubercle-bacilli; (2) That inoculations with cultures of Toussaint's pure micrococci not containing any tubercle-bacilli did not produce tuberculosis in animals; (3) That Koch's assertions as regards the constant occurrence of the tubercle-bacilli in the tubercles of animals

¹ *Wiener med. Blätter*, 1888.

² *Ibid.* 10, 1884.

artificially tuberculised are quite correct ; (4) That material other than tuberculous does not produce tuberculosis, that is to say, that the cases of artificial tuberculosis in guinea-pigs observed by Wilson Fox and Burdon Sanderson, and in the older experiments of Cohnheim and Fraenkel, viz. those in which chronic inflammation and caseation (*i.e.* artificial tuberculosis) was thought to have been induced by other than tuberculous matter, *e.g.* by non-tuberculous caseous matter, setons, indifferent substances like bits of gutta-percha inserted into the peritoneal cavity, &c., were really due to accidental contamination with tuberculous material.

According to my own experience extending over a very large number of cases of human miliary tuberculosis and tuberculosis of cattle, I cannot for a moment accept the statement that the bacilli found in the two affections are identical ; for I find that in the two diseases their morphological characters and distribution are very different. The bacilli of human tuberculosis are conspicuously larger than those of the tuberculosis of cattle, and in many instances more regularly granular. As is seen in Figs. 83-85, those of human sputum are nearly half, or at least one-third, as large again as those of the caseous masses from the lungs of cattle.

The bacilli in the tuberculous deposits of cattle are always contained in the cells ; the larger the cell the more numerous the bacilli. This fact comes out very strikingly in thin and well-stained sections. Around many of the smaller and larger clumps of bacilli the cell-outline is still recognisable, and when the cell disintegrates, as it does sooner or later, the bacilli become free in groups ; in this respect there exists a remarkable similarity between leprosy and bovine tuberculosis. But in the human tubercles the bacilli are always scattered between the cells.

I cannot agree with Koch, Watson Cheyne, and others, who maintain that each tubercle owes its origin to the immigration of the bacilli, for there is no difficulty in ascertaining that in human tuberculosis, in tuberculosis of cattle, and in artificially induced tuberculosis of guinea-pigs and rabbits, there are met with tubercles in various stages—young and adult—in which no trace of a bacillus is to be found ; whereas in the same section cheesy tubercles may be present containing numbers of tubercle-bacilli.

Schuchardt and Krause¹ have found tubercle-bacilli, though

¹ *Fortschritte d. Med.* 9, 1883.

sparingly, in fungoid and scrofulous inflammations; Demme¹ and Doutrelepont² found the tubercle-bacilli also in the tissue of lupus. But the bacilli occurring in the lupus-tissue, as far as I am able to see, are morphologically different from the tubercle-bacilli. In a preparation made of the juice of lupus-



FIG. 87.—FROM A SECTION THROUGH THE KIDNEY OF RABBIT DEAD OF ARTIFICIAL TUBERCULOSIS.

- a. Blood-vessel filled with caseous matter, and in it numerous tubercle-bacilli.
 b. Nuclei of cells of the tuberculous new growth.
 c. Capillary vessel in cross section.

Magnifying power 700.

tissue, large transparent cells with several nuclei are found in the cell-substance of which are noticed groups of thickish short bacilli, thicker and shorter than tubercle-bacilli. These bacilli are either placed singly or in chains of two.

The so-called Bacillus of Cholera.—In the reports from India by Dr. Koch, at the head of the German Commission sent to investigate the recent outbreak of cholera in Egypt, we notice that, like the French Commission, they failed to communicate the disease to animals; that the German Commission failed to discover any specific organism in the blood of patients suffering from cholera that the intestines contained in their cavity and wall numerous peculiar "comma-shaped" bacilli, which Koch considers to have a special relation to

¹ *Berliner klin. Woch* 15, 1883.

² *Monatshefte f. praktische Dermatologie*, 6, 1883.



FIG. 88.—FROM THE SAME KIDNEY AS IN PRECEDING FIGURE.

- a. Large artery filled with caseous matter, and in it numerous tubercle-bacilli
- b. Coat of artery.
- c. Nuclei of the tuberculous new growth.
- d. A Malpighian corpuscle.

Magnifying power about 500.



FIG. 89.—FROM THE JUICE OF LUPUS-TISSUE PREPARED AFTER THE KOCH-WEIGERT METHOD OF DRYING A THIN LAYER ON A COVER-GLASS.

Magnifying power about 700.

the disease. Considering the state of the intestine in this disease, the presence of the bacilli, however peculiar, in its wall is in itself not convincing proof of their specific nature. Considering also that animals are as yet found insusceptible to cholera, artificial cultivations of these bacilli, successfully accomplished by Koch, cannot be tested.

From the artificial cultivations of these comma-shaped bacilli, Koch learned that it is necessary that the nourishing medium should have an alkaline reaction, and that the bacilli are easily killed by drying. Koch found these comma-shaped bacilli in linen soiled with the cholera dejecta, also in the water of a tank that had produced cholera in several people who had partaken of it. As soon as the bacilli disappeared from this water cholera cases ceased.

My friend Mr. A. Lingard has placed at my disposal sections through the human intestine from cases of dysentery; there are seen in the superficial parts of the necrosed mucous membrane large numbers of putrefactive bacilli. In



FIG. 90.—FROM A SECTION THROUGH THE MUCOUS MEMBRANE OF THE INTESTINE OF A PATIENT DEAD OF DYSENTERY.

A number of blood discs (extravasated into the tissue of the mucous membrane) and between them long thin bacilli.

Magnifying power 700. (Stained with methyl-blue.)

some cases, however, in the depth of the tissue there are found, amongst the extravasated blood-corpuscles, numbers of very fine, long, straight, or more commonly curved, bacilli and bacillus filaments; some are distinctly made up of a chain of long bacilli. They stain well and conspicuously in methyl-blue.

CHAPTER XII.

VIBRIO.

VIBRIONES are characterised by being rod-shaped, but not straight; they are more or less wavy; and they are motile.

(a) *Vibrio rugula* consist of rods of about 0.008 to 0.016 mm. in length, and curved either like a C or like an S. They



FIG. 91.—*VIBRIO RUGULA*
(AFTER COHN).



FIG. 92.—*VIBRIO SERPENS*,
ISOLATED (AFTER COHN).

are single, or form chains of two. Their protoplasm is always slightly granular. They are found in putrefying organic substances, and often form continuous masses, the individuals interlacing in all directions.

(b) *Vibrio serpens*.—This is also a septic organism, much thinner and longer than the previous one, more wavy, as a



FIG. 93.—VIBRIO SERPENS IN SWARMS (AFTER COHN).

rule, curved into a single or double wave. The length varies between 0.011 and 0.025 mm. It is motile; and also forms continuous masses, the individuals interlacing in all directions.

NOTE.—Many of the longer pathogenic bacilli, as bacillus of anthrax, of symptomatic charbon, of tubercle, of ulcerative stomatitis, &c., often present themselves in forms closely resembling vibrios.

CHAPTER XIII.

SPIROBACTERIUM (*Spirillum*).

SPIRILLA are filaments of a spiral shape, motile, and owing to their shape follow a spiral course when moving. They are probably capable of forming minute bright spores.

1. *Septic spirilla*.—These are found in all kinds of putrefying organic substances, and are of three kinds.



FIG. 94.—SPIRILLUM TENUE, (1) SINGLY AND (2) IN SWARMS (AFTER COHN).

(a) *Spirillum tenue*.—This is much finer and more wavy than *vibrio serpens*, the turns being closer together and spiral. Its length varies between 0.002 and 0.005 mm. ; it often forms continuous felted masses ; it is motile.

Occasionally the spirilla grow to a great length—two, three, and more of them forming a chain ; the individual spirilla are not arranged in a linear series, but folded into a zigzag. This form, which in reality is not a special kind of spirillum, is

called by Cohn¹ *spirochæta plicatilis*. The spirillum found in the tartar of the teeth is of this form, *spirochæta denticola*. But there exist all intermediate forms between a single spirillum tenue and a spirochæta. In stained specimens the construction of the spirochæta from several spirilla tenua is very distinct.

(b) *Spirillum undula* is much thicker and shorter than the former ; there are all forms between such as are only half a



FIG. 95.—SPIRILLUM UNDULA
(AFTER COHN).



FIG. 96.—SPIRILLUM VOLUTANS
(AFTER COHN).

turn to such as are of a whole turn of a spiral. It is motile and forms chains of two or more elements, occurring also in continuous masses, occasionally held together by a hyaline interstitial substance.

(c) *Spirillum volutans*.—These organisms are giant spirilla long and thick, with granular protoplasm ; 0.025 to 0.03 mm long ; motile, and with a flagellum at each end.

¹ *Beiträge zur Biologie d. Pflanzen*, vol. ii.

Pigment spirilla.

I have seen on paste a spirillum, morphologically identical spirillum undula; it is of a pale pink or rosy colour,¹ motile, and forms a kind of zooglycea, the individuals being closely placed and therefore producing a rosy colour of a more



FIG. 97.—BLOOD OF RELAPSING FEVER (HUMAN).
Blood-corpuscles and spirilla Obermeyeri.
Magnifying power 700. (After Koch.)

led tint. Where they form continuous masses, the naked eye can detect the rosy tint.

) *Spirillum sanguineum* (*Ophidomonas sanguinea* Ehrenberg).—This was observed by Cohn and Warming² in pond-water. Morphologically it is identical with spirillum volutans.

On a Rose-coloured Spirillum," *Quar. Journ. of Micr. Sci.*, vol. xv. New
Abtr. z. Biol. d. Pflanzen, vol. i.

It is motile, with a flagellum either at one or both ends. Warming occasionally saw two and three flagella at one end. It is about 0.003 mm. thick; all forms occur between such as have half and such as have two and a-half turns of a spiral. Lankester also saw the same kind of organism amongst peach-coloured bacteria.¹

3. Pathogenic spirilla.

Spirillum Obermeyer (of relapsing fever) is morphologically identical with *Spirillum tenue* (or *Spirochaeta plicatilis* of Col



FIG. 93.—BLOOD OF APE INOCULATED WITH BLOOD SHOWN IN PRECEDING FIGURE.

Blood-corpuscles and spirilla.
Magnifying power 700. (After Koch.)

It was discovered in great numbers by Obermeyer² in the blood of the general circulation in patients suffering

¹ *Quarterly Journ. of Micr. Science*, vol. xiii. New Series.

² *Centralbl. f. med. Wiss.* 10, 1873.

relapsing fever. The spirilla disappear from the blood during the non-febrile stages, gradually decreasing in numbers. They are motile; they come out well in specimens of blood made after the Weigert-Koch method of drying the blood in a very thin layer, and then staining with methyl-violet or Bismarck-brown.¹ H. V. Carter² succeeded in producing relapsing fever in monkeys by inoculation with human blood containing the spirillum Obermeyer. The blood in the monkey contained the same spirilla in great numbers. Koch has cultivated artificially the spirilla Obermeyer, and saw them growing into long spiral threads.³

¹ Weigert, *Deutsche med. Woch.* 1876; Heydenreich, Berlin, 1877.

² *Lancet*, 1879, vol. i. p. 84; and 1880, vol. i. p. 662.

³ *Deutsche med. Woch.* 19, 1879.

CHAPTER XIV.

YEAST FUNGI : TORULACEÆ, SACCHAROMYCES.

YEAST, *torula* (Pasteur), or *saccharomyces*, is not a bacterium, but belongs to an altogether different order of fungi—the *Blastomyces*. It consists of spherical or oval cells, very much larger than the largest micrococci, and as in the case of these each cell consists of a membrane and contents. The contents are either homogeneous or finely granular protoplasm; in the latter case there are generally present one, two, or more small vacuoles.

There are a great many species of *Torula*, varying from one another morphologically chiefly in their size, and physiologically by their action on various fluids (see below).

The cells multiply in suitable media by gemmation, a minute knob-like projection appearing at one side of the cell, and enlarging till it reaches nearly the size of the original or mother-cell. It finally becomes altogether constricted off from this latter, or having reached its full size remains fixed to the mother-cell, and each cell again producing by gemmation a new cell. In this way aggregations of four, six, eight, or more cells are formed, which may be arranged either as a chain when the production proceeds in a linear manner, or as a group if the gemmation takes place laterally.

Under varying conditions of growth, *e.g.* on transplanting ordinary yeast growing in sugar-containing fluids on to potato, but sometimes also in the same nutritive fluid, it is observed that some of the yeast cells enlarge twice, thrice, and more times; they then form in their interior two, three, or more small cells by endogenous formation; these new cells are regarded as spores¹—the mother-cell being an *ascospore*—and

¹ T. de Seynes, *Comptes Rendus*, 1866; Rees, *Bot. Zeitschr.* 1869; Hansen, *Carlsberg Laborat.* 1885.

become free by finally bursting the membrane of the mother-cell. On sowing these new cells into sugar-containing fluids they multiply by the process of gemmation.

Classifying them according to physiological function there are various species of torula or saccharomyces. They all have the power to split up sugar into alcohol and carbonic acid, but this power is not possessed by all to the same degree.




FIG. 99.—TORULA, OR SACCHAROMYCES.

In the lower part of the figure an ascospore and four isolated spores (after Rees) are shown.

Magnifying power about 700.

(a) *Saccharomyces cerevisiæ* (*torula cerevisiæ*). This is the ordinary yeast used in the production of beer. The individual full-grown cells vary in diameter from 0.008 to 0.01 mm.; they form beautiful long chains. They produce ascospores.

(b) *Saccharomyces vini* is very common in the air, and produces alcoholic fermentation of grape-juice; it is therefore the proper yeast of wine-production. Its cells are elliptical, slightly smaller than the former; it forms ascospores.

(c) *Saccharomyces pastorianus* is of various kinds (Hansen): in some the cells are about 0.002 to 0.005 mm. in diameter, in others larger. Some form ascospores, others do not. Most of them can be found in wine-fermentation and in cider-fermentation, but only after the first alcoholic fermentation is completed. They are very common in the air. I have sown a saccharomyces, which was contained in ordinary water, on solid nourishing media (gelatine, and gelatine and broth). It grew up copiously and formed groups of a distinct pink col

When growing in the depth of the nourishing medium it grew as a colourless torula, no ascospores were formed, multiplication taking place by gemmation only.¹

(d) *Saccharomyces mycoderma* (*mycoderma vini*). This yeast is found forming the scum or pellicle on the surface of wine, beer, and fermented cabbage (*Sauerkraut*); its cells are oval, about 0.006 mm. long and 0.002 broad. It forms chains; the ascospores are two or three times larger. It has nothing to do with the alcoholic fermentation, and is not to be confounded with *mycoderma aceti*,² which is a bacterium and the efficient cause of acid fermentation in wine and beer.




FIG. 100.—*SACCHAROMYCES MYCODERMA*, OR *OIDIUM ALBICANS*.
(After Grawitz.)

From an artificial cultivation in dilute nourishing material.

- d. Branched mycelium
- f. a. Torula stage.
- f. β. Mycelial stage.

The *saccharomyces mycoderma* does not grow well in the depth of liquids, but when sown into a liquid of acid reaction and containing but little sugar, Cienkowsky saw the cells elongating into cylindrical elements; each of which by gemmation produced a new cell which also elongated, and so on.

¹ *Quart. Journ. of Micr. Science*, 1883.

² *Wiggall*, see chapter viii. 2.

till a linear series of cylindrical cells was formed, separated from one another only by a thin septum; a mass of filaments very much resembling a mycelium was thus formed. The cylindrical cells give origin by gemmation to spherical and elliptical torula-cells.

Such a growth, in which the torula-cells are capable of forming a sort of mycelium, was formerly called *oidium*, and as *oidium albicans* is recognised as the cause of "thrush;" the well-known white patches which form on the mucous membrane of the mouth and pharynx in suckling infants and debilitated patients.

Grawitz¹ has proved by observations on artificial cultures that this fungus is identical with the *oidium* variety of *Saccharomyces mycoderma*; the cells are spherical or cylindrical, the former about 0·003 to 0·005 mm. in diameter, the latter up to 0·03 or 0·05 mm. long. As above described it forms mycelium-like filaments from which, by lateral and terminal gemmation, spring spherical or oval torula-cells. It also forms ascospores containing four to eight spores.

¹ *Virchow's Archiv*, vol. lxx

CHAPTER XV.

MOULD-FUNGI: HYPHOMYCETES OR MYCELIAL FUNGI.

Of this class of fungi only those are of special interest to the pathologist which in some way or other are connected with disease. The fungi consist of branched and septate threads or hyphæ; each filament or hypha is composed of a row of cylindrical cells, consisting of a membrane and clear protoplasm, the individual cells being separated from one another by a thin transverse septum; they increase in number by fission, and in this way the filaments increase in length. The growing ends of the hyphæ are filled, not with transparent, but with highly-refractive protoplasm. Some cells, by budding out laterally, produce cylindrical threads, which subdivide into a series of cylindrical cells, these by division and lengthening forming a new branch-hypha. The filaments form by their branches an interlacing feltwork, called thallus or mycelium. The mycelial fungi which interest us, belong to the order known to botanists as the *Ascomycetes*. They are characterised by the fact that one or other branch of the mycelial-hyphæ produces at its end a series of spherical or oval cells—the conidia-spores or *conidia*. In addition to this some of the hyphæ form peculiar large mother-cells, or *sporangia*, in the interior of which spores are formed by endogenous formation. When these sporangia are cylindrical or club-shaped, they include eight spores, and are called *asci*; the spores being *ascospores*. All conidia or spores by germination grow into the mycelial threads which become septate or subdivided into a row of cylindrical cells; these by division cause the lengthening of the mycelial threads.

(a) *Oidium lactis*.—Here the mycelium is composed of septate branched filaments of various thicknesses. Some branches

of the mycelium at their ends or laterally at a septum produce by division a series of spherical or oval conidia-spores, about 0·007 to 0·01 mm. long. These ultimately become isolated, and then germinate into a short cylindrical filament, which subdivides by transverse septa into a series of cylindrical cells ;




FIG. 101.—*OIDIUM LACTIS*.
Mycelium and spores.

these by continued growth and division give origin to the ordinary septate branch-hyphæ. The formation of conidia proceeds at the ends of these in the same manner as before. The *oidium lactis* forms a whitish mould on milk, bread, paste, potato, &c.

Favus, Herpes tonsurans, and Pityriasis versicolor of man and animals, are, according to the researches of Grawitz,¹ due to a fungus in morphological respects identical with *oidium lactis*. In favus it is known as *Achorion Schoenleini*, in Herpes

¹ *Virchow's Archiv*, vol. lxx. p. 560.

tonsurans as *Trichophyton tonsurans*, in Pityriasis versicolor as *Microsporon furfur*. Grawitz has shown by artificial cultures on gelatine, that the spherical or oval conidia germinate into shorter or longer cylindrical filaments, which by subdivision




FIG. 102.—FUNGI FROM A FAVUS-PATCH (NEUMANN).

form septate mycelial hyphæ. These and their branches give origin in their turn to spherical or oval spores or conidia. They, as well as the hyphæ, differ in size in the various species.

Malcolm Morris and G. C. Henderson,¹ on the other hand, maintain, that by artificial cultivation of the spores of *Trichophyton* in the substance of gelatine-peptone, at temperatures varying from 15° to 25° C., these grow into branched septate mycelial filaments, which by their mode of fructification are seen to be identical with the mycelium of *Penicillium*. Compare also with Babes.²

(b) *Aspergillus*.—Some of the branches of the mycelium of this fungus assume an upright position, are thicker and not at

¹ *Journal of the Royal Microscopical Society*, April 11, 1883.

² *Archives de Physiologie*, 8, 1853, p. 466.




FIG. 103.—*ASPERGILLUS GLAUCUS* (AFTER DE BARY.)

A. Hypha, the end of which c bears.
 c. The basidia.
 a.s. Ascogonium.




FIG. 104.—*E. PERITHECIUM*, HIGHLY MAGNIFIED.
 a.s. Ascogonium.
 w. Cells of the pollinodia.

all, or only slightly, septate, and at their end form flask-shaped enlargements, from which grow out radially short cylindrical cells—*basidia*; and these again at their distal or free ends produce chains of spherical spores or conidia. This is a very common mould, and according to differences in the coloration of the mycelium and spores is subdivided into different species: *A. glaucus*, *candidus*, *flavescens*, *fumigatus*, &c.

Besides this mode of spore-formation (asexual), there is another (sexual), which according to de Bary consists in this: some terminal branch of the mycelium becomes twisted like a spiral, this is considered the female organ of fructification or *carpogonium*; from the same thread branches grow towards the carpogonium; one of these branches becomes fused with the terminal portion of the carpogonium called the ascogonium, while the others—the *pollinodia*—branch and surround the carpogonium like a capsule, the whole organ is now called a *perithecium*. Finally the ascogonium by rapid division gives origin to a number of oval septate tubes, inside of which by endogenous formation numerous spores make their appearance.

Grohe¹ was the first to show that the introduction of the spores of some species of *aspergillus* into the vascular system of rabbits sometimes produces death, with symptoms of metastasis into the various organs due to localised foci, where these spores grow into mycelial filaments. Lichtheim² showed that such mycoses in rabbits cannot be produced by the spores of *Aspergillus glaucus*, but by those of *Aspergillus flavescens* and *fumigatus*. Grawitz³ studied this process more minutely, and found, that no matter whether the spores are injected into the vascular system or into the peritoneal cavity, there are established in the kidneys, liver, intestines, lungs, muscles, and occasionally in the spleen, marrowbones, lymphatic glands, nervous system, and skin, minute metastatic foci, due to the growth of the spores into mycelial filaments with imperfect organs of fructification, but no spore-formation. Grawitz thought that the spores of ordinary moulds (*penicillium* and *aspergillus*) are capable of assuming these pathogenic properties if cultivated at higher temperatures (39°—40° C.), and in alkaline media. These fungi, as is well known, grow well at ordinary temperatures and in acid media, and are then innocuous when introduced into the animal body; but by gradual acclimatisation they can also be made to grow at higher temperatures and in alkaline media, when they assume pathogenic

¹ *Berl. klin. Woch.* 1871.

² *Ibid.* 9 and 10, 1882.

³ *Virchow's Archiv*, vol. lxxxI. p. 855.



FIG. 105.—FROM A SECTION THROUGH THE KIDNEY OF A RABBIT DEAD 86 HOURS AFTER THE INJECTION OF SPORES INTO THE JUGULAR VEIN.

F. Fat droplets.

T. Tyrosin crystals.

In the upper part of the figure is a metastatic focus composed of *Aspergillus* spores and mycelium. In the lower half of the figure the urinary tubules and two Malpighian corpuscles are seen. (After Grawitz.)

properties, becoming capable of resisting the action of living tissues and of growing in them. This view has been proved to be incorrect by Gaffky,¹ Koch,² and Leber.³ Those spo

¹ *Mittheil. a. d. kais. Gesundheitsamte*, 1860.

² *Berl. klin. Woch.* 1881.

³ *Ibid.* 1862.

that do exert such pathogenic properties are not at all dependent on such acclimatisation, and are not ordinary moulds, but a distinct species of aspergillus (Lichtheim), which grows well at higher temperatures (38°—48° C.), and the spores of which under all conditions of growth are capable of producing in rabbits the mycosis in question.

(c) *Penicillium*.—In this fungus hyphæ, which are not septate, grow out from the mycelium; from the end of each of these arise like the fingers of the hand a number of short branched cylindrical cells, which give origin to chains of spherical spores.

The following two fungi belong to the order of fungi called *Phycomycetes*.

(d) *Mucor* is characterised by this, that from the mycelium hyphæ grow out which are not septate, and at the end of these a large spherical cell originates, *sporangium*, in which by endogenous formation a large number of spherical spores are developed; the wall of the sporangium giving way, the spores become free.

(e) *Saprolegnia*; colourless tubular threads, forming gelatinous masses on living and dead animal and vegetable matter in fresh water. The cylindrical or flask-shaped ends of the threads—*zoosporangia*—form in their interior numbers of spherical or oval spores—*zoospores*, possessed of locomotion (one flagellum at each pole) and which finally escape from the threads. These zoospores after some time become resting, surround themselves with a membrane, and finally germinate into a cylindrical mass which becomes transformed into the mycelium. Besides this asexual there is, however, a second or sexual mode of fructification, consisting in this: At the end of a mycelial thread a cell grows up into a spherical large ball, the *oogonium*. From the same thread, thin threads—*antheridia*—grow towards the oogonium, with the protoplasm of which they merge. This latter then differentiates into a number of spherical masses, the *oospores*, which become invested with a membrane. These become free and then germinate and grow into a mycelium. *Saprolegnia* grows on the skin of living fish, and causes here severe illness often terminating in death. Thus the salmon disease, as Professor Huxley has shown¹ is caused by this parasite. The zoospores of this salmon saprolegnia are, however, as Huxley has shown, as a rule non-motile. The hyphæ of the fungus traverse the epidermis in the diseased patches of the salmon, and they bore through the superficial layer of the derma, a stem-part being situated in the epidermis, and a root-part in the derma; each of these

¹ *Proceedings of the Royal Society*, No. 219, 1862.

elongates and branches out. "The free ends of the stem-hyphæ rise above the surface of the epidermis and become converted into zoosporangia, more or fewer of the spores of which attach




FIG. 106.—SAPROLEGNIA OF SALMON DISEASE.

A sporangium filled with zoospores; in connexion with them several young-mycelial threads.

themselves to the surrounding epidermis and repeat the process of penetration." In saprolegnia associated with the salmon-disease Professor Huxley observed only the asexual mode of fructification.

CHAPTER XVI.

ACTINOMYCES.

IN cattle there occurs a fatal disease, which is characterised by the formation of firm nodules of various sizes, due to a growth of small cells. In the centre of the nodules lie dense groups of peculiar club-shaped corpuscles—*actinomyces*—radiating from a firm homogeneous centre, and joined to this by longer or shorter, single or branched, filamentous stalks. Each of these actinomyces-corporuscles appears homogeneous, and of a bright slightly greenish lustre. These masses consist of what is called *Actinomyces* (Bollinger), and the disease is termed actinomycosis. In cattle the disease manifests itself by firm tumours in the jaw, in the alveoli of the teeth, and particularly by a great enlargement and induration of the tongue—"wooden tongue." On making sections through this latter organ there are found present in all parts microscopic tumours of small-cell growth. In the centre of each tumour is a clump of actinomyces. This clump is surrounded by a zone of largish cells, with one to four nuclei. The periphery of the tumour is made up of a fibrous capsule, with spindle-shaped cells. Occasionally the tumours are to be seen also in the skin and in the lung; in the latter organ they appear as whitish nodules, easily mistaken for tubercles.¹ Bollinger first described the disease in cattle.² Israel³ was the first to point out a disease in man characterised by metastatic abscesses (spreading it seems from a primary abscess of the jaw) in various internal organs, due to the presence of a fungus,

¹ Pflüg. *Centralbl. f. med. Wiss.* 14, 1882; Hink, *ibid.* 46, 1882.

² *Ibid.* 27, 1877.

³ *Verehow's Archiv*, vols. lxxiv. and lxxviii.

which afterwards was identified as actinomyces, and Ponfick¹ has clearly established that in man it is not a rare disease.




FIG. 107.—FROM A SECTION THROUGH THE TONGUE OF A COW DEAD OF ACTINOMYCOSIS.

A nodule is shown composed of round cells, in the centre is the clump of actinomyces surrounded by large transparent cells. Magnifying power 350.



FIG. 108.—A CLUMP OF ACTINOMYCES MORE HIGHLY MAGNIFIED, 700.

According to careful observations, Johne² succeeded transmitting the disease from cattle to cattle by inocul

¹ *Die Actinomykose des Menschen*, Berlin, 1881.

² *Deutsche Zeitschr. f. Tiermedizin*, vii. 1881.

but not by feeding. He also found¹ in twenty out of twenty-one healthy pigs examined, the actinomyces present in the crypts of the tonsils.

Israel² succeeded in transmitting the disease to a rabbit by inserting into the peritoneal cavity a piece of a human actinomyces-tumour.

R. Virchow quite recently,³ in conjunction with O. Israel and Duncker, ascertained that pork occasionally contained whitish chalky nodules, larger than those due to trichinæ, and containing in their interior the actinomyces.

O. Israel⁴ claims to have succeeded in artificially cultivating the actinomyces on solid ox-serum ; in fluid media the growth does not succeed, owing to the swelling up and death of the actinomyces-corpules.

¹ *Centralbl. f. med. Wiss.* 15, 1881.

² *Ibid.* xxvii. 1883.

³ *Virchow's Archiv*, vol. xcv. p. 544

⁴ *Ibid.* vol. xcv. p. 112.

CHAPTER XVII.

ON RELATIONS OF SEPTIC TO PATHOGENIC ORGANISMS.¹

THERE is hardly any question which to the pathologist and sanitary officer can be of greater importance than the relation of septic to pathogenic organisms. To the pathologist the life history of a micro-organism, outside and within the animal body, must ever remain an important field of inquiry; to the sanitary officer all conditions affecting the life and death of those organisms which produce, or at least are intimately bound up with, infectious diseases, such as the distribution and growth of these micro-organisms outside the animal body, the agencies which affect it in a favourable and unfavourable sense, are the points which he has particularly to consider in dealing with the spread and prevention of infectious maladies. Now, it is known of many micro-organisms, both those that are associated with putrefactive processes as well as those that are bound up with infectious disease, that temperature, the medium in which they grow, presence and absence of certain chemical compounds are capable of materially affecting them. I need not for this purpose enumerate all that is known already by direct experiment, but will only limit myself to reference to the researches of Schröter, Cohn, and Wernich on that group of micro-organisms known as pigment bacteria, i.e. bacteria which only under certain conditions, notably temperature and soil, produce definite pigments (Cohn's *Beiträge zur Biologie d. Pflanzen*); to those of Hansen (Carlsberg Laboratory) on yeast; to those of Neelsen on the bacilli producing the blue colour of milk, the bacillus syncyanus (*Beitr. zur Biol. d. Pflanzen*, iii. 2, p. 187); to the works of Toussaint, Pasteur, Chauveau, Koch,

¹ The greater part of this chapter is copied from an interim report by myself to the Medical Officer of the Local Government Board, 1884.

and others on the bacillus anthracis ; Arloing, Thomas, and Cornevin on the bacillus of symptomatic charbon ; of Koch on the bacillus of tuberculosis ; of Israel on actinomyces, and many others ; and particularly would I refer to the many valuable suggestions and considerations expressed by v. Nägeli in these respects in his book, *Die niederen Pilze*, München, 1877 and 1882.

While from these observations it would appear that both septic and pathogenic micro-organisms are capable of suffering some modifications in their morphological and physiological behaviour, it is nevertheless still an open question whether an organism which under ordinary conditions is only associated with putrefactive changes in dead organic material, and which cannot under these ordinary conditions grow and multiply within the living body, can, under certain extraordinary circumstances, become endowed with the power of growing and multiplying within the body of a living animal, creating there a pathological condition, inducing there an infectious disease.

Three distinct septic micro-organisms have, after numerous experiments and careful observations, been mentioned, as being capable when growing under certain extraordinary conditions of assuming pathogenic properties. These three organisms are : (A) the common bacillus of hay infusion is said by Buchner to be capable of transformation into bacillus anthracis ; (B) a bacillus subtilis, present in the air, which, although quite harmless in itself, assumes distinct pathogenic properties when growing in an infusion of the seeds of *Abrus precatorius*, becoming hereby endowed with the power of causing severe ophthalmia (Sattler) ; (C) a common mould, aspergillus, which harmless in itself, when grown on neutral and alkaline material at about body-temperature (38° C.) assumes, according to Grawitz, very poisonous properties, producing in rabbits inoculated with it death, with metastasis of aspergillus and its spores in the various internal organs.

There are in the literature of micro-organisms other cases mentioned, in which such a transformation has been *supposed*, but without any experimental proof, and we need not therefore trouble ourselves more about them.

Let us now review *seriatim* the above three cases :

(A) Dr. Hans Buchner in a paper, which for many reasons may be considered an important one, "Ueber d. experim. Erzeugung des Milzbrandcontagiums, &c.," published in the *Sitzungsberichte d. math. physik. classe d. k. Bair. Akademie d. Wiss.* 1880, heft iii. p. 369, states that he succeeded in

transforming the common bacillus of hay infusion, the hay bacillus, into the bacillus anthracis.

The hay bacillus and the bacillus anthracis rank together morphologically under that form which Cohn has named bacillus subtilis.

The two are, however, not quite identical in morphological respects. The hay bacillus is a minute rod or cylindrical-shaped bacillus, which by elongation and division produces chains and further threads just like the bacillus anthracis, but in the latter (*i.e.* bacillus anthracis) the bacilli and their threads are composed of cubical elements, as is shown in stained specimens, and as has been mentioned in a former chapter, whereas in those of the hay bacillus the elements are rods or cylinders. I have seen, however, many of the short hay bacilli which being constricted, *i.e.* in the act of division, appear as two short more or less cubical elements placed end to end. It is generally assumed that in hay bacillus the bacilli are always rounded at their ends, whereas the bacilli anthracis are as if straight cut at their ends; but this is not universally the case, since I have seen in cultures the bacilli anthracis with distinctly rounded ends. But, speaking generally, the hay bacillus is a rod more distinctly rounded at its ends, the bacillus anthracis of the blood is not so.

The bacillus anthracis is slightly thicker than the hay bacillus. In artificial cultivations carried on in neutral broth the bacillus anthracis is about twice as thick as the hay bacillus growing in the same fluid, and when both are growing in neutralised hay infusion the two are very conspicuously different from one another, and can at a glance be distinguished from one another; the hay bacillus being about half the thickness of the bacillus anthracis. In stained specimens, too, the latter is beautifully made up of a row of cubical cells, whereas the former consists of cylinders only.

True, the bacillus anthracis is not always of the same thickness, for, as I have shown, when growing in neutral pork broth it is decidedly thicker than in the blood of an animal dead of anthrax. And also in the blood of different animals the bacillus anthracis slightly varies in thickness, for in the guinea-pig's blood it is slightly thicker than in that of the rabbit or sheep.

The hay bacillus is motile, possessed of a flagellum, and therefore capable of locomotion; the bacillus anthracis is not motile. I am quite aware that Cossar Ewart (*Quarterly Journal of Microscopical Science*, April 1878) states that he has seen in a specimen kept artificially heated under micro-

scopic observation that the bacillus anthracis, at first non-motile, is capable of becoming motile. At one or both ends a flagellum grows out from its body. But this observation is unreliable, since Ewart did not guard himself in any way from the accidental introduction of septic bacilli, many of which are motile. Besides, he says of the bacilli, which he figures as anthrax bacilli, that they are connected with one another by two fine threads, and that they probably separate from one another and each retains one filament, which is its flagellum. But his observations, so far as they have application to anthrax bacilli, are capable of quite a different interpretation. In every specimen of blood and in every artificial culture bacilli can be seen, in which at one place or more the protoplasm is wanting, owing as I have shown, to degeneration; in such places only the empty sheath is present, and of course in fresh specimens this gives the appearance as if the two protoplasmic portions of the bacillus were connected with one another by two fine threads, i.e. the sheath being transparent is seen here edgewise.

In no instance has the bacillus anthracis been observed to be motile. I have examined thousands of specimens of fresh bacillus anthracis in the blood and in artificial cultures, and I have never seen anything that in the least would lead me to differ from this proposition.

As regards the spores they are of the same aspect and size in both the hay bacillus and bacillus anthracis. The threads in good cultures form in both cases the same bundles more or less twisted and forming convolutions, but in certain cultures of the bacillus anthracis, e.g. in broth, in which the growth is limited to the bottom of the fluid, the convolutions and the twisted condition of the threads are more pronounced, more cable-like.

Hay bacillus being motile, every culture of it is uniformly turbid, the bacilli being capable of moving about, and after a day or two of incubation at 35° C. they form a distinct pellicle on the surface of the fluid, which in further stages becomes complete and thick. These pellicles are composed of a dense feltwork of threads of the bacilli, and in them spore-formation is going on.

By shaking the fluid the pellicle sinks to the bottom, and if the fluid is not exhausted yet, a new pellicle is formed of the same nature.

In no culture of hay bacillus are there ever observed those cloudy, fluffy, whitish and transparent convolutions that are *seen in cultivations* of bacillus anthracis carried on at the

bottom of fluid broth, and which have been so accurately described by Pasteur.

Both the hay bacillus and the anthrax bacillus when growing on gelatine mixtures liquefy the gelatine; both when growing in meat broth turn the at first colourless fluid in the course of incubation to an amber, and further to a brown, tint.

The hay bacillus is capable of thriving well in acid solutions, it grows copiously in hay infusion, which is of a distinct acid reaction; the bacillus anthracis, although capable of making a slight progress in acid hay infusion, does not get far, for degeneration soon sets in; it thrives best in neutral solutions. Hay bacillus thrives also very well in neutral solutions.

Buchner states, that by successive cultivation of bacillus anthracis under *constant variation of the nutritive material*, he saw it assume gradually the properties of hay bacillus. Thus he saw that its mode of growth gradually changed, inasmuch as instead of forming, as the typical bacillus anthracis does, fluffy convolutions at the bottom of the fluid-nourishing medium, it gradually showed a tendency to stick to the glass and to the surface of the fluid, and to form a sort of pellicle just like the hay bacillus does. This I consider to be an erroneous interpretation of an easily explained and simple fact. It does not want any of the many successive generations of bacillus anthracis, in which Buchner says he has achieved this transformation, it simply requires two nourishing fluids, in both of which the bacillus anthracis will thrive well, but which fluids differ in specific gravity. Let Buchner do as I have done, let him take two test-tubes, both containing sterile broth, but in one the broth concentrated, in the other dilute. Let him inoculate the two test-tubes with bacillus taken from the same blood, say of a guinea-pig dead of anthrax, let him place them in the incubator at a temperature of 35°-42° C. After two or three days, and more decidedly later, he will notice this very difference in the aspect of the cultures that he lays so much stress on as indicating a change in the physiological character of the bacillus. One test-tube, containing the dilute broth, shows the typical fluffy convolutions at the bottom of the fluid; while the other, containing concentrated broth, shows a distinct attempt at the formation of a pellicle. Let him now take out a droplet from this second test-tube and inoculate with it two test-tubes of the same nature as above, *i.e.* one containing concentrated broth, the other dilute broth. After two or three or more days of incubation he will find exactly the same differences as above.

Buchner states that the bacillus anthracis when carried

through a large number of successive cultures, at a temperature of 35° — 37° C., gradually loses its pathogenic properties. Of this assertion I have said already a great deal in my Report for 1881—1882, and I mention it here merely in connexion with Buchner's other assertions. I have shown, that even assuming that Buchner has had in all his cultures the true bacillus anthracis, but for which there is no definite proof, as Koch has so ably pointed out in his critical review of Buchner's work (*Mittheilungen aus dem k. Gesundheitsamte*, Berlin, 1881, Bnd. I.), Buchner, having tested his cultures on white mice only, has fallen into a serious error, for, as I have shown (Reports for 1881—1882), a culture of bacillus anthracis may have become quite harmless to white mice, but be still very virulent to other animals. In fact, therefore, Buchner's result does not require for its achievement more than one culture, provided this has been kept for several days or weeks without spore-formation, as was the case in Buchner's experiments.

As regards Buchner's statement that by successive cultivation of bacillus anthracis at 35° — 37° C., this assumes the morphological and physiological characters of hay bacillus, I agree with Koch in regarding this as a complete error. If the cultures are quite safe from contamination nothing of the sort ever happens. I have now for several years carried on such cultures, and have not seen anything of the sort. It is of course clear that if by any accidental contamination, say at the time of inoculating a fresh tube, a motile septic non-pathogenic bacillus, with which, or with the spores of which, the air sometimes abounds, is introduced, every new culture established from this one will abound in this bacillus, and as it grows quicker and more easily than the bacillus anthracis, the next cultivations become barren of all the bacillus anthracis, and only the non-pathogenic motile bacillus will be found present. This criticism has been applied by Koch to Buchner's experiments, and I must fully endorse it.

But there is a much more serious statement of Buchner's—serious, because if true in nature, it is dreadful to contemplate to what amount of anthrax man and brute may become subject—viz., that he maintains to have succeeded in transforming the hay bacillus into bacillus anthracis, by carrying the former through many generations under ever varying change of soil. It is needless to detail here all these experiments of Buchner, since I do not attach any great value to them, and I should not have troubled myself much about them were it not that one meets in mycological literature, particularly on *the part of botanists*, an acceptance of Buchner's statement

that hay bacillus can change into the pathogenic bacillus anthracis (see Zopf, *Die Spaltpilze*, Breslau, 1883).

I have repeated Buchner's experiments on rabbits, guinea-pigs, and white mice. I have grown the hay bacillus in various kinds of broth, in gelatine broth mixtures, in hydrocele fluid, in peptone fluid, in Agar-Agar and peptone, at temperatures varying between 30° and 38° C., and I have, to put it shortly, never seen that it shows the least tendency to change its morphological characters, or that it ever assumes any morphological or physiological character like the bacillus anthracis. I consider this a perfectly hopeless task, and I feel sure any one might as soon attempt to transform the bulb of the common onion into the bulb of the poisonous colchicum.

But Buchner states that with his cultures of hay bacillus, carried through many generations under varying conditions of soil, he inoculated white mice, which died under symptoms of anthrax, and whose blood contained the typical bacillus anthracis. I do not for a moment doubt that he really had mice dying from anthrax after inoculation with cultures of hay bacillus, but I question the admissibility of his interpretation. I believe that some accidental contamination of the culture of hay bacillus with anthrax spores or otherwise may have occurred and have got overlooked. How liable one kind of infective material is to be invaded by foreign infective matter may be understood from the following examples of its actual occurrence.

It is now admitted on all hands that the results of Villemin in producing what is called artificial tuberculosis in guinea-pigs, by inoculating the animals subcutaneously with cheesy matter derived from human and bovine tuberculosis or from a guinea-pig suffering from artificial tuberculosis, cannot be produced by any other means; it cannot be produced by ordinary, *i.e.* non-tubercular cheesy or other pus,¹ nor by setons (as once thought by Wilson Fox and Sanderson) setting up chronic caseous inflammations in the skin of guinea-pigs, nor by chronic mechanical irritation, *e.g.* insertion into the peritoneal cavity of bits of gutta-percha or other substances producing chronic peritonitis (as was thought by Cohnheim and Frænkel), but, as Cohnheim now tersely puts it, tuberculosis can be produced only by matter derived from a tuberculous source, and anything that produces this tuberculosis is derived from a tuberculous source. Dr. Wilson Fox, after the very important experiments performed by Dr. Dawson Williams, according to which chronic inflammation in th

¹ Compare Watson Cheyne, *Practitioner*, April 1883.

skin of guinea-pigs produced by setons, is in no case followed by tuberculosis, has conceded that in his earlier experiments there must have entered some error in the use of the materials. Cohnheim has conceded the same. It is clear that these observers, while working at the same period with both tuberculous and non-tuberculous matter, must have had, in the course of experiments with the latter substance, accidental contamination with the former, and hence had the guinea-pigs inoculated by them with non-tuberculous matter nevertheless affected with tuberculosis. Dr. Williams, who had no contamination to fear, working with non-tuberculous matter only, had consequently no accidental contamination. This shows us how dangerous, as regards reliability of results, it is to work in one laboratory with different infective materials at the same period.

I have myself experienced some very curious results bearing on this very point. During the last year I have seen the following cases of accidental contamination occur. I work in the laboratory of the Brown Institution, which comprises a suite of rooms. Although working extensively on anthrax, I generally limit myself to one room only. A friend of mine, who one day injected into a vein of a guinea-pig blood taken from a blood-vessel of a dog suffering from distemper, found, to his great disappointment, the guinea-pig dead after two days under the typical symptoms of anthrax, the blood of this animal teeming with the characteristic bacilli. The hypodermic syringe used in this experiment for injection had not been previously used by me in my anthrax experiments, since I never use a syringe in my inoculations, but only glass pipettes freshly made and drawn out into a fine tube. The experiment was performed in the room adjoining the one in which my anthrax investigations were being carried on, but I was in the habit of making every day a good many specimens of anthrax cultivations and spores, so that there must have been a good many of these spores distributed on the table and floor, and probably found their way into the wound of the guinea-pig at the time the above experiment was made.

Another gentleman working in the laboratory of the Brown Institution intended to inoculate several guinea-pigs with human tubercles. For this end he mashed up in saline solution, in a clean mortar, a bit of human lung studded with tubercles. He did this in my room on the same table on which I was working with anthrax. One of these guinea-pigs, inoculated with human tubercle, died before the second day was over of typical anthrax. Its blood was teeming with the *bacillus anthracis*. Such an accidental anthrax of a

guinea-pig inoculated with tuberculous matter occurred several times. In all cases freshly drawn-out glass capillary pipettes had been used for performing the inoculation, and also the other instruments had been carefully cleaned before the inoculation.

I myself had the following accidental contaminations :—

A guinea-pig had been inoculated with a culture of bacillus anthracis, which I did not expect would produce anthrax, the culture not being capable of starting new cultures, the bacillar threads being all in a state of degeneration. The animal, of course, remained unaffected. Some weeks afterwards inspecting the guinea-pig, to my surprise, I found the inguinal lymphatic glands at the side of the former inoculation greatly swollen, filled with cheesy pus. The animal was killed, and was found to be affected with general tuberculosis, the cheesy matter of the tubercular deposits containing the tubercle bacilli. Comparing my notes on this animal with those of my friend Lingard, we found that on the very day on which I inoculated the animal with my anthrax culture we had inoculated several other guinea-pigs with tuberculous matter. This tuberculous matter was prepared in the same room in which I prepared the fluid for my anthrax inoculation, but the instruments in the two sets of experiments had not been the same.

A rabbit was inoculated with a culture of bacillus anthracis which I did not expect would produce anthrax. The animal remained unaffected with anthrax, but died after four weeks with the symptoms of extremely well-marked tuberculosis—in fact, the best marked case that I have seen—of both lungs, spleen, liver, and kidney. The tubercular deposits contained the tubercle bacilli.

Also in this instance inoculations with tuberculous matter had been going on at the same time, when I meant to have inoculated nothing else but a culture of anthrax bacilli.

I think all these facts taken together prove unmistakably that working with two contagia in the same laboratory and at the same period, accidental contamination is of no rare occurrence. And this applies with equal force to Buchner's experiments. Buchner worked extensively with anthrax cultures in the same laboratory, and at the same time he had those successful cases of anthrax in mice which he thought to have inoculated with cultures of hay bacillus, and accidental contamination probably was the result. Buchner himself has experimentally shown that anthrax virus in the shape of spores can by inhalation produce anthrax, and, therefore, this is another argument against his above cases of positive result. I am assuming that his cultures of hay bacillus were real

free of spores of bacillus anthracis; but, seeing that his anthrax cultures were probably contaminated with hay bacillus, I do not see why, by some chance, one of his tubes which he thought he inoculated with hay bacillus should not have been accidentally contaminated with the spores of bacillus anthracis, of which there must have been many about in the air of the laboratory.

If Buchner could show us that in a laboratory, in which for some considerable time anthrax cultures, anthrax animals, and examinations of anthrax bacilli had not been carried on, cultivation of hay bacillus ultimately yields a fluid which produces typical anthrax, then I should be perhaps prepared to concede his proposition of a transmutation of hay bacillus into bacillus anthracis. Such a proposition is of the widest importance, and therefore its proof ought to be beyond cavil, there ought to be no chance of a possibility of error. Such proof Buchner has not given, and I cannot therefore accept his interpretation.

(B) The second instance in which the transformation of a common septic into a specific or pathogenic organism has been experimentally achieved, or I should rather say has been stated to have been achieved, is the jequirity bacillus. In 1882 the well-known ophthalmologist M. L. de Wecker in Paris drew attention to the therapeutic value of the seeds or beans of *Abrus precatorius*, a leguminosa common in India and South America. The people of Brazil use it under the name jequirity as a means to cure trachoma or granular lids. De Wecker after many experiments found that a few drops of an infusion made of these seeds causes severe conjunctivitis, in the course of which, no doubt, trachoma is brought to disappearance and cure, and it is accordingly on the Continent and in this country now used for this therapeutic object. [I am informed by my friend Dr. T. Lewis, formerly of India, now pathologist at the Netley Army Medical School, that the people in some parts of India know the poisonous properties of these seeds, and use it for inoculating with them subcutaneously, cattle; in consequence a severe inflammation is set up, and the animals die of some sort of septicæmia. This is done for the sake of simply obtaining the hides of the beasts.]

Sattler, in a very important and extensive research (*Wiener medic. Wochenschrift*, N. 17-21, 1883, and *Klin. Monatsbl. f. Augenheilk.*, June 1883) ascertained that when an infusion of the jequirity seeds is made of the strength of about half per

cent., this infusion after some hours to a few days contains numerous bacilli, motile, capable of forming spores, and in all respects identical with a bacillus subtilis. The bacilli are about 0.00058 mm. thick, and from 0.002 to 0.0045 mm. long. They form a pellicle on the surface of the infusion, and in the bacilli of this pellicle active spore formation is going on. The bacilli grow and multiply well at a temperature of about 35° C., but also, only slower, at ordinary temperature. Sattler cultivated artificially the bacilli on blood-serum gelatine and meat-extract peptone gelatine, both solid media, and continued their growth through several successive cultivations. Both the infusions of the jequirity and the bacilli taken from these artificial cultures inoculated into the conjunctiva of healthy rabbits produce severe ophthalmia, leading to the production of great œdematous swelling of the conjunctiva and eyelids, and temporary closure of the latter, and to the secretion of purulent exudation. Both the exudation and the swollen lids are said to contain infective bacilli and their spores. Sattler ascertained by many experiments, that none of the bacilli and the spores distributed in the atmosphere had those specific properties, viz., to excite ophthalmia, as long as they grow in other than jequirity fluid, but having had access, *i.e.* having entered the jequirity infusion, assume here this specific power. There is no doubt that Sattler worked the whole problem with great care, worked out all points connected with it in great detail, and for this reason his work was considered to have for the first time unmistakably established that a harmless bacillus, owing to the particular soil in which it grew, assumes definite specific or pathogenic properties. To me this jequirity bacillus had a great interest, since I was particularly anxious to get hold of such an organism, in order to see whether and how far it can again be made harmless. For if ever there was a good case, a case in which a previously harmless micro-organism had by some peculiar conditions become specific, this was a case; and therefore it must be here possible by altering its conditions of life again to transform it into a harmless being. The theoretical and practical importance of such a case must be evident to every one who has at all devoted any thought to the relation of micro-organisms with disease. The whole doctrine of the infectious diseases, I might almost say, is involved in such a case, for if in one case it can be unmistakably proved that a harmless bacterium can be transformed into a pathogenic organism, *i.e.* into a specific virus of an infectious malady, and if this again can under altered conditions resume its harmless property, then we should :

once be relieved of searching for the initial cause in the outbreak of an epidemic. But in that case we should be forced to contemplate, as floating in the air, in the water, in the soil, everywhere, millions of bacteria which, owing to some peculiar unknown condition, are capable at once to start any kind of infectious disorder, say anthrax (Buchner), infectious ophthalmia (Sattler), and probably a host of other infectious diseases, and thus to form the starting-point of epidemics. And the only redeeming feature, if redeeming it can be called, in this calamity would be the thought that the particular bacterium would by and by, owing to some accidental new conditions, again become harmless.

These were the reasons, and good reasons I think they were, which prompted me to inquire into the jequirity bacillus and jequirity ophthalmia, and after a very careful and extensive series of experiments, to be described presently, I have proved beyond any doubt that the jequirity bacillus, *per se*, has no more power to create an infectious ophthalmia than Buchner's hay-bacillus had of creating anthrax.

The following experiments prove this conclusively :—

The seeds of jequirity (*Abrus precatorius*) are crushed and powdered, the perisperm is removed, and of the rest an infusion is made of about the strength of half per cent. with distilled water, previously boiled and contained in a flask previously sterilised (by heat) and plugged with sterile cotton-wool. The infusion is made while the water is still tepid. After half an hour the infusion is filtered into a fresh sterile flask, plugged with sterile cotton-wool, the access of air being limited as much as possible. This is effected by keeping the cotton-wool in the mouth of the flask around the end of the glass filter. The filtered fluid is of a slightly yellowish-green colour, and is almost neutral and limpid. A small quantity is withdrawn with a capillary glass pipette freshly drawn out, and from this several test-tubes containing sterile nourishing material (peptone solution, broth, Agar-Agar and peptone) are inoculated; and from the same pipette, and at the same time, several eyeballs of healthy rabbits are inoculated, by placing a drop or two of the infusion under the conjunctiva bulbi. The test-tubes are placed in the incubator and kept there at 35° C. After twenty-four hours all eyeballs are intensely inflamed, the eyelids closed and swollen, and a large amount of purulent secretion is present in the conjunctival sac, but all the test-tubes remain perfectly limpid; no growth has made its appearance, and they remain so.

In a second series the infusion prepared in the above manner

is used fifteen minutes after it is made and used as above, for inoculation of test-tubes and eyeballs. The fluid in the test-tubes after incubation remains limpid, the eyeballs all become inflamed. In both series the amount of fluid inoculated into the test-tubes is more than twice as great as that injected into the eyeballs. From this it is quite clear that the fluid used for inoculation of the test-tubes was barren of any micro-organisms, and nevertheless it possessed a powerful poisonous principle. I do not mean to say that the infusion as a whole contained in the flask contains no organisms, but that the small quantity of the fresh infusion that was used for the inoculation of the test-tubes and eyeballs contained none is absolutely certain. When such a flask is placed in the incubator, after twenty-four to forty-eight hours or later there are found in it large quantities of bacilli, the spores of which must have entered from the air during the process of preparing the infusion. The bacilli are such as described by Sattler; they soon form spores in the usual way. Such an infusion is very poisonous, just like the fresh one. Sattler has shown, and this is easily confirmed, that the spores of these bacilli stand boiling for a few minutes without losing their power to germinate. Consequently, if such a poisonous infusion full of bacilli and spores be boiled for half a minute the spores are not killed; proof for this: that if with a minute dose of this spore containing boiled infusion any suitable sterile nourishing material in test-tubes be inoculated, and then these test-tubes be placed in the incubator at 35° C., after twenty-four to forty-eight hours the nourishing fluids are found teeming with the jequirity bacilli; *but no amount of this material produces the least symptom of ophthalmia. Every infusion of jequirity loses its poisonous activity by boiling it a short time, $\frac{1}{2}$ to 1 minute, and hence the above result.*

In this respect the poisonous principle of jequirity infusion comports itself similar to the pepsin ferment, which, as is well known, is destroyed by short boiling.

If an infusion is made as above, and after fifteen minutes it is filtered and then subjected to boiling for $\frac{1}{2}$ -1 minute, it will be found to have become absolutely non-poisonous, but not sterile, placing it in the incubator after twenty-four to forty-eight hours, vast numbers of the jequirity bacillus are found in it. But no amount of this fluid is capable of producing the slightest symptom of ophthalmia.

A large per-centage of the rabbits, whose conjunctiva has been inoculated with the fresh unboiled poisonous infusion, die after several, three to eight, days. The eyeballs and

eyelids are intensely inflamed, as stated above, the skin and subcutaneous tissue of the face, neck, chest, and even abdomen, is found enormously œdematous, the pericardium, pleura, lungs, and peritoneum very much inflamed, their cavities filled with a large quantity of exudation. The exudations of the conjunctiva, pericardium, peritoneum, the œdematous skin and subcutaneous tissues contain no infectious property, and as a rule no bacilli or spores of any kind, if examined in the living animal or immediately after death.

There is one point which requires careful consideration, it is this : Sattler states that he has cultivated the bacillus, taken from a poisonous jequirity infusion, through several successive generations on solid material, and with the new cultures he was able to produce the jequirity ophthalmia. I have no doubt whatever that this is really the case, but it bears an interpretation different from the one Sattler gave it. Sattler, and most pathologists, would, of course, say this : if any micro-organism taken from a soil that possesses infective properties be carried through many successive artificial cultivations, all accidentally adhering matter would hereby become so diluted that it may be considered as practically lost ; that is to say, the organisms of the further generations have become altogether free of that matter. If the organisms of these further generations still possess the same poisonous property as the original material, then we must conclude that this poisonous principle is identical with the micro-organism. I do not agree with this whole chain of propositions, although I agree with some parts. If a micro-organism be carried through several successive cultivations in a *fluid medium*, always using for inoculation of a new culture an infinitesimal dose, and as nourishing medium a comparatively large quantity of fluid, then, no doubt, carrying on the cultivations through four, five, or six successive cultures, any accidentally adhering original matter becomes practically lost, and if then the organism still possesses the same poisonous action to the same degree as the original material, then no doubt the conclusion that organism and poison are in this case identical becomes inevitable. But this is not the case with the jequirity bacillus. Taking from a poisonous jequirity infusion full of the bacilli one to two drops, and inoculating with it a test-tube containing about four to five cc. of nourishing material, and using this at once *without previous incubation*, we find that even a few drops of this so diluted fluid still possess poisonous action. Precisely the same result is obtained when taking from a perfectly fresh *jequirity infusion*, *i.e.* before any organisms have made their

appearance, one to two drops, and diluting them with four to five cc. of distilled water, and using of this diluted fluid one to two drops for inoculating the conjunctiva of healthy rabbits, severe ophthalmia will be the result. Carrying on the cultivation of these bacilli started from a poisonous infusion, for a second generation in fluid medium, no trace of any poisonous action can be now detected, any quantity of such a cultivation is incapable of producing ophthalmia. Sattler used in his cultivations solid nourishing material, on the surface of which he deposited his drop of poisonous jequirity infusion containing the bacilli; after some days' incubation, the bacilli having become greatly multiplied, he took out from this second culture a drop, and transferred it to a new culture tube of solid material, and so he went on; every one of these cultures possessed poisonous action. Clearly it would, since he always used part of the original fluid deposited on the surface of the solid nourishing material. Part of this (being gelatine) became by the growth liquefied, but considering that Sattler started with infusions of considerable concentration—he left the seeds for many hours and days in the infusion—it is not to be wondered at that this would bear a considerable amount of dilution, and still retain its poisonous properties. From all this we see, then, that the jequirity bacillus *per se* has nothing to do with the poisonous principle of the jequirity seeds, but that this principle is a chemical ferment in some respects (in its inability to withstand boiling) similar to the pepsin ferment.¹

(C) The third case, in which an experimental attempt has been made to transform a common septic into a specific or pathogenic micro-organism, is exemplified by the common mould, *aspergillus*, a mycelial fungus. But since this point has been discussed already in Chapter XV. I need not here enter into it again, suffice it to say that certain species of *aspergillus* possess the power of making in the rabbit a general mycosis, and this power they possess *ab initio*; other kinds of *aspergillus* do not possess this character and cannot acquire it under any conditions.

¹ Since this has been in print, I became aware that Messrs. Warden and Waddell published in Calcutta during the present year a most valuable memoir, detailing a large number of observations on the jequirity poison, which are in complete harmony with my own observations. They have definitely proved, that the active principle is a proteid—*abrin*—closely allied to native albumen; that its action is similar to that of a soluble ferment, that it can be isolate and that it is contained, not only in the seeds but also in the root and stem *Abrus precatorius*

Thus also this third case of a transformation of a common into a specific organism due to altered conditions of growth falls to the ground.

It might be now asked, how about those cases in which by injection of very small quantities of putrid organic substances, pyæmia or septicæmia, has been produced in rodents? Take the case of Davaine's septicæmia in rabbits. This disease has been produced in rabbits by Davaine, Coze and Feltz, and by many other observers, by injecting into the subcutaneous tissue of healthy rabbits small quantities of putrid ox's blood. The rabbits die in the course of a day or two, and their blood is found teeming with minute organisms, which prove to be bacterium termo; every drop of this blood possesses infective properties; when inoculated into a rabbit it produces septicæmia with precisely the same appearances as before. Pasteur and Koch have succeeded in producing septicæmia in mice and rabbits, and especially in guinea-pigs, by inoculating them subcutaneously with garden earth or with putrid fluid. This is Pasteur's septicæmia, or Koch's malignant œdema; it is characterised by œdema at the seat of inoculation, and spreading hence in the subcutaneous tissue of the surrounding parts. The animals die generally in twenty-four to seventy-two hours.

Koch has produced by injection of small quantities of putrid fluids into the subcutaneous tissue of mice a peculiar septicæmia; the animals sometimes die in forty to sixty hours, and the white corpuscles of the blood are found crowded with exceedingly minute bacilli. Koch succeeded also in producing a pyæmia in rabbits by injection of putrid fluids, and this pyæmia is characterised by zooglœa of minute micrococci being present in the blood-vessels. Further, a progressive necrosis in mice by inoculating them with putrid fluids, the necrosis being due to the growth of micrococci and spreading from the seat of inoculation, and destroying as they spread all the elements of the tissue. All these cases have been minutely described by Koch in his classical work, *Die Aetologie der Wundinfectionskrankheiten*, Leipzig, 1879. I have in addition mentioned in Chapter VII. § 13 a micrococcus causing abscess and pyæmia in mice.

Now do these cases prove that septic micro-organisms, living and thriving in putrid organic fluids, can, when introduced into the body of animals, owing to some peculiar unknown condition, so change as to produce a fatal infectious disease?

av, with Koch, who has very ably discussed all these
2. Those organisms which are connected with the

above morbid processes possess this pathogenic power *ab initio*, not due to any peculiar condition of growth.

Amongst the legion of different species of micrococci and bacilli occurring in putrid substances, the great majority are quite harmless; when introduced into the body of an animal they are unable to grow and to multiply, and therefore are unable to produce any disturbance. But some few species there are which, although ordinarily growing and thriving in putrid substances, possess this power, that when introduced into the body of a suitable animal they set up here a specific disease.

One of the best studied cases is that of the bacillus anthracis. This organism is capable of growing well and copiously outside the body of an animal, it thrives well wherever it finds the necessary conditions of temperature, moisture, and nitrogenous material; when it finds access into the body of a suitable animal it produces the highly infectious fatal malady known as anthrax. The micrococcus of erysipelas is now well known through the admirable researches of Fehleisen to be capable of existence and multiplication outside the animal body; it grows well in artificial cultures, so does the tubercle bacillus of Koch, so does the bacillus which I described of swine-plague, mentioned in a former chapter, and so do other micro-organisms. Davaine's septicæmia in rabbits, Koch's septicæmia in mice, &c., &c., cannot be produced by every putrid blood or putrid organic fluid, only by some, only now and then, *i.e.* when the particular micro-organism capable of inducing the disease is present in those substances, and then only when it finds access into a suitable animal. Davaine's septicæmia of rabbits cannot be induced in guinea-pigs, Koch's septicæmia of mice cannot be induced in guinea-pigs, anthrax bacilli cannot induce the disease in dogs, and so with the other micro-organisms.

We conclude then from this that some definite micro-organisms, although as a rule existing and growing in various substances of the outside world, have the power when finding access into the body of a suitable animal to grow and thrive here also, and to induce a definite pathological condition. But this power they have *ab initio*. Those that do not possess this power cannot acquire it by any means whatever. Just as there are species of plants which act as poisons to the animal body, and other species of plants which, although belonging to the same group and family, and although very much alike to the others, have no such power and cannot acquire such a power by any means, so there are micro-organisms which

are pathogenic while others are quite harmless. The latter remain so no matter under what conditions and for how long they grow.

I have made a series of experiments with the view to obtain pure cultivations of definite septic micro-organisms: various species of micrococci, bacterium termo, and bacillus subtilis, of which the morphological characters could with precision be ascertained and which at starting were tested to be barren of any power of inducing disease. I have cultivated these in pure cultivations for many generations, and under varying conditions, and then I have inoculated with them a large number of animals (mice, rabbits, and guinea-pigs); and to put it briefly, I have not found that hereby any of them acquired the least pathogenic power.

CHAPTER XVIII.

VITAL PHENOMENA OF NON-PATHOGENIC ORGANISMS.

As has been stated in a former chapter, all putrefaction of organic matter is associated with micro-organisms. It is now generally admitted, because based on a large number of exact experiments (by Schwann, Mitscherlich, Helmholtz, Pasteur, Cohn, Burdon Sanderson, Lister, W. Roberts, Tyndall, and many others), that organic matter kept safe from becoming contaminated with micro-organisms of the air, water, and filth, remains free of them, and consequently of the form of decomposition which is generally considered as putrefactive; namely, the changing of proteids into soluble peptones; then the splitting up of these into leucin and tyrosin; further the decomposition of these and other crystallisable nitrogenous bodies into comparatively low compounds. These in their turn by oxidation ultimately yield ammonia, and its salts and nitrates of inorganic elements, with the simultaneous development of certain gases, *e.g.* ammonia, sulphuretted hydrogen, and other products, belonging to the aromatic series. The view now generally entertained is that the organisms cause disintegration of nitrogenous compounds by withdrawing from them certain molecules of nitrogen, building up with these their own protoplasm. Similarly carbohydrates and inorganic salts, as phosphates, potassium, and sodium salts, are dissociated by them, inasmuch as they require a certain amount of carbon, phosphorus, potassium, and sodium, for building up their own bodies. In this process of decomposition certain alkaloids are produced, the composition of which is not accurately known, and which are called by the collective name of ptomaines (Selmi and others). These alkaloids are known to have a poisonous (toxic) effect when introduced in sufficient quantities into the system of a living animal. Very possibly the poisonous property of some articles of food, that have

undergone putrefaction or some unknown kind of fermentation, is caused by some ferment, the product of micro-organisms; (sausage poisoning, poisoning by bad fish and other articles.)

Gaspard, Panum, Bergmann, Billroth, Burdon Sanderson, Gutmann and Semmer, and many others have shown, that by putrefaction of animal substances, a substance can be obtained—the septic poison or sepsin—which can be isolated by various chemical processes destructive of every living micro-organism, and which on injection into the vascular system of animals, especially dogs, in sufficient quantities, produces a marked febrile rise of temperature, and is capable of causing death with the symptoms of acute poisoning, *e.g.* shivering, vomiting and purging, spasms, torpor, collapse and death. On post-mortem there is found severe congestion and hæmorrhage of the intestine, particularly the duodenum and rectum; hæmorrhage in the pleura, lungs, pericardium, and endocardium; congestion and hæmorrhage in the peritoneum. This *putrid infection*, or putrid intoxication, leads to death in twelve to twenty-four hours, or even less. On injecting smaller quantities only a febrile disturbance is noticed, severe symptoms and death only following after injection of considerable quantities, such as several centimetres of putrid fluid. There is *a priori* no reason why something like putrid intoxication should not occur as a pyæmic affection in the human subject; if, for instance, at an extensive wound, *e.g.* after amputation of a limb, a large surface of suppurating tissue is established, on which, as is well known, numbers of putrefactive organisms are capable of growing, it is possible and quite probable, that here these organisms produce the septic poison, which when absorbed into the system in sufficient quantities produces septic intoxication. From this affection septicæmia proper, due to absorption of a specific organism by a small open wound or a vein, which increases within the body, and therefore is a living, growing, and self-multiplying thing producing foetal septicæmia, must be carefully distinguished. As Lister has shown, under careful antiseptic dressing of wounds, putrid intoxication as well as septicæmic infection is rare or does not occur at all.

These putrefactive processes must be distinguished from certain fermentative processes, in the course of which by introducing a definite micro-organism—zymogenic organism—into a definite substance, definite chemical products are produced. Thus the *torula cervisiæ* or *saccharomyces* introduced into a solution containing sugar, produces alcoholic fermenta-

tion, *i.e.* oxidation and splitting up of sugar into alcohol and carbonic acid.

The bacterium *lactis* introduced into substances containing lactic sugar, moist, or grape sugar, produces by oxidation a conversion of the sugar into lactic acid and carbonic acid (?). A micrococcus (see a former chapter) produces, according to Pasteur, the conversion of dextrose into a sort of gum, called by Béchamp *viscose*, and recognised by Pasteur as the cause of the viscous change of wine and beer. The urea in the urine is converted by the micrococcus *ureæ* (Pasteur) into carbonate of ammonium. Solutions containing starch, dextrin, or sugar, infected with the bacillus *amylobacter* yield, as mentioned in a former chapter, butyric acid. The same bacillus converts glycerine (Fitz) into butyric acid, ethyl-alcohol, &c. Alcohol is oxidised by the presence of a definite micrococcus (Pasteur) into acetic acid.

A species of minute bacillus *subtilis* produces from fats butyric acid (Pasteur, Cohn), and many kinds of micro-organisms form pigmentary bodies, *e.g.* those producing the blue colour of milk.

What the chemical influence of pathogenic organisms on animal tissues may be is not yet known; and even when they grow outside the body, *i.e.* in artificial cultures, it is not yet known what their chemical effect on the nourishing material is, except that, as is the case with all other organisms, putrefactive and pathogenic, they continue to grow and multiply as long as there are present the necessary substances, *i.e.* until the medium is "exhausted."

From the enormous number of micro-organisms present in the outer world, it is clear that the *rôle* they play in the disintegration of higher organic bodies into lower compounds, as well as in the building up of new compounds, is a very important one; to mention but one series, to wit, the enormous importance they have for the vegetable kingdom in reducing nitrogenous compounds to soluble nitrates of inorganic salts, so essential to the existence and growth of our common field crops (Lawes).

One of the most interesting facts observed in the growth of septic micro-organisms is this, that the products of the decomposition started and maintained by them have a most detrimental influence on themselves, inhibiting their power of multiplication, in fact, after a certain amount of these products has accumulated, the organisms become arrested in their growth, and finally may be altogether killed. Thus the substances belonging to the aromatic series, *e.g.* indol, skatol,

phenol, and others, which are produced in the course of putrefaction of proteids, have a most detrimental influence on the life of many micro-organisms, as has been shown by Wernich and others (see Antiseptics).

It is not well known whether in all or in some of these instances the organisms produce the chemical effects by creating a special zymogen or ferment, and by this create the chemical disturbance, or whether they merely dissociate the compounds by abstracting for their own use certain molecules ; but this much is known, that in consequence of this chemical disturbance definite chemical substances are produced. It is quite possible that the pathogenic, like the zymogenic, organisms have this special character, that if the soil (animal body) contains a certain chemical substance, they are capable of growing and thus capable of producing a definite zymogen or ferment.

In many cases of putrefactive and zymogenic organisms a definite soil may be capable of furnishing suitable material for various organisms at the same time ; as a matter of fact this is what one constantly meets with in ordinary putrefaction of vegetable and animal matter, which teem with various species of micro-organisms. But as a rule it will be observed that one species is more apt to find a suitable soil in this substance than others ; and then it will be found that this one organism, above all others, grows more numerously ; and when it has done growing, that is, when it has exhausted its own peculiar pabulum, another organism, not dependent on this, but on some other substance still present, makes a good start and multiplies accordingly. Thus one finds constantly that, a fluid, supposing it contains a variety of proteids, carbohydrates, and salts, having become infected with micrococcus, and various species of bacilli, in the first days or weeks chiefly micrococci are present ; afterwards when the micrococci have done multiplying and sink to the bottom of the fluid, this latter gradually becomes filled with a variety of bacilli. Or if micrococcus and bacterium termo be sown at the same time in a suitable fluid, we find that at first only the bacterium termo increases rapidly ; afterwards, when this has ceased multiplying, the micrococcus takes the field. In still other cases, as in putrid blood, exudation fluid, particularly in putrefying solid parenchymatous or other substances, various kinds of micro-organisms grow on simultaneously in different parts of the material.

The same holds good for the zymogenic organisms. A *solution containing sugar is a very fit soil for saccharomyces ; when this has exhausted the material, i.e. when all the sugar*

has been split up into alcohol and carbonic acid, the latter escaping as gas, then the material is ready for the organisms capable of converting the alcohol into acetic acid. The two kinds of organisms may be, however, growing at the same time, the *saccharomyces* leading.

Septic or putrefactive organisms then, like zymogenic and pathogenic organisms, are *ceteris paribus*, dependent for their growth on the presence of the suitable nourishing material. And, as we have seen, they differ materially from one another in this respect; while putrefactive organisms find in almost all animal and vegetable fluids the necessary substances for nutrition, the zymogenic and pathogenic organisms are much more limited in these respects. Where there is no alcohol present the organisms producing the acetic acid fermentation cannot grow; where there is no sugar or a similar substance present the *saccharomyces* cannot grow, and so also a particular pathogenic organism—the *bacillus anthracis*—cannot grow in the living tissues of the living pig, dog, or cat, but grows well in those of rodents, ruminants, and man; the *bacillus* of swine-plague grows well in the pig, rabbit, and mouse, but not in the guinea-pig or man.

Septic organisms differ also from pathogenic organisms in this, that the former are capable of growing in fluids containing only simple nitrogenous compounds, *e.g.* tartrate of ammonia, whereas pathogenic organisms require more complex combinations, proteids, or allied nitrogenous substances. Thus, for instance, in Cohn's and Pasteur's fluids septic micrococcus, bacterium, and *bacillus* grow well and copiously, but pathogenic organisms absolutely refuse to grow in them; even *bacillus anthracis*, which appears the least selective, cannot make a start in it. Some organisms, *e.g.* tubercle-bacilli, require the most complex nitrogenous compounds; they refuse to grow, for instance, in broth in which anthrax-bacilli, bacilli of swine-plague, micrococcus diphtheriticus and erysipelatosus can grow well.

All septic and zymogenic organisms properly so-called, and described in former chapters, differ in this essential respect from pathogenic organisms, that the former two absolutely refuse to grow in the living tissues of a living animal.

As was stated in former chapters, it is not at all uncommon to find masses of micrococci in tissues which during the life of the subject have become dead or necrotic, or so severely changed by inflammation or otherwise that they may be considered as practically dead. In the diseased, necrotic, intestine, the liver, the spleen, in abscesses, in the subcutaneous, sub-

mucous, or parenchymatous tissues, masses of micrococci have been noticed which in no way bear any intimate relation to disease, merely finding in the dead or severely diseased tissues a suitable nidus for their growth and multiplication. But they may be present even in organs which show no severe disorganisation; thus, for instance, in fatal cases of small-pox, typhoid fever, pyæmia, even infantile diarrhoea, masses of micrococci may be found in and around the blood-vessels in the liver and spleen. In all these cases the micrococci are capable of growing, because owing to the severe general disorder these tissues have before the actual death of the patient lost their vitality, and, consequently, are unable to resist the immigration and settlement of the micrococci. Of the same character are the masses of bacilli one meets with sometimes in the intestinal wall, liver, and mesenteric glands after death from severe disorder of the bowels, *e.g.* typhoid fever and dysentery. I cannot for a moment accept the view of Klebs and Koch, that the presence of the bacilli mentioned in a former chapter necessarily stand in any causal relation to typhoid fever, seeing that they are not constant, and particularly that they are found in organs directly connected with the intestines, which we know are in this disease in an intense state of disorganisation.

The question arises: Where do the micrococci and bacilli come from which are thus capable of settling in a disorganised tissue even during the life of the subject? There can be no doubt that as regards the intestinal wall, the mesenteric glands, the liver, and the spleen, the organisms could readily, in cases of severe disorganisation of the intestine, immigrate from the cavity of the bowel, where they are normally present, into the wall of the intestine, and moreover be absorbed together with the products of disorganised tissue into the mesenteric lymphatic glands, the liver, and the spleen. Further, it is not difficult to explain that if a focus of inflammation or necrosis be set up at various internal places in consequence of emboli carried from an inflammatory focus to which micrococci or bacilli from the outer world have access, *e.g.* the skin, alimentary canal, respiratory organs, these internal places or metastases would harbour the same organisms, and as soon as disintegration—abscess, caseation, or necrosis—takes place in these metastases, also the imported organisms would multiply to a great extent, the tissue being shut out from the circulation and practically dead.

All this I say is not difficult of explanation if we bear in mind that the products of an inflammatory focus to which

organisms have ready access from the outer world become themselves a ready nidus for these organisms; and when some of these products are absorbed and taken into the general circulation to act as emboli and thus to set up inflammation in distant regions, the organisms, embedded in and shielded by those products from any destructive action the living blood in the circulation may be capable of exerting, are thus transported to the new foci of inflammation and disintegration, resulting from the emboli. All this is self-evident, and does not require more proof than what is already known, and it follows from such considerations that the presence of micrococci and bacilli in the tissues of internal organs in severe cases of disease, when some of the organs become disorganised before the actual death of the body, or in secondary foci of severe inflammation and necrosis, may have no connection whatever with the original cause of the disease or necrosis, may be, and probably are, simply due to a transportation and immigration of non-pathogenic putrefactive organisms.

A most striking case of this kind I met with in mice dead of swine-plague; the bowels were severely inflamed, and in the liver there were present necrotic patches, an almost constant symptom of the disorder; in such necrotic patches the capillary blood-vessels are sometimes, not always, found distended and plugged with the zooglœa of putrefactive micrococci, which have nothing to do, specifically, with the real disease (see a former chapter).

The cavity of the alimentary canal, small and large intestine, especially the latter, contains under normal conditions innumerable masses of putrefactive micro-organisms. These being much smaller than chyle-globules, must of necessity become as easily absorbed as the latter by the lacteals, and by these are carried into the general circulation; but being putrefactive they are unable to exist in the normal blood and normal tissues, and therefore in healthy condition perish. But if there be in any part a focus of disorganisation they can settle there and propagate, provided they get there through the blood in a living condition. Many experiments prove that they cannot pass unscathed through the normal healthy blood, and therefore it is not probable that they would reach such a focus in a living state; but let them be well inclosed in a solid particle, say of disorganised tissue, and then carried through the vascular system, and we can quite understand that in this state, *i.e.* in and with that particle, they may reach the distant focus in a living state, and if in this focus the conditions are favourable for their growth, *e.g.* if there is

inflammation and necrosis, we may expect them to multiply accordingly.

It is now admitted by most competent observers that in the healthy and normal state the blood and tissues contain no micro-organisms whatever, and that the assertions to the contrary are due to errors in the experiment, *i.e.* to accidental contamination. I will on this point merely refer, amongst many others, to the observations of Watson Cheyne¹ and F. W. Zahn.² Consequently it cannot be maintained that if in any focus of disintegration micro-organisms make their appearance they are derived from those normally present; we must, on the contrary, believe that putrefactive organisms can be imported from parts connected with the outer world into distant localities in which disorganisation of tissues has taken place.

It is clear from the foregoing that after death micro-organisms will readily immigrate into the various tissues, and in this respect those organs situated near places where under all conditions micro-organisms exist will be the first to be invaded by them; *e.g.* the lungs, from micro-organisms present in the bronchi and air-cells derived from the outer air, the walls of the alimentary canal, the mesenteric glands, the peritoneal cavity, the liver, and the spleen. The bacilli possessed of locomotion are particularly to be mentioned in this respect, but other non-motile bacilli and micrococci also find their way into these organs; thus Koch³ saw only a few hours after death bacilli (non-motile) present in the blood of the arteries of a healthy person who had died by strangulation.

¹ *Pathological Transactions*, vol. xxx.

² *Virchow's Archiv*, vol. xcv.

³ *Pathogene Micro-organismen*.

CHAPTER XIX.

VITAL PHENOMENA OF PATHOGENIC ORGANISMS.

As has been stated in the preceding chapter, the specific micro-organisms have the great differential character that they are capable of existing and propagating themselves in healthy living tissues. In those species in which the complete series of proofs has been furnished to establish the fact that the micro-organisms are intimately associated with the cause of the malady (*e.g.* malignant anthrax, tuberculosis, swine-plague erysipelas), in which it has been shown beyond doubt: (*a*) that an animal suffering from the malady contains in definite distribution the particular micro-organism, (*b*) that the micro-organisms, cleared by successive artificial cultures from any adhering hypothetical chemical virus, when introduced into a suitable animal produce the malady, (*c*) that every such affected animal contains the micro-organism, in the same distribution and relation to the diseased organs as the original animal dead after disease—in those instances, I say, the only way of understanding the effect of the micro-organisms is to assume, what is actually the case, that the micro-organisms introduced into the living tissue go on multiplying, and directly or indirectly, *i.e.* themselves or by their products, as will be stated below, produce certain definite disorders in the different parts. In the most favourable cases (anthrax, tuberculosis), a single organism introduced into a suitable locality in the animal body will be capable of starting readily a new brood. But in other cases it is necessary that an appreciable number of the organisms be introduced in order to start a brood. This was the case in some of the septicæmic processes in rodents studied by Koch.¹ The period between the time of introduction of the organism into the body (blood, sk¹

¹ *Infektionskrankheiten*, *loc. cit.*

or mucous membranes, subcutaneous tissue, lungs, alimentary canal) and the production of the new brood large enough to produce a definite effect locally or generally, corresponds to the incubation-period of the disease, and, as is well known, there is in this respect a great difference in the different diseases. Thus in anthrax the introduction of the bacilli into the subcutaneous tissue of a suitable animal is followed after from sixteen to twenty-four hours or more by a local effect (œdematous swelling), and a few hours after by general constitutional illness, when bacilli can as a rule be found in the blood. On the other hand, in tuberculosis after the introduction of the *bacilli tuberculosis* into the subcutaneous tissue, the nearest lymph-glands show the first signs of swelling and inflammation after one week or even later, and the general disease of the internal viscera does not follow until one, two, or more weeks have elapsed. This is also borne out by observations of the behaviour of these bacilli in artificial cultures; whereas a suitable material inoculated with the bacillus anthracis and kept at the temperature of the animal body (38° to 39° C.), shows already after twenty-four hours a good crop of the bacilli; in the case of the tubercle-bacilli the first signs of a new brood are not noticed, as Koch has pointed out, and as I have had in several instances occasion to verify, before ten to fourteen days have passed.

One of the most important points, and the most difficult of comprehension, is this power of the pathogenic organisms to resist the influence of the healthy tissues of the living animals, a power which we said above is not possessed by the non-pathogenic organisms. A careful analysis shows at the outset that this power of the pathogenic organisms is not possessed by them indiscriminately, for while a particular species is in some animals capable of overcoming the influence of the living tissue, *i.e.* to multiply and to produce the particular disease, in other animals it is not capable of doing this, and hence the animal remains unaffected—it is said to be not susceptible to the disease. Thus, for instance, the bacillus anthracis when introduced into a human being, or a herbivorous animal, is capable of multiplying and of producing anthrax, whereas in carnivorous animals and even in the omnivorous pig it is not capable of doing so. Or again, the bacillus of swine-plague while capable of producing the disease in swine, rabbits, and mice, is not capable of doing

bird, the guinea-pig, or carnivorous animals.

to look for this difference in behaviour?

* of a pig when obtained as infusion or
nourishing material for the bacillus

anthracis as the tissues and juices obtained from a herbivorous animal; artificial cultures of the former and of the latter behave in exactly the same manner, both as regards copiousness of growth and virulence of the bacilli. Again, artificial cultures of the bacillus of swine-plague made in juice of the tissues of the guinea-pig or fowl are exactly the same as those made of the juice of the tissues of a rabbit or pig. The tissues, therefore, *per se* cannot be said to possess any inimical action on the organisms. The living condition *per se* can also not have this power, since we see that the power to overcome the influence of the living tissue is precisely the great distinguishing character of pathogenic organisms. There remains, therefore, only one thing, that is that there is something or other present in a particular tissue to which this latter owes its immunity, and this something must of necessity be connected with the tissue while alive, as we said before. Now, the life of the tissue in the pig cannot be different from that of the mouse, if by life is understood the function of the tissue, the connexion with the vascular and nervous system, and all the rest of it. The subcutaneous connective tissue has no different function, no different relation to the vascular and nervous system in the pig from what it has in the mouse, and nevertheless we find that it behaves so differently in the two cases towards the bacillus anthracis. This something then, which inhibits the growth and multiplication of the bacillus anthracis in the tissue of the pig but not in the mouse, must be something which, although dependent on the life of the tissue, is not identical with any of the characters constituting the life of the tissue, but must be some product of that life. To assume then, as is done by some observers, that the living state of the cells *per se* is the inhibitory power does not cover the facts, as we have just shown. The most feasible theory seems to me to be this, that this inhibitory power is due to the *presence of a chemical substance* produced by the living tissues. It does not require any great effort to conceive, and it does not seem at all improbable, that the blood and tissues of the pig contain certain chemical substances which are not present in the mouse, substances which like so many others chemistry is not yet capable of demonstrating. But that there exist vast and gross differences in the chemical constitution of the blood and tissues of different species of animals there can be no reasonable doubt; it is a fact with which physiological chemistry is quite familiar.

We arrive then, after all this, at the conclusion that to the presence in the blood and tissues of particular c

substances, present only during life, and a result of the life of the tissue, the organisms in a particular case cannot thrive and produce the disease. And further, that for each particular species of organism there is a particular chemical substance required to exert this inhibitory power, for, as we have seen, while the anthrax-bacillus is not capable of thriving in the pig, it does well in the guinea-pig, while the bacillus of swine-plague thrives well in the pig, it does not in the guinea-pig. The incapability of non-pathogenic organisms to thrive in healthy living tissues would on this theory be explained by the assumption that these chemical substances present in every healthy living tissue are inimical to all putrefactive organisms.

What we have said hitherto refers only to the healthy living tissues. It is quite possible to imagine that, owing to the presence of a particular chemical substance in the healthy living tissue, a pathogenic organism is not able to thrive in a particular animal; but under certain abnormal conditions, when for instance owing to a diseased or altered state of the tissue that substance is absent, the organism might be enabled to exist and multiply and to produce the disease. Supposing it to be true that a person so long as he is healthy and well is able to successfully withstand the growth of a particular pathogenic organism, he is then unsusceptible to the disease; but we can understand that if the alimentary canal or the lungs be diseased, then the organism passing into the bowels by ingestion or into the lungs by inspiration would find there a tissue in which the necessary inhibitory substance, present in the healthy state, might be absent, and the organism would be capable of thriving and of producing the disease.

The next point to consider is the relation of the specific micro-organism to the essence of the disease, or in other words the question whether the organism itself is the virus or whether this latter is the product of the former, something in the same way as the septic ferment is the product of the putrefactive organisms?

That the virus in the infectious diseases is intimately connected with the organisms is proved by the fact, that the virus introduced into the body multiplies *ad infinitum*; for instance, in anthrax or tuberculosis after the introduction of an infinitesimal dose, we find the disease (malignant anthrax or general tuberculosis respectively) sets in in its virulent form; in the *first case every* drop of blood teems with the bacillus anthracis; *in the second* (tuberculosis) every tuberculous caseous particle

in the lymph-glands, lungs, spleen, and liver contains the bacilli; in both instances crops of the bacilli are produced in the afflicted body, and every particle of the tissue containing the bacilli is capable of starting the disease when introduced into a fresh subject. Moreover the artificial cultures of the pure bacilli are possessed of the same pathogenic power. The same holds good for leprosy, for erisypelae, for swine-plague, &c. So that the proposition that the organisms are intimately connected with the virus must be considered as well established.

But even after this it remains an open question whether the organism is identical with the virus, or whether the organism is concerned in elaborating the virus—a sort of ferment; and further, whether the virus being the latter's product, is obtainable apart from the organism.

Let us start with the proposition that the virus is a product of the organism, a sort of non-organised ferment, but not the organism itself, although this latter is essential for the creation of the latter.

Inoculating a few bacilli anthracis into the subcutaneous tissue of a suitable animal, e.g. a guinea-pig, we find after twelve to twenty-four hours the first indications of illness, consisting in a local swelling and a general rise of the body-temperature. At this time there are present in the local swelling bacilli, but only in small numbers; in the blood the bacilli are very scarce indeed, so scarce that it is difficult to meet with one bacillus in an appreciable quantity of blood. By this time then the bacilli could not have produced the change by their "numbers" alone. Immediately before death, sometimes some hours, we find in most instances the blood teeming with the bacilli, but this is by no means in all cases; I have seen a considerable number of deaths from typical anthrax in the mouse, guinea-pig, rabbit, and sheep, occurring from forty-eight to sixty hours after inoculation, in which the number of bacilli of the blood and tissues was extremely small; they were present, but only here and there was there one to be found. That the bacilli are present some hours before death in the shape of spores, as has been maintained by Archangelski, I have disproved in a former chapter. In those cases in which the bacilli are scarce even *in articulo mortis* and immediately after death, the scarcity is not due to the bacilli having already degenerated, since the degenerating bacilli are not noticeable in any way in these instances. It remains then to assume that death occurs; these cases not owing to the presence of the bacilli in numbers

that is to say, that it is neither owing to their appropriating from the blood-corpuscles the available oxygen necessary for their multiplication (Bollinger) and thus producing death by asphyxia, nor to their mechanical effect in plugging up the capillaries of vital organs, a theory upheld by some observers from the fact that in most cases the capillaries appear filled with the bacilli, and in some cases in extensive regions, lung, kidney, and spleen, the capillaries are almost occluded by the bacilli. We must assume, then, that although as a rule immediately before death all the conditions are present to enable the bacilli to multiply readily and to produce a large crop, this is not necessarily connected with the cause of death, being in fact in consequence of the animal being *in articulo mortis*; but that the immediate cause of death is the chemical alteration produced by the bacilli in the blood and tissues. For producing this effect it is not necessary to have more than a certain number of the bacilli. As soon as this number is reached death follows. The same may be said of other pathogenic organisms. Thus for instance in the case of tubercle-bacilli, after the introduction of these into the subcutaneous tissue of a guinea-pig, multiplication takes place, and after they reach a certain number, the nearest lymph-glands become swollen and inflamed and then caseous; but this stands in no relation to the number of the bacilli, for in some instances the microscopic examination reveals only very few bacilli, they are scattered in very small numbers over very wide areas. And the same is observed in the tuberculous deposits of the internal organs. In some of them the bacilli are exceedingly scarce, while in others neither more nor less advanced they are numerous. Here also we must assume that as soon as a certain, perhaps even small, number of the bacilli has been produced the chemical effect produced is sufficient to be the cause of a certain pathological change. In glanders, the nodules in the skin and lung reveal sometimes, even under the most careful examination after approved methods, the presence of but very few bacilli. In swine-plague, in the lungs, which in severe cases are enormously affected, sometimes only very few of the pathogenic organisms can be discovered. It follows then that the pathological condition brought about by the organisms is not due to the direct action of their numbers; but is an indirect sequence, brought about by definite chemical alterations in the blood or tissues as the case may be.

In this we may assume two theories as possible: (*a*) It is possible that these chemical effects are produced by the presence and growth of the organisms, as truly as in the

alcoholic fermentation of sugar the alcohol produced is a result of the presence of the saccharomyces ; this is only in so far a product of the organism as this, in its multiplication, assimilates some molecules of carbon and hydrogen, which it abstracts from the sugar, and in consequence of which the sugar yields alcohol ; but it is not, as it were, a secretion of the organism, a special ferment. (b) But it is likewise possible that the organism elaborates a special ferment, which, after a certain amount has been produced, sets up the particular pathological changes. From these considerations it follows that the virus cannot be considered independent of the organism ; we cannot assume that the two can be separated from one another ; for, as we have just now shown, the most feasible assumption, and the one borne out by observation, is that owing to the multiplication of the organisms, certain chemical changes are produced in the blood and tissues, or that a special ferment is created, which sets up the anatomical changes characteristic of the particular disease.

CHAPTER XX.

VACCINATION AND IMMUNITY.

WE have in the foregoing chapter tried to show that owing to the presence in the normal blood and tissues of a living animal of some chemical substances varying in the different species, and inimical to particular pathogenic organisms, the latter, when introduced into the tissues of the particular species, cannot thrive, and that it is for this reason that the animal is not susceptible to the corresponding disease. Now, how do we explain the fact that a human being or an animal having been once the subject of a particular infectious disease, becomes thereby in some cases unsusceptible to a second attack? The oldest and perhaps the most favoured theory to explain this immunity is that which assumes that during the first attack the organisms growing in the body consume, or are instrumental in eliminating or destroying, some chemical compound necessary for the existence and multiplication of the organism. As soon as this substance has become consumed or destroyed the organisms cannot further multiply, and therefore the disease ceases; and further, that owing to the subsequent absence of this same chemical compound, a new infection by the same organisms is not possible, *i.e.* the individual is protected. Thus this theory puts the case on a level with, say, the relation of the saccharomyces to the alcoholic fermentation; as long as a solution contains sugar, the saccharomyces is capable of multiplying, but as soon as all the sugar has disappeared as such, *i.e.* has become split up into alcohol and carbonic acid, the fermentation ceases, the solution being now exhausted as regards the saccharomyces; a new charge of saccharomyces put into the solution is not *capable* of multiplication. This theory, then, to explain the *immunity*, is generally spoken of as the *Exhaustion Theory*.

On careful analysis, it will be found that it is not capable of explaining all the facts of the case. As we mentioned in a former chapter, cattle inoculated with blood of a guinea-pig dead of anthrax become affected with anthrax, which, although not fatal, is nevertheless sometimes very severe. The animal recovers, and is now, for a time at least, protected against a second attack. But there is absolutely no ground for the assumption that if an infusion of the tissues of this animal were made, the bacillus anthracis sown in it would not thrive luxuriantly, seeing that bacillus anthracis grows on almost anything that contains a trace of proteids. Similarly when of the tissues of a guinea-pig, or mouse or rabbit, dead of anthrax, an infusion is made, and this is used as nourishing material for bacillus anthracis in artificial cultures, it is found that these latter thrive splendidly. The same fact I have observed in the case of swine-plague. There is then no reason whatever for assuming that, after one attack of illness, the blood and tissues become an unfavourable soil for a second invasion of the same organism, and that this should be due to the exhaustion of some necessary chemical compound.

There is another theory, commonly spoken of as the *Antidote Theory* (Klebs). According to this, the organisms growing and multiplying in the body during the first attack produce, directly or indirectly, some substance which acts as a sort of poison against a second immigration of the same organism. I am inclined to think that this theory is in harmony with the facts. There is nothing known, from the observations before us, which would negative the possibility of the correctness of this theory; nay, I would almost say all our knowledge of the life of the micro-organisms points to the conclusion that the different species are associated with different kinds of chemical processes, and that as a result of the activity we find different chemical substances produced.

The different fermentations connected with the different species of fungi afford striking illustrations of this view. According to this theory, we can well understand that—just as in the case of an animal, say a pig, unsusceptible to anthrax—the unsusceptibility being due to the presence in the blood and tissues of a particular chemical substance inimical to the growth of the bacillus anthracis—so also in the case of a sheep or ox that has once passed through anthrax—there is now present in the blood and tissues a chemical substance inimical to the growth and multiplication of the bacillus anthracis whereby these animals become possessed of immunity against a second attack of anthrax.

Whether this chemical substance has been elaborated directly by the bacilli, or whether it is a result of the chemical processes induced in the body by the bacilli during the first illness, matters not at all ; it is only necessary to assume that the blood and tissues of the living animal contain this chemical substance.

Some observers (Grawitz, &c.) are not satisfied with this theory, but assume that owing to the first attack the cells of the tissues so change their nature that they become capable of resisting the immigration of a new generation of the same organism. There is absolutely nothing that I know of in favour of such a theory ; it is impossible to imagine that the cells of the connective tissues, of the blood and of other organs, owing to a past attack of scarlatina, become possessed of new functions or of some new power, as, for instance, a greater power of oxidising or the like. Connective tissue-cells, blood-corpuscles, liver-cells, and other tissues are, so far as we know, possessed of precisely the same characters and functions after an attack of scarlatina as before.

On the whole then, we may, it seems, take it as probable, that owing to the presence in the normal blood and tissues in a living animal of a chemical substance inimical to the growth of a particular micro-organism, this animal is unsusceptible to the disease dependent on the growth and multiplication of this micro-organism ; and further, that in those infectious maladies in which one attack gives immunity against a second attack of the same kind, one attack produces a chemical substance in the blood and tissues which acts inimically to a new immigration of the same organism ; hence the animal becomes unsusceptible to a new attack, or is "protected." This is not the case with all infectious maladies, for, as is well known, in a good many instances a single attack does not protect against a second ; and, as is also well known, a first attack may protect but only for a limited period, or for a period greatly differing in different individuals. All this would be explained by our theory in the same way as it is explained by the other theories ; viz., when one attack does not protect, no inhibitory chemical substance has been produced ; while in those diseases in which one attack does protect only for a limited period, the necessary inhibitory substance has only lasted for a limited period, and so on.

CHAPTER XXI.

ANTISEPTICS.

IN former chapters we have on several occasions mentioned that a variety of substances and conditions are capable of exerting a detrimental influence on the life and growth of micro-organisms. Amongst these are—The presence of certain substances in the nutrient soil, the temperature, and some chemical products, such as those belonging to the aromatic series, phenol, indol, skatol, &c. The presence of certain substances in the nourishing material is, as we have seen, an essential condition, *cæteris paribus*, for the growth and multiplication of micro-organisms. Thus pathogenic organisms cannot thrive in an acid medium, they cannot thrive if proteids or allied compounds and certain inorganic salts are absent; putrefactive and zymogenic organisms, on the other hand, or, at any rate, some of them, are capable of thriving well in acid media (*e.g.* the bacillus subtilis in acid hay-infusion, the micrococcus ureæ in acid urine). Further, many (not all) pathogenic organisms cannot thrive unless they are exposed to a certain degree of warmth; they thrive best at blood-heat, while putrefactive and many zymogenic organisms thrive well at ordinary temperatures, though of course their growth is more rapid at higher temperatures, such as 30°—38° C. Heat above 50° or 60° C. arrests the growth of and even kills many organisms, except the spores of bacilli, which, as we find on a former page, survive even when exposed to the temperature of boiling water for several minutes. The presence of carbolic acid, phenol, thymol, salicylic acid, perchloride of mercury,

&c., prevent even when in great dilution the growth of micro-organisms.

In any inquiry into the influence of one substance or another on micro-organisms it is necessary to bear in mind that the influence of certain conditions on the micro-organisms may be a twofold one : (1) the condition may be unfavourable to the growth of the organism in question, and (2) the condition may be fatal to the life and existence of it. The second condition involves, *a fortiori*, the first ; but the reverse is not the case. Owing to the failure to distinguish between these two propositions a great deal of confusion has arisen on the subject. One hears constantly this or that substance is an "antiseptic," meaning by this a substance inimical to the growth of micro-organisms, or a substance is a "germicide," implying by this that this substance kills the organisms ; but when one comes to analyse the observations that are said to establish this reputation for a particular substance, one finds that the substances in question are really only detrimental to the growth of the organisms.

By sowing any micro-organism into a nourishing medium, to which has been added a certain substance (*e.g.* carbolic acid to the amount of 1 per cent.), and exposing this medium to the conditions of temperature, moisture, &c., otherwise favourable to the growth of the organism, if we find that after the lapse of a due period the growth is retarded or altogether inhibited, the conclusion is drawn that this substance (*viz.*, the carbolic acid of 1 per cent.) is an antiseptic. There is nothing more fallacious than this method of reasoning ; a great many micro-organisms can be exposed to a 1 per cent. solution of carbolic acid for hours without in the least being affected, for on being then transferred to a suitable nourishing medium they grow and thrive well. Similarly by placing the spores of bacillus anthracis in a proteid medium containing perchloride of mercury of the strength of 1 in 300,000, it is found (as Koch has shown) that the spores are absolutely incapable of germinating. But if from this the conclusion is drawn, that perchloride of mercury of the strength of 1 in 300,000 is a germicide, I should most strongly dissent, for perchloride of mercury even of the strength of 1 per cent. is not a germicide any more than vinegar ; for on placing the spores of bacillus anthracis in a proteid medium, to which so much vinegar or any other acid has been added as makes it decidedly acid, it will be found that the spores do not *germinate*.

In order to pronounce a certain substance an antiseptic in the strict sense of the word, it is necessary to place the organisms in this substance for a definite time, then to remove them thence, and to place them in a suitable nourishing medium; if they then refuse to grow the conclusion is justified that the exposure has injured or destroyed the life of the organisms. In the case of pathogenic organisms a substance to be pronounced a germicide must be shown to have this power, that when the organism is exposed to the substance and then introduced into a suitable artificial medium it refuses to grow; and it must also be shown that when introduced into a suitable animal it is incapable of producing the disease which the same organism, unexposed to the substance in question, does produce.

I have made a good many observations on the influence of antiseptics on micro-organisms, both putrefactive and pathogenic, and I have found that many assertions hitherto made on this subject, treated in the above light, are absolutely untrustworthy and erroneous.

Various species of putrefactive micrococci, bacterium termo, bacillus subtilis, various pathogenic micro-organisms, as bacillus anthracis, bacillus of swine fever, absolutely refuse to grow in media to which is added phenyl-propionic acid, or phenyl-acetic acid, to an amount so small as 1 in 1,600; but if the same organisms are exposed to these substances in much stronger solutions, 1 in 800, 1 in 400, or even 1 in 200, and then transferred to a suitable nourishing material, it is found that they have completely retained their vitality, they multiply as if nothing had been done to them. I have exposed the spores of bacillus anthracis to the above acids of the strength of 1 in 200 for forty-eight hours and longer, and then inoculated guinea-pigs with them, and I found that the animals died of typical anthrax in exactly the same way as if they had been inoculated with pure spores of the bacillus anthracis.

Koch has published a large series of systematic and most valuable observations¹ made in testing the influence on spores of bacillus anthracis of a large number of antiseptics (thymol, arsenate of potassium, turpentine, clove-oil, iodine, hydrochloric acid, permanganate of potassium, eucalyptol, camphor, quinine, salicylic acid, benzoic acid, and many others), and amongst them he found perchloride of mercury to be the most powerful, since even a solution of 1 in 600,000 is capable of impeding, one of 1 in 300,000 of completely checking, the germinating

¹ *Mittheil. aus d. k. Gesundheitsamte, Berlin, 1881.*

power of the spores. To regard these substances, from these observations, in any way as antiseptics for the spores of bacillus anthracis would be no more justifiable than to consider weak vinegar as such. Perchloride of mercury in a solution of 1 in 300,000 is no more capable of interfering with the life and functions of the spores of bacillus anthracis than water or salt solution, for the spores may be steeped in this solution for any length of time, and yet on being transferred to a suitable medium they grow and multiply splendidly, and when inoculated into rodents they produce fatal anthrax with absolute certainty. With my friend Dr. Blyth, Medical Officer of Health for the Marylebone district in London, I have tried the action of a number of substances in common use as antiseptics (*e.g.* Calvert's fluid, pure terebene, phenol 10 per cent., perchloride of mercury 1 per cent.) on the spores of bacillus anthracis, exposing these in comparatively large quantities of the above fluids (the two being well mixed) for twenty-four hours, and then inoculating guinea-pigs with them (spores and antiseptic). The animals died with symptoms of typical anthrax, the blood teeming with the bacillus anthracis.

These substances then are no more antiseptics, and still less germicides for the spores of bacillus anthracis, than water is.

In all these inquiries, particularly in those upon pathogenic organisms capable of forming spores, the influence of the substances must be judged not merely by their action on the organisms, but also on the spores; for, in this very case of the bacillus anthracis, the bacilli taken from the blood of an animal dead of anthrax are killed after an exposure of say ten minutes to a solution of phenyl-propionic acid of the strength of 1 in 400, or even 1 in 800, whereas the spores of the bacilli (produced in artificial cultures) withstand completely exposure to this acid of any strength and for any length of time.

It is not my object to pass here in review all that has been done in this interesting field of research, important because of its obviously great practical importance. The work hitherto done has been enormous, but, I fear, of less utility than at first sight appears, for in most of it the point most prominent in the mind of the worker was to ascertain whether the particular antiseptic, mixed with the nourishing medium in a solution of definite strength, has or has not the power of inhibiting the growth of the micro-organisms. This point no doubt is of some interest, and perhaps of great interest, but whether a particular substance is an antiseptic in the proper *sense* of the word, *i.e.* whether on exposing the organisms to

this substance in a solution of definite strength and for a definite period, the organisms become afterwards incapacitated from growing or multiplying ; or still more, whether or not the substance is a germicide, *i.e.* is capable of altogether annihilating the life of the organisms ; these are questions which require special attention, and represent a wide and rich field of inquiry ; but, as far as I can see, it has received only in very few instances due attention.

1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025

1941-1945

1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025


INDEX.

- ABRIN**, 165
Abrus precatorius, 160
Abcess in rabbit, 57
Achorion Schönleini, 141
Actinomyces, 148
Acute inflammations, 43
Aerobic, 34
Agar-Agar, 16
Air examination, 32
Amylobacter, 80
Anæmia perniciosa, 55
Anaerobic, 34
Anilin dyes, 6
Anilin oil, 7
Antheridia, 146
Anthrax bacillus, 105
Antiseptics, 137
Aromatic products, 171
Artificial cultures, 25
Artificial tuberculosis, 125
Ascococcus Billrothi, 42
Ascogonium, 144
Ascomycetes, 140
Ascospores, 140
Aspergillus, 142
Attenuation, 116
- BACILLUS**, general characters of, 66
 septic, 75
 zymogenic, 80
 chromogenic, 81
 pathogenic, 83
Bacterium, general characters of, 60
 termo, 60
 lactis, 61
 lineola, 61
Bacteria, pathogenic, 63
Bacteridia, 105
Basidia, 144
Beggiatoa, 79
Broth, preparation of, 11
Buchner's fluid, 13
- CAPILLARY pipette**, 19
Carbolic acid, 188
Carpegonium, 144
Cattle plague, 54
Charbon symptomatique, 103
Cheyne's method of staining, 7
Cholera bacillus, 126
Choleraic diarrhoea, 67
Cladotrix dichotoma, 78
Clathrocystis, 42
Clostridium butyricum, 81
Cohn's fluid, 14
Conidia, 140
Cotton-wool, 19
- DAVAINE'S septicæmia**, 64
Desmobacteria, 66
Diphtheria micrococcus, 48
Diplococcus, 38
Dispora caucasica, 81
Dumb-bell micrococcus, 37
Dysentery, 128
- EHRlich's method of staining**, 120
Endocarditis ulcerosa, 53
Erysipelas micrococcus, 48
- FAVUS fungus**, 141
Flasks, sterilisation of, 17
Fowl cholera, 64
- GELATINE**, 14
Gelose, 16
Germicide, 188
Gibbes' method of staining, 6
Glanders bacillus, 92
Glass-cell, 30
Gonorrhœa micrococcus, 53

- HAY** bacillus, 75
 Hay infusion, 76
 Herpes tonsurans, 141
 Hot-air chamber, 18
 Hydrocele fluid, 13
 Hyphæ, 140
- IMMUNITY**, 148
 Incubator, 9
 Indol, 171
 Infectious diseases, 2
 Inoculation, method of, 25
- JEQUIRITY** bacillus, 160
 Jequirity infusion, 161
 Jequirity ophthalmia, 161
 Jequirity seeds, 160
- KLEBS'S** method of cultivation, 26
 Koch's malignant œdema, 101
 Koch's method of cultivation, 28
 inoculation, 28
 staining, 7, 119
 progressive necrosis, 56
 septicæmia of mouse, 83
 of rabbit, 58
 Koch-Weigert method, 6
- LEPROSY** bacillus, 93
 Leptothrix, 66
 Leptothrix buccalis, 68
 Lister's method of cultivation, 26
- MALARIA** bacillus, 89
 Malignant œdema, 101
 Meat extract, 18
 Meat poisoning, 87
 Microbe of fowl cholera, 61
 Micrococcus, 37
 general character, 37
 chromogenic, 40
 pathogenic, 43
 septic, 40
 zymogenic, 40
 Microsporon furfur, 142
 Microzyma bombycis, 59
 Monas, 42
 Mucor, 146
 Mycelium, 140
 Mycoderma aceti, 62
 Mycoderma saccharomyces, 138
 Mycosis intestinalis, 107
 Mycothrix, 37
- NAGEL'S* method of culture, 26
Noma, 92
- NOURISHING** media, 9
 fluids, 11
 solids, 14
 preparation of, 21
 sterilisation of, 12
 Nosema bombycis, 59
- ŒDEMA**, malignant, 101
 Oidium albicans, 139
 Oidium lactis, 140
 Oogonium, 146
 Oospores, 146
 Osteomyelitis, 55
- PAPAYOTIN**, 103
 Pasteur's fluid, 14
 Pasteur's septicæmia, 102
 Pasteur's vaccin charbonneux, 116
 Pathogenic organisms, 2
 vital phenomena of, 177
 Penicillium, 146
 Peptone, 13
 Perithecium, 144
 Phenol, 171
 Pityriasis versicolor, 141
 Pleuro-pneumonia, 51
 Pneumococcus, 49
 Pollinodia, 144
 Progressive necrosis in mouse, 56
 Ptomaines, 169
 Puerperal fever, 54
 Pyæmia of mouse, 57
 rabbit, 57
- SACCHAROMYCES**, 136
 Salmon disease, 146
 Saprolegnia, 146
 Sarcina, 39, 43
 Scarlatina micrococcus, 54
 Schizomycetes, 34
 Septic ferment, 170
 Septicæmia of man, 84
 mouse, 83
 rabbit, 58
 Serum of blood, 13
 Silk-worm diseases, 59
 Spherobacterium, 37
 Spirillum, 131
 septic, 131
 pigment, 133
 pathogenic, 134
 Spirochæta, 132
 Sporangium, 140
 Spores, general characters, 73
 formation of, 69
 germination, 74
 of anthrax bacilli, 112
 Stomatitis ulcerative, 90

- Sterilisation of instruments, 19
 Streptothrix, 78
 Swine plague, 94
- TEST-TUBES, 21
 Thallus, 140
 Thrush, 139
 Torula, 136
 Torula form of bacillus, 109
 Trichophyton tonsurans, 142
 Tuberculosis bacillus, 118
 Typhoid fever bacillus, 86
- UREÆ micrococcus, 40
- VACCIN charbonneux, 116
 Vaccination, 184
 Vaccinia and Variola micrococcus, 46
- Vibrio, 129
 Virus, general character of, 180
- WATER examination, 31
- XANTHINUM bacterium, 62
- YEAST fungus, 136
- ZOOLCEA, 39
 Zooglyca ramigera, 60
 Zoospores, 146
 Zymogen, 172
 Zymogenic bacteria, 61
 bacilli, 80
 micrococci, 40

THE END.



1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236
2237
2238
2239
2240
2241
2242
2243
2244
2245
2246
2247
2248
2249
2250
2251
2252
2253
2254
2255
2256
2257
2258
2259
2260
2261
2262
2263
2264
2265
2266
2267
2268
2269
2270
2271
2272
2273
2274
2275
2276
2277
2278
2279
2280
2281
2282
2283
2284
2285
2286
2287
2288
2289
2290
2291
2292
2293
2294
2295
2296
2297
2298
2299
2300
2301
2302
2303
2304
2305
2306
2307
2308
2309
2310
2311
2312
2313
2314
2315
2316
2317
2318
2319
2320
2321
2322
2323
2324
2325
2326
2327
2328
2329
2330
2331
2332
2333
2334
2335
2336
2337
2338
2339
2340
2341
2342
2343
2344
2345
2346
2347
2348
2349
2350
2351
2352
2353
2354
2355
2356
2357
2358
2359
2360
2361
2362
2363
2364
2365
2366
2367
2368
2369
2370
2371
2372
2373
2374
2375
2376
2377
2378
2379
2380
2381
2382
2383
2384
2385
2386
2387
2388
2389
2390
2391
2392
2393
2394
2395
2396
2397
2398
2399
2400
2401
2402
2403
2404
2405
2406
2407
2408
2409
2410
2411
2412
2413
2414
2415
2416
2417
2418
2419
2420
2421
2422
2423
2424
2425
2426
2427
2428
2429
2430
2431
2432
2433
2434
2435
2436
2437
2438
2439
2440
2441
2442
2443
2444
2445
2446
2447
2448
2449
2450
2451
2452
2453
2454
2455
2456
2457
2458
2459
2460
2461
2462
2463
2464
2465
2466
2467
2468
2469
2470
2471
2472
2473
2474
2475
2476
2477
2478
2479
2480
2481
2482
2483
2484
2485
2486
2487
2488
2489
2490
2491
2492
2493
2494
2495
2496
2497
2498
2499
2500
2501
2502
2503
2504
2505
2506
2507
2508
2509
2510
2511
2512
2513
2514
2515
2516
2517
2518
2519
2520
2521
2522
2523
2524
2525
2526
2527
2528
2529
2530
2531
2532
2533
2534
2535
2536
2537
2538
2539
2540
2541
2542
2543
2544
2545
2546
2547
2548
2549
2550
2551
2552
2553
2554
2555
2556
2557
2558
2559
2560
2561
2562
2563
2564
2565
2566
2567
2568
2569
2570
2571
2572
2573
2574
2575
2576
2577
2578
2579
2580
2581
2582
2583
2584
2585
2586
2587
2588
2589
2590
2591
2592
2593
2594
2595
2596
2597
2598
2599
2600
2601
2602
2603
2604
2605
2606
2607
2608
2609
2610
2611
2612
2613
2614
2615
2616
2617
2618
2619
2620
2621
2622
2623
2624
2625
2626
2627
2628
2629
2630
2631
2632
2633
2634
2635
2636
2637
2638
2639
2640
2641
2642
2643
2644
2645
2646
2647
2648
2649
2650
2651
2652
2653
2654
2655
2656
2657
2658
2659
2660
2661
2662
2663
2664
2665
2666
2667
2668
2669
2670
2671
2672
2673
2674
2675
2676
2677
2678
2679
2680
2681
2682
2683
2684
2685
2686
2687
2688
2689
2690
2691
2692
2693
2694
2695
2696
2697
2698
2699
2700
2701
2702
2703
2704
2705
2706
2707
2708
2709
2710
2711
2712
2713
2714
2715
2716
2717
2718
2719
2720
2721
2722
2723
2724
2725
2726
2727
2728
2729
2730
2731
2732
2733
2734
2735
2736
2737
2738
2739
2740
2741
2742
2743
2744
2745
2746
2747
2748
2749
2750
2751
2752
2753
2754
2755
2756
2757
2758
2759
2760
2761
2762
2763
2764
2765
2766
2767
2768
2769
2770
2771
2772
2773
2774
2775
2776
2777
2778
2779
2780
2781
2782
2783
2784
2785
2786
2787
2788
2789
2790
2791
2792
2793
2794
2795
2796
2797
2798
2799
2800
2801
2802
2803
2804
2805
2806
2807
2808
2809
2810
2811
2812
2813
2814
2815
2816
2817
2818
2819
2820
2821
2822
2823
2824
2825
2826
2827
2828
2829
2830
2831
2832
2833
2834
2835
2836
2837
2838
2839
2840
2841
2842
2843
2844
2845
2846
2847
2848
2849
2850
2851
2852
2853
2854
2855
2856
2857
2858
2859
2860
2861
2862
2863
2864
2865
2866
2867
2868
2869
2870
2871
2872
2873
2874
2875
2876
2877
2878
2879
2880
2881
2882
2883
2884
2885
2886
2887
2888
2889
2890
2891
2892
2893
2894
2895
2896
2897
2898
2899
2900
2901
2902
2903
2904
2905
2906
2907
2908
2909
2910
2911
2912
2913
2914
2915
2916
2917
2918
2919
2920
2921
2922
2923
2924
2925
2926
2927
2928
2929
2930
2931
2932
2933
2934
2935
2936
2937
2938
2939
2940
2941
2942
2943
2944
2945
2946
2947
2948
2949
2950
2951
2952
2953
2954
2955
2956
2957
2958
2959
2960
2961
2962
2963
2964
2965
2966
2967
2968
2969
2970
2971
2972
2973
2974
2975
2976
2977
2978
2979
2980
2981
2982
2983
2984
2985
2986
2987
2988
2989
2990
2991
2992
2993
2994
2995
2996
2997
2998
2999
3000
3001
3002
3003
3004
3005
3006
3007
3008
3009
3010
3011
3012
3013
3014
3015
3016
3017
3018
3019
3020
3021
3022
3023
3024
3025
3026
3027
3028
3029
3030
3031
3032
3033
3034
3035
3036
3037
3038
3039
3040
3041
3042
3043
3044
3045
3046
3047
3048
3049
3050
3051
3052
3053
3054
3055
3056
3057
3058
3059
3060
3061
3062
3063
3064
3065
3066
3067
3068
3069
3070
3071
3072
3073
3074
3075
3076
3077
3078
3079
3080
3081
3082
3083
3084
3085
3086
3087
3088
3089
3090
3091
3092
3093
3094
3095
3096
3097
3098
3099
3100
3101
3102
3103
3104
3105
3106
3107
3108
3109
3110
3111
3112
3113
3114
3115
3116
3117
3118
3119
3120
3121
3122
3123
3124
3125
3126
3127
3128
3129
3130
3131
3132
3133
3134
3135
3136
3137
3138
3139
3140
3141
3142
3143
3144
3145
3146
3147
3148
3149
3150
3151
3152
3153
3154
3155
3156
3157
3158
3159
3160
3161
3162
3163
3164
3165
3166
3167
3168
3169
3170
3171
3172
3173
3174
3175
3176
3177
3178
3179
3180
3181
3182
3183
3184
3185
3186
3187
3188
3189
3190
3191
3192
3193
3194
3195
3196
3197
3198
3199
3200
3201
3202
3203
3204
3205
3206
3207
3208
3209
3210
3211
3212
3213
3214
3215
3216
3217
3218
3219
3220
3221
3222
3223
3224
3225
3226
3227
3228
3229
3230
3231
3232
3233
3234
3235
3236
3237
3238
3239
3240
3241
3242
3243
3244
3245
3246
3247
3248
3249
3250
3251
3252
3253
3254
3255
3256
3257
3258
3259
3260
3261
3262
3263
3264
3265
3266
3267
3268
3269
3270
3271
3272
3273
3274
3275
3276
3277
3278
3279
3280
3281
3282
3283
3284
3285
3286
3287
3288
3289
3290
3291
3292
3293
3294
3295
3296
3297
3298
3299
3300
3301
3302
3303
3304
3305
3306
3307
3308
3309
3310
3311
3312
3313
3314
3315
3316
3317
3318
3319
3320
3321
3322
3323
3324
3325
3326
3327
3328
3329
3330
3331
3332
3333
3334
3335
3336
3337
3338
3339
3340
3341
3342
3343
3344
3345
3346
3347
3348
3349
3350
3351
3352
3353
3354
3355
3356
3357
3358
3359
3360
3361
3362
3363
3364
3365
3366
3367
3368
3369
3370
3371
3372
3373
3374
3375
3376
3377
3378
3379
3380
3381
3382
3383
3384
3385
3386
3387
3388
3389
3390
3391
3392
3393
3394
3395
3396
3397
3398
3399
3400
3401
3402
3403
3404
3405
3406
3407
3408
3409
3410
3411
3412
3413
3414
3415
3416
3417
3418
3419
3420
3421
3422
3423
3424
3425
3426
3427
3428
3429
3430
3431
3432
3433
3434
3435
3436
3437
3438
3439
3440
3441
3442
3443
3444
3445
3446
3447
3448
3449
3450
3451
3452
3453
3454
3455
3456
3457
3458
3459
3460
3461
3462
3463
3464
3465
3466
3467
3468
3469
3470
3471
3472
3473
3474
3475
3476
3477
3478
3479
3480
3481
3482
3483
3484
3485
3486
3487
3488
3489
3490
3491
3492
3493
3494
3495
3496
3497
3498
3499
3500
3501
3502
3503
3504
3505
3506
3507
3508
3509
3510
3511
3512
3513
3514
3515
3516
3517
3518
3519
3520
3521
3522
3523
3524
3525
3526
3527
3528
3529
3530
3531
3532
3533
3534
3535
3536
3537
3538
3539
3540
3541
3542
3543
3544
3545
3546
3547
3548
3549
3550
3551
3552
3553
3554
3555
3556
3557
3558
3559
3560
3561
3562
3563
3564
3565
3566
3567
3568
3569
3570
3571
3572
3573
3574
3575
3576
3577
3578
3579
3580
3581
3582
3583
3584
3585
3586
3587
3588
3589
3590
3591
3592
3593
3594
3595
3596
3597
3598
3599
3600
3601
3602
3603
3604
3605
3606
3607
3608
3609
3610
3611
3612
3613
3614
3615
3616
3617
3618
3619
3620
3621
3622
3623
3624
3625
3626
3627
3628
3629
3630
3631
3632
3633
3634
3635
3636
3637
3638
3639
3640
3641
3642
3643
3644
3645
3646
3647
3648
3649
3650
3651
3652
3653
3654
3655
3656
3657
3658
3659
3660
3661
3662
3663
3664
3665
3666
3667
3668
3669
3670
3671
3672
3673
3674
3675
3676
3677
3678
3679
3680
3681
3682
3683
3684
3685
3686
3687
3688
3689
3690
3691
3692
3693
3694
3695
3696
3697
3698
3699
3700
3701
3702
3703
3704
3705
3706
3707
3708
3709
3710
3711
3712
3713
3714
3715
3716
3717
3718
3719
3720
3721
3722
3723
3724
3725
3726
3727
3728
3729
3730
3731
3732
3733
3734
3735
3736
3737
3738
3739
3740
3741
3742
3743
3744
3745
3746
3747
3748
3749
3750
3751
3752
3753
3754
3755
3756
3757
3758
3759
3760
3761
3762
3763
3764
3765
3766
3767
3768
3769
3770
3771
3772
3773
3774
3775
3776
3777
3778
3779
3780
3781
3782
3783
3784
3785
3786
3787
3788
3789
3790
3791
3792
3793
3794
3795
3796
3797
3798
3799
3800
3801
3802
3803
3804
3805
3806
3807
3808
3809
3810
3811
3812
3813
3814
3815
3816
3817
3818
3819
3820
3821
3822
3823
3824
3825
3826
3827
3828
3829
3830
3831
3832
3833
3834
3835
3836
3837
3838
3839
3840
3841
3842
3843
3844
3845
3846
3847
3848
3849
3850
3851
3852
3853
3854
3855
3856
3857
3858
3859
3860
3861
3862
3863
3864
3865
3866
3867
3868
3869
3870
3871
3872
3873
3874
3875
3876
3877
3878
3879
3880
3881
3882
3883
3884
3885
3886
3887
3888
3889
3890
3891
3892
3893
3894
3895
3896
3897
3898
3899
3900
3901
3902
3903
3904
3905
3906
3907
3908
3909
3910
3911
3912
3913
3914
3915
3916
3917
3918
3919
3920
3921
3922
3923
3924
3925
3926
3927
3928
3929
3930
3931
3932
3933
3934

LANE MEDICAL LIBRARY

To avoid fine, this book should be returned
on or before the date last stamped below.

--	--	--

